Index to Volume 52

Author Index

A

Abbas J (*see* Gupta V et al). 2010;52:1671-3
Abbate R (*see* Sticchi E et al). 2010;52:97-102
Abbott AL (*see* Nicolaides AN et al). 2010;52:1486-96
Abbruzzese TA (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Abouliatim I, Gouicem D, Kobeiter H, Majeski M, Becquemin J-P. Early type III endoleak with an Endurant endograft. 2010;52:1665-7
Abu Hamad G (*see* Chaer RA et al). 2010;52:569-75
Abu-Halimah S (*see* AbuRahma AF et al). 2010;52:1180-7
Abularrage CJ, Conrad MF, Hackney LA, Paruchuri V, Crawford RS, Kwolek CJ, LaMuraglia GM, Cambria RP. Long-term outcomes of diabetic patients undergoing endovascular infrainguinal interventions. 2010;52:314-22
Abularrage CJ, Crawford RS, Conrad MF, Lee H, Kwolek CJ, Brewster DC, Cambria RP, LaMuraglia GM. Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair. 2010;52:19-24
AbuRahma AF, Abu-Halimah S, Hass SM, Nanjundappa A, Stone PA, Mousa A, Lough E, Dean LS. Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis. 2010;52:1180-7
Acker MA (*see* Foley PJ et al). 2010;52:850-3
Acosta S (*see* Block TA et al). 2010;52:959-66
Adam DJ (*see* Bradbury AW et al). 2010;52:939-45 (Case rep.)
Adelman M (*see* Fonseca R et al). 2010;52:556-61
Adiseshiah M, Boardley D, Agu O, Raphael M. Twelve-year outcomes after endovascular aneurysm repair using earliest available endograft components from a single center. 2010;52:49-54
Agaimy A (*see* Bal A et al). 2010;52:456-9
Aggarwal R (*see* Willaert W et al). 2010;52:1700-5
Agko M (*see* Keilani ZM et al). 2010;52:1052-7
Agu O (*see* Adiseshiah M et al). 2010;52:49-54
Aguilera CM (*see* Lyon CA et al). 2010;52:1301-9
Aho P-S (*see* Arvela E et al). 2010;52:616-23
Aizaki M (*see* Kumakura H et al). 2010;52:110-7
Akcalı Y (*see* Kurtoglu M et al). 2010;52:1262-71
Akkersdijk GJM (*see* Jongkind V et al). 2010;52:1376-83
Al-Barjas HS (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Al-Omran M (*see* Altoijry A et al). 2010;52:1-2
Albäck A (*see* Arvela E et al). 2010;52:1218-25
Albadawi H (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Alegret JM, Calvo N, Ligero C, Palomares R, Millá L, Martín-Paredero V, Montero M, Joven J. Dilated aortic root is related to a global aortic dilating diathesis. 2010;52:867-72
Algra A (*see* Hartog AG et al). 2010;52:1062-71 (Presidential address)
Alimi YS (*see* Boufi M et al). 2010;52:1211-7
Alimi YS (*see* Hartung O et al). 2010;52:738-41
Altavilla D (*see* Bitto A et al). 2010;52:1015-21
Altoijry A, Mamdani M, Al-Omran M. Trends in abdominal aortic aneurysm repair in the era of endovascular technology in Ontario. 2010;52:1-2
Amann-Vesti B (*see* Willenberg T et al). 2010;52:664-8
Ambrosini V (*see* Stabile E et al). 2010;52:1505-10
Andersen C (*see* Causey MW et al). 2010;52:734-7
Andersen ND, Chopra A, Monahan TS, Malek JY, Jain M, Pradhan L, Ferran C, LoGerfo FW. Endothelial cells are susceptible to rapid siRNA transfection and gene silencing ex vivo. 2010;52:1608-15
Andreadou I (*see* Sigala F et al). 2010;52:704-13
Andrews JS (*see* Godshall CJ et al). 2010;52:775-81
Andrews JS (*see* Moore PS et al). 2010;52:118-26
Anil G, Taneja M. Revascularization of an occluded brachiocephalic vein using Outback-LTD re-entry catheter. 2010;52:1038-40
Annambhotla S (*see* Bechara CF et al). 2010;52:1682-96
Annex BH (*see* Powell RJ et al). 2010;52:1525-30
Aoyama T (*see* Ishii H et al). 2010;52:854-9
Aper T (*see* Bisdas T et al). 2010;52:323-30
Araki Y (*see* Kumakura H et al). 2010;52:110-7
Ariëns RAS (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Arvela E, Söderström M, Albäck A, Aho P-S, Venermo M, Lepäntalo M. Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia. 2010;52:616-23
Arvela E, Söderström M, Korhonen M, Halmesmäki K, Albäck A, Lepäntalo M, Venermo M, Biancari F. Finnvasc score and modified Prevent III score predict long-term outcome after infrainguinal surgical and endovascular revascularization for critical limb ischemia. 2010;52:1218-25
Ascher E (*see* Usoh F et al). 2010;52:394-9
Ashton HA (*see* Thompson A et al). 2010;52:55-61
Assi Z (*see* Comerota AJ et al). 2010;52:243-7
Asthana S (*see* Spark JI et al). 2010;52:632-6
Asymptomatic Carotid Stenosis and Risk of Stroke (ACSRS) Study Group (*see* Nicolaides AN et al). 2010;52:1486-96
Auquier P (*see* Boufi M et al). 2010;52:1211-7
Auricchio S (*see* Bonforte G et al). 2010;52:1551-6
Azghari A (*see* Hartung O et al). 2010;52:738-41

B

Bünger CM, Schareck W, Klar E, Kröger J-C. Laparoscopic treatment of renal artery entrapment. 2010;52:1357-61
Badger SA (*see* Jones C et al). 2010;52:298-302
Bai W (*see* Li T et al). 2010;52:1242-5
Bakaeen FG (*see* Schwill S et al). 2010;52:1034-7 (Case rep.)
Bal A, Schönleben F, Agaimy A, Gessner A, Lang W. *Listeria monocytogenes* as a rare cause of mycotic aortic aneurysm. 2010;52:456-9
Bannon PG (*see* Byrom MJ et al). 2010;52:176-95
Bao Q, Chen L. Thrombotic mesenteric ischemia due to aortic dissection. 2010;52:1674
Baptista-Silva JCC (*see* Erling N et al). 2010;52:1321-9
Baranowska-Bosiacka I (*see* Wiernicki I et al). 2010;52:691-6
Barnett C (*see* Mazari FAK et al). 2010;52:1564-71
Barshes NR, (*see* Nguyen LL et al). 2010;52:768-74
Barthelemy P (*see* Hartung O et al). 2010;52:738-41
Bate G (*see* Bradbury AW et al). 2010;52:939-45 (Case rep.)
Baumgartner I (*see* Willenberg T et al). 2010;52:664-8
Bavaria JE (*see* Brinster CJ et al). 2010;52:460-3
Beaver TM (*see* Lee WA et al). 2010;52:1459-63
Bechara CF, Annambhotla S, Lin PH. Access site management with vascular closure devices for percutaneous transarterial procedures. 2010;52:1682-96
Becher RD, Corriere MA, Edwards MS, Godshall CJ. Late erosion

Index to Volume 52

Author Index

A

Abbas J (*see* Gupta V et al). 2010;52:1671-3
Abbate R (*see* Sticchi E et al). 2010;52:97-102
Abbott AL (*see* Nicolaides AN et al). 2010;52:1486-96
Abbruzzese TA (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Abouliatim I, Gouicem D, Kobeiter H, Majeski M, Becquemin J-P. Early type III endoleak with an Endurant endograft. 2010;52:1665-7
Abu Hamad G (*see* Chaer RA et al). 2010;52:569-75
Abu-Halimah S (*see* AbuRahma AF et al). 2010;52:1180-7
Abularrage CJ, Conrad MF, Hackney LA, Paruchuri V, Crawford RS, Kwolek CJ, LaMuraglia GM, Cambria RP. Long-term outcomes of diabetic patients undergoing endovascular infrainguinal interventions. 2010;52:314-22
Abularrage CJ, Crawford RS, Conrad MF, Lee H, Kwolek CJ, Brewster DC, Cambria RP, LaMuraglia GM. Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair. 2010;52:19-24
AbuRahma AF, Abu-Halimah S, Hass SM, Nanjundappa A, Stone PA, Mousa A, Lough E, Dean LS. Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis. 2010;52:1180-7
Acker MA (*see* Foley PJ et al). 2010;52:850-3
Acosta S (*see* Block TA et al). 2010;52:959-66
Adam DJ (*see* Bradbury AW et al). 2010;52:939-45 (Case rep.)
Adelman M (*see* Fonseca R et al). 2010;52:556-61
Adiseshiah M, Boardley D, Agu O, Raphael M. Twelve-year outcomes after endovascular aneurysm repair using earliest available endograft components from a single center. 2010;52:49-54
Agaimy A (*see* Bal A et al). 2010;52:456-9
Aggarwal R (*see* Willaert W et al). 2010;52:1700-5
Agko M (*see* Keilani ZM et al). 2010;52:1052-7
Agu O (*see* Adiseshiah M et al). 2010;52:49-54
Aguilera CM (*see* Lyon CA et al). 2010;52:1301-9
Aho P-S (*see* Arvela E et al). 2010;52:616-23
Aizaki M (*see* Kumakura H et al). 2010;52:110-7
Akcalı Y (*see* Kurtoglu M et al). 2010;52:1262-71
Akkersdijk GJM (*see* Jongkind V et al). 2010;52:1376-83
Al-Barjas HS (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Al-Omran M (*see* Altoijry A et al). 2010;52:1-2
Albäck A (*see* Arvela E et al). 2010;52:1218-25
Albadawi H (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Alegret JM, Calvo N, Ligero C, Palomares R, Millá L, Martín-Paredero V, Montero M, Joven J. Dilated aortic root is related to a global aortic dilating diathesis. 2010;52:867-72
Algra A (*see* Hartog AG et al). 2010;52:1062-71 (Presidential address)
Alimi YS (*see* Boufi M et al). 2010;52:1211-7
Alimi YS (*see* Hartung O et al). 2010;52:738-41
Altavilla D (*see* Bitto A et al). 2010;52:1015-21
Altoijry A, Mamdani M, Al-Omran M. Trends in abdominal aortic aneurysm repair in the era of endovascular technology in Ontario. 2010;52:1-2
Amann-Vesti B (*see* Willenberg T et al). 2010;52:664-8
Ambrosini V (*see* Stabile E et al). 2010;52:1505-10
Andersen C (*see* Causey MW et al). 2010;52:734-7
Andersen ND, Chopra A, Monahan TS, Malek JY, Jain M, Pradhan L, Ferran C, LoGerfo FW. Endothelial cells are susceptible to rapid siRNA transfection and gene silencing ex vivo. 2010;52:1608-15
Andreadou I (*see* Sigala F et al). 2010;52:704-13
Andrews JS (*see* Godshall CJ et al). 2010;52:775-81
Andrews JS (*see* Moore PS et al). 2010;52:118-26
Anil G, Taneja M. Revascularization of an occluded brachiocephalic vein using Outback-LTD re-entry catheter. 2010;52:1038-40
Annambhotla S (*see* Bechara CF et al). 2010;52:1682-96
Annex BH (*see* Powell RJ et al). 2010;52:1525-30
Aoyama T (*see* Ishii H et al). 2010;52:854-9
Aper T (*see* Bisdas T et al). 2010;52:323-30
Araki Y (*see* Kumakura H et al). 2010;52:110-7
Ariëns RAS (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Arvela E, Söderström M, Albäck A, Aho P-S, Venermo M, Lepäntalo M. Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia. 2010;52:616-23
Arvela E, Söderström M, Korhonen M, Halmesmäki K, Albäck A, Lepäntalo M, Venermo M, Biancari F. Finnvasc score and modified Prevent III score predict long-term outcome after infrainguinal surgical and endovascular revascularization for critical limb ischemia. 2010;52:1218-25
Ascher E (*see* Usoh F et al). 2010;52:394-9
Ashton HA (*see* Thompson A et al). 2010;52:55-61
Assi Z (*see* Comerota AJ et al). 2010;52:243-7
Asthana S (*see* Spark JI et al). 2010;52:632-6
Asymptomatic Carotid Stenosis and Risk of Stroke (ACSRS) Study Group (*see* Nicolaides AN et al). 2010;52:1486-96
Auquier P (*see* Boufi M et al). 2010;52:1211-7
Auricchio S (*see* Bonforte G et al). 2010;52:1551-6
Azghari A (*see* Hartung O et al). 2010;52:738-41

B

Bünger CM, Schareck W, Klar E, Kröger J-C. Laparoscopic treatment of renal artery entrapment. 2010;52:1357-61
Badger SA (*see* Jones C et al). 2010;52:298-302
Bai W (*see* Li T et al). 2010;52:1242-5
Bakaeen FG (*see* Schwill S et al). 2010;52:1034-7 (Case rep.)
Bal A, Schönleben F, Agaimy A, Gessner A, Lang W. *Listeria monocytogenes* as a rare cause of mycotic aortic aneurysm. 2010;52:456-9
Bannon PG (*see* Byrom MJ et al). 2010;52:176-95
Bao Q, Chen L. Thrombotic mesenteric ischemia due to aortic dissection. 2010;52:1674
Baptista-Silva JCC (*see* Erling N et al). 2010;52:1321-9
Baranowska-Bosiacka I (*see* Wiernicki I et al). 2010;52:691-6
Barnett C (*see* Mazari FAK et al). 2010;52:1564-71
Barshes NR, (*see* Nguyen LL et al). 2010;52:768-74
Barthelemy P (*see* Hartung O et al). 2010;52:738-41
Bate G (*see* Bradbury AW et al). 2010;52:939-45 (Case rep.)
Baumgartner I (*see* Willenberg T et al). 2010;52:664-8
Bavaria JE (*see* Brinster CJ et al). 2010;52:460-3
Beaver TM (*see* Lee WA et al). 2010;52:1459-63
Bechara CF, Annambhotla S, Lin PH. Access site management with vascular closure devices for percutaneous transarterial procedures. 2010;52:1682-96
Becher RD, Corriere MA, Edwards MS, Godshall CJ. Late erosion

of a prophylactic Celect IVC filter into the aorta, right renal artery, and duodenal wall. 2010;52:1041-4
Becquemin J-P (*see* Abouliatim I et al). 2010;52:1665-7
Becquemin J-P (*see* Houbballah R et al). 2010;52:878-83
Beeman BR, Murtha K, Doerr K, McAfee-Bennett S, Dougherty MJ, Calligaro KD. Duplex ultrasound factors predicting persistent type II endoleak and increasing AAA sac diameter after EVAR. 2010;52:1147-52
Belch JJF, Dormandy J and the, CASPAR Writing Committee. Results of the randomized, placebo-controlled clopidogrel and acetylsalicylic acid in bypass surgery for peripheral arterial disease (CASPAR) trial. 2010;52:825-33
Beller CJ, Gebhard MM, Karck M, Labrosse MR. Usefulness and limitations of computational models in aortic disease risk stratification. 2010;52:1572-9 (Basic research studies)
Bena JF (*see* Kashyap VS et al). 2010;52:369-74
Berard X, Corpataux J-M, Taoufiq H, Sassoust G, Brizzi V, Midy D. Don't trust a vein graft to treat carotid aneurysm in patients with Behçet disease. 2010;52:471-4
Berguer R (*see* Criado E et al). 2010;52:406-11
Berne JD (*see* Keilani ZM et al). 2010;52:1052-7
Bertges DJ (*see* De Martino RR et al). 2010;52:5-12
Bertges DJ, Goodney PP, Zhao Y, Schanzer A, Nolan BW, Likosky DS, Eldrup-Jorgensen J, Cronenwett JL, Vascular Study Group of New England. The Vascular Study Group of New England Cardiac Risk Index (VSG-CRI) predicts cardiac complications more accurately than the Revised Cardiac Risk Index in vascular surgery patients. 2010;52:674-83
Bertoglio L (*see* Civilini E et al). 2010;52:758-9
Beyer J (*see* Schwarz T et al). 2010;52:1246-50
Biamino G (*see* Stabile E et al). 2010;52:1505-10
Biancari F (*see* Arvela E et al). 2010;52:1218-25
Bicknell C (*see* Willaert W et al). 2010;52:1700-5
Bien S (*see* Wiegand S et al). 2010;52:742-5
Bisdas S (*see* Bisdas T et al). 2010;52:323-30
Bisdas T, Bredt M, Pichlmaier M, Aper T, Wilhelmi M, Bisdas S, Haverich A, Teebken OE. Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections. 2010;52:323-30
Bishop PD (*see* Kashyap VS et al). 2010;52:369-74
Bismuth J (*see* Davies MG et al). 2010;52:331-9
Bismuth J, Donovan MA, O'Malley MK, El Sayed HF, Naoum JJ, Peden EK, Davies MG, Lumsden AB. Incorporating simulation in vascular surgery education. 2010;52:1072-80
Bitto A, De Caridi G, Polito F, Calò M, Irrera N, Altavilla D, Spinelli F, Squadrito F. Evidence for markers of hypoxia and apoptosis in explanted human carotid atherosclerotic plaques. 2010;52:1015-21
Björck M (*see* Block TA et al). 2010;52:959-66
Bjerk AA (*see* Stone DH et al). 2010;52:127-31
Bjurholm A (*see* Mani K et al). 2010;52:1653-7 (Case rep.)
Black JH (*see* Guzzo JL et al). 2010;52:658-63
Black S (*see* McQuade K et al). 2010;52:584-91
Blaschuk OW (*see* Lyon CA et al). 2010;52:1301-9
Blecha MJ (*see* Willson TD et al). 2010;52:219-21
Blest N (*see* Spark JI et al). 2010;52:632-6
Blevins SM (*see* Gardner AW et al). 2010;52:1204-10
Block TA, Acosta S, Björck M. Endovascular and open surgery for acute occlusion of the superior mesenteric artery. 2010;52:959-66
Boardley D (*see* Adiseshiah M et al). 2010;52:49-54
Bobadilla JL, Hoch JR, Leverson GE, Tefera G. The effect of warfarin therapy on endoleak development after endovascular aneurysm repair (EVAR) of the abdominal aorta. 2010;52:267-71
Bode C (*see* Duerschmied D et al). 2010;52:1531-6
Bogey WM (*see* Durham CA et al). 2010;52:884-90
Boiteau G (*see* Grisafi JL et al). 2010;52:1346-9
Boldt O (*see* Donas KP et al). 2010;52:356-61
Bolia A (*see* Markose G et al). 2010;52:1410-6
Bond D (*see* Griffin MB et al). 2010;52:69-76
Bonforte G, Rossi E, Auricchio S, Pogliani D, Mangano S, Mandolfo S, Galli F, Genovesi S. The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease. 2010;52:1551-6
Bosman W-MPF, van der Steenhoven TJ, Suárez DR, Valstar ER, de Vries AC, Brom HLF, Jacobs MJ, Hamming JF. The effect of injectable biocompatible elastomer (PDMS) on the strength of the proximal fixation of endovascular aneurysm repair grafts: An in vitro study. 2010;52:152-8
Bosson J-L (*see* Galanaud J-P et al). 2010;52:932-8
Boufi M (*see* Hartung O et al). 2010;52:738-41
Boufi M, Dona B, Orsini B, Auquier P, Hartung O, Alimi YS. A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions. 2010;52:1211-7
Bounameaux H (*see* Christenson JT et al). 2010;52:1234-41
Bowen BP (*see* Lopes LB et al). 2010;52:1596-607
Boyd CS (*see* Jones C et al). 2010;52:298-302
Boyne NS (*see* Kruger AJ et al). 2010;52:1518-24
Bradbury AW, Bate G, Pang K, Darvall KA, Adam DJ. Ultrasound-guided foam sclerotherapy is a safe and clinically effective treatment for superficial venous reflux. 2010;52:939-45 (Case rep.)
Brawley J (*see* Lee GS et al). 2010;52:1058-60
Bredt M (*see* Bisdas T et al). 2010;52:323-30
Brewster DC (*see* Abularrage CJ et al). 2010;52:19-24
Brewster LP, Greisler HP. There may be nothing new under the sun, but this is ridiculous. 2010;52:1697-8 (Surgical ethics challenges)
Brinster CJ, Szeto WY, Bavaria JE, Woo EY, Fairman RM, Jackson BM. Endovascular repair of extent I thoracoabdominal aneurysms with landing zone extension into the aortic arch and mesenteric portion of the abdominal aorta. 2010;52:460-3
Brizzi V (*see* Berard X et al). 2010;52:471-4
Brom HLF (*see* Bosman W-MPF et al). 2010;52:152-8
Brophy CM (*see* Lopes LB et al). 2010;52:1596-607
Brown B (*see* Mazari FAK et al). 2010;52:1564-71
Brown LC (*see* Sweeting MJ et al). 2010;52:1-4
Buckley SD (*see* Buckley CJ et al). 2010;52:237-41 (Special communications)
Buckley CJ, Rutherford RB, Diethrich EB, Buckley SD. Inherent problems with randomized clinical trials with observational/no treatment arms. 2010;52:237-41 (Special communications)
Burchardt ER (*see* Comerota AJ et al). 2010;52:723-9
Burdess A, Nimmo AF, Campbell N, Harding SA, Garden OJ, Dawson AR, Newby DE. Perioperative platelet and monocyte activation in patients with critical limb ischemia. 2010;52:697-703
Burkhardt CGE, Cox M, Clouse WD, Porras C, Gifford CSM, Williams K, Propper CBW, Rasmussen TE. Outcomes of selective tibial artery repair following combat-related extremity injury. 2010;52:91-6
Burris DG (*see* Rich NM et al). 2010;52:508-11
Burris DG (*see* Welling DR et al). 2010;52:790-3
Buschmann L (*see* Schwarz T et al). 2010;52:1246-50

Bush HL (*see* Goldstein LJ et al). 2010;52:1140-6
Bush RL. Invited commentary. 2010;52:1140-6
Bustami R (*see* Sales CM et al). 2010;52:1251-4
Byrom MJ, Bannon PG, White GH, Ng MKC. Animal models for the assessment of novel vascular conduits. 2010;52:176-95

C

Cabral GM (*see* da Gama AD et al). 2010;52:1668-70
Calò M (*see* Bitto A et al). 2010;52:1015-21
Calderin J (*see* De Marco Garcia LP et al). 2010;52:139-44
Caldito G (*see* West CA et al). 2010;52:1164-72
Calligaro KD (*see* Beeman BR et al). 2010;52:1147-52
Calvo N (*see* Alegret JM et al). 2010;52:867-72
Cambria R (*see* De Martino RR et al). 2010;52:5-12
Cambria RP (*see* Abularrage CJ et al). 2010;52:314-22
Cambria RP (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Campagna C (*see* Yu P et al). 2010;52:444-52
Campbell N (*see* Burdess A et al). 2010;52:697-703
CAPTURE 2 Investigators and Executive Committee (*see* Matsumura JS et al). 2010;52:576-83
Carpenter JP (*see* Patel MS et al). 2010;52:1135-9
Carradice D (*see* Mazari FAK et al). 2010;52:77-84
Carroll M (*see* Schmieder GC et al). 2010;52:362-8
Carson S (*see* West CA et al). 2010;52:1164-72
Cartañá R (*see* Pereda D et al). 2010;52:1362-4
Carter V (*see* Lawrence PF et al). 2010;52:388-93
Caselli G (*see* Gabrielli R et al). 2010;52:464-6
Casey PJ (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
CASPAR Writing Committee (*see* Belch JJF et al). 2010;52:825-33
Castaldo J (*see* Li Y et al). 2010;52:1716-22
Causey MW, Oguntoye MO, Miller S, Andersen C, Singh N. Limb salvage after delayed diagnosis for blunt traumatic infrapopliteal occlusion. 2010;52:734-7
Causey MW, Quan R, Hamawy A, Singh N. Superficialization of arteriovenous fistulae employing minimally invasive liposuction. 2010;52:1397-400
Cavatassi W (*see* Endean ED et al). 2010;52:720-2 (Case rep.)
Cayne N (*see* Fonseca R et al). 2010;52:556-61
Chaer RA (*see* Cho J-S et al). 2010;52:1127-34
Chaer RA (*see* Fernandez N et al). 2010;52:834-42
Chaer RA (*see* Wagner JK et al). 2010;52:216-8
Chaer RA, Shen J, Rao A, Cho JS, Abu Hamad G, Makaroun MS. Cerebral reserve is decreased in elderly patients with carotid stenosis. 2010;52:569-75
Chahim M (*see* Uhi J-F et al). 2010;52:714-9
Chaikof EL (*see* Chung J et al). 2010;52:1435-41
Chaikof EL (*see* Fillinger MF et al). 2010;52:1022-33
Chandra A (*see* Lawrence PF et al). 2010;52:388-93
Chang C-J (*see* Hsiao J-F et al). 2010;52:669-73
Chang CK (*see* De Martino RR et al). 2010;52:5-12
Chang CK (*see* Stone DH et al). 2010;52:127-31
Chang K (*see* Guzzo JL et al). 2010;52:658-63
Chang L (*see* Hamblin M et al). 2010;52:984-93
Chao C-C (*see* Lin M-S et al). 2010;52:308-13
Chao C-L (*see* Lin M-S et al). 2010;52:308-13
Chapot R (*see* Wiegand S et al). 2010;52:742-5
Chappell L (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Charalabidis P (*see* Spark JI et al). 2010;52:632-6
Chaturvedi S (*see* Matsumura JS et al). 2010;52:576-83
Chaux A (*see* Song L et al). 2010;52:994-1002
Cheever D (*see* Riles TS et al). 2010;52:914-9
Chen B (*see* Dong Z et al). 2010;52:1450-8
Chen L (*see* Bao Q et al). 2010;52:1674
Chen M-F (*see* Lin M-S et al). 2010;52:308-13
Chen Y-H (*see* Lin M-S et al). 2010;52:308-13
Chen YE (*see* Hamblin M et al). 2010;52:984-93
Chen Y-S, Lu M-J, Huang H-S, Ma M-C. Mechanosensitive transient receptor potential vanilloid type 1 channels contribute to vascular remodeling of rat fistula veins. 2010;52:1310-20
Cheng J (*see* Matsumura JS et al). 2010;52:576-83
Cheshire N (*see* Willaert W et al). 2010;52:1700-5
Chetter IC (*see* Mazari FAK et al). 2010;52:1564-71
Chiappa R (*see* Gabrielli R et al). 2010;52:464-6
Chiesa R (*see* Civilini E et al). 2010;52:758-9
Chiesa R (*see* Marone EM et al). 2010;52:1511-7
Ching YA (*see* Giles KA et al). 2010;52:1471-7
Cho J (*see* Crisostomo PR et al). 2010;52:1272-7
Cho JS (*see* Chaer RA et al). 2010;52:569-75
Cho J-S, Park T, Kim JY, Chaer RA, Rhee RY, Makaroun MS. Prior endovascular abdominal aortic aneurysm repair provides no survival benefits when the aneurysm ruptures. 2010;52:1127-34
Choke E, Cockerill GW, Dawson J, Howe F, Wilson WRW, Loftus IM, Thompson MM. Vascular endothelial growth factor enhances angiotensin II-induced aneurysm formation in apolipoprotein E-deficient mice. 2010;52:159-66
Chopra A (*see* Andersen ND et al). 2010;52:1608-15
Chou H-H (*see* Hsiao J-F et al). 2010;52:669-73
Chou M-M (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Christenson JT, Gueddi S, Gemayel G, Bounameaux H. Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up. 2010;52:1234-41
Chung J, Corriere MA, Milner R, Kasirajan K, Salam A, Dodson TF, Chaikof EL, Veeraswamy RK. Midterm results of adjunctive neck therapies performed during elective infrarenal aortic aneurysm repair. 2010;52:1435-41
Chung J, Corriere MA, Veeraswamy RK, Kasirajan K, Milner R, Dodson TF, Salam AA, Chaikof EL. Risk factors for late mortality after endovascular repair of the thoracic aorta. 2010;52:549-55
Chuter TAM (*see* Monahan TS et al). 2010;52:303-7
Civilini E, Melissano G, Bertoglio L, Chiesa R. Collateral pathways visualization of the innominate vein. 2010;52:758-9
Clagett GP (*see* Lang JL et al). 2010;52:290-7
Clagett GP (*see* Timaran CH et al). 2010;52:1188-95
Clouse WD (*see* Burkhardt CGE et al). 2010;52:91-6
Cochery T (*see* Mahe G et al). 2010;52:62-8
Cockerill GW (*see* Choke E et al). 2010;52:159-66
Collins A (*see* Gordon P et al). 2010;52:467-70
Comerota AJ (*see* Gale SS et al). 2010;52:645-50
Comerota AJ (*see* Thakur S et al). 2010;52:208-11
Comerota AJ, Grewal NK, Thakur S, Assi Z. Endovenectomy of the common femoral vein and intraoperative iliac vein recanalization for chronic iliofemoral venous occlusion. 2010;52:243-7
Comerota AJ, Link A, Douville J, Burchardt ER. Upper extremity ischemia treated with tissue repair cells from adult bone marrow. 2010;52:723-9
Conforti M (*see* Rispoli P et al). 2010;52:212-5

Conrad MF (*see* Abularrage CJ et al). 2010;52:314-22
Constantinescu I (*see* Stoica AL et al). 2010;52:103-9
Cooper JA (*see* Thompson A et al). 2010;52:55-61
Coppi G (*see* Marone EM et al). 2010;52:1511-7
Corpataux J-M (*see* Berard X et al). 2010;52:471-4
Corriere MA (*see* Becher RD et al). 2010;52:1041-4
Corriere MA (*see* Chung J et al). 2010;52:1435-41
Corriere MA (*see* Moore PS et al). 2010;52:118-26
Coscas R, Greenberg RK, Mastracci TM, Eagleton M, Kang WC, Morales C, Hernandez AV. Associated factors, timing, and technical aspects of late failure following open surgical aneurysm repairs. 2010;52:272-81
Coselli JS (*see* Schwill S et al). 2010;52:1034-7 (Case rep.)
Costanzo A (*see* Gabrielli R et al). 2010;52:464-6
Cox M (*see* Burkhardt CGE et al). 2010;52:91-6
Craven TE (*see* Fleming SH et al). 2010;52:953-8
Craven TE (*see* Moore PS et al). 2010;52:118-26
Crawford RS (*see* Abularrage CJ et al). 2010;52:314-22
Criado E, Berguer R, Greenfield L. The spectrum of arterial compression at the thoracic outlet. 2010;52:406-11
Criqui MH (*see* Singh N et al). 2010;52:624-31
Criqui MH, Ho LA, Denenberg JO, Ridker PM, Wassel CL, McDermott MM. Biomarkers in peripheral arterial disease patients and near- and longer-term mortality. 2010;52:85-90
Crisostomo PR, Cho J, Feliciano B, Klein J, Jones D, Dalsing MC. Period frequency of iliofemoral venous occlusive disease by Doppler ultrasound and corresponding treatment in a tertiary care facility. 2010;52:1272-7
Croft BA (*see* Davies MG et al). 2010;52:1478-85
Cronenwett JL (*see* Bertges DJ et al). 2010;52:674-83
Cronenwett JL (*see* De Martino RR et al). 2010;52:5-12
Cronenwett JL (*see* Goodney PP et al). 2010;52:897-905
Cruz JWMC (*see* Erling N et al). 2010;52:1321-9

D

da Gama AD, Cabral GM. Carotid body tumor presenting with carotid sinus syndrome. 2010;52:1668-70
Daamen WF (*see* Geutjes PJ et al). 2010;52:1330-8
Dalsing MC (*see* Crisostomo PR et al). 2010;52:1272-7
Darling RC (*see* Mehta M et al). 2010;52:1442-9
Darling RC (*see* Rahimi SA et al). 2010;52:891-6
Darvall KA (*see* Bradbury AW et al). 2010;52:939-45 (Case rep.)
Darzi A (*see* Willaert W et al). 2010;52:1700-5
Davidovic L (*see* Dragas M et al). 2010;52:1658-61
Davies MG (*see* Bismuth J et al). 2010;52:1072-80
Davies MG, Bismuth J, Saad WE, Naoum JJ, Peden EK, Lumsden AB. Outcomes of interventions for recurrent disease after endoluminal intervention for superficial femoral artery disease. 2010;52:331-9
Davies MG, Younes HK, Harris PW, Masud F, Croft BA, Reardon MJ, Lumsden AB. Outcomes before and after initiation of an acute aortic treatment center. 2010;52:1478-85
Davila-Santini LR (*see* De Marco Garcia LP et al). 2010;52:139-44
Davis KA (*see* Jonker FHW et al). 2010;52:39-44
Davis RP (*see* Fleming SH et al). 2010;52:953-8
Davis RP (*see* Moore PS et al). 2010;52:118-26
Dawson AR (*see* Burdess A et al). 2010;52:697-703
Dawson J (*see* Choke E et al). 2010;52:159-66
Dayal R (*see* Giacovelli JK et al). 2010;52:
de Borst GJ (*see* Hartog AG et al). 2010;52:1062-71 (Presidential address)
De Bray J-M (*see* Mahe G et al). 2010;52:62-8
De Caridi G (*see* Bitto A et al). 2010;52:1015-21
de Kleijn D (*see* Derksen WJM et al). 2010;52:592-9
de Vries AC (*see* Bosman W-MPF et al). 2010;52:152-8
de Vries J-PPM (*see* Derksen WJM et al). 2010;52:592-9
de Vries N (*see* Geutjes PJ et al). 2010;52:1330-8
de Wit GA (*see* Sprengers RW et al). 2010;52:843-9
De Marco Garcia LP, Davila-Santini LR, Feng Q, Calderin J, Krishnasastry KV, Panetta TF. Primary balloon angioplasty plus balloon angioplasty maturation to upgrade small-caliber veins (<3 mm) for arteriovenous fistulas. 2010;52:139-44
De Martino RR, Nolan BW, Goodney PP, Chang CK, Schanzer A, Cambria R, Bertges DJ, Cronenwett JL, Vascular Study Group of Northern New England. Outcomes of symptomatic abdominal aortic aneurysm repair. 2010;52:5-12
Dean LS (*see* AbuRahma AF et al). 2010;52:1180-7
deFreitas DJ (*see* Stoner MC et al). 2010;52:223-31 (Review article)
DeLoughery TG (*see* Liem TK et al). 2010;52:651-7
Demos J (*see* Guzzo JL et al). 2010;52:658-63
Denenberg JO (*see* Criqui MH et al). 2010;52:85-90
Denollet J (*see* Smolderen KG et al). 2010;52:637-44
Deonanan JK (*see* Fleming SH et al). 2010;52:953-8
Derksen WJM, de Vries J-PPM, Vink A, Velema E, Vos J-A, de Kleijn D, Moll FL, Pasterkamp G. Histologic atherosclerotic plaque characteristics are associated with restenosis rates after endarterectomy of the common and superficial femoral arteries. 2010;52:592-9
DeRose G (*see* Forbes TL et al). 2010;52:45-8
DeRubertis B (*see* Lawrence PF et al). 2010;52:388-93
Derubertis BG (*see* Tulloch AW et al). 2010;52:1283-9
Detschelt E (*see* Grisafi JL et al). 2010;52:1346-9
Diehm N (*see* Willenberg T et al). 2010;52:664-8
Diethrich EB (*see* Buckley CJ et al). 2010;52:237-41 (Special communications)
Dillavou ED (*see* Spangler EL et al). 2010;52:
Dimick JB (*see* Osborne NH et al). 2010;52:400-5
Ding X (*see* Jiang J et al). 2010;52:1278-82
Ding P-X, Li Y-D, Han X-W, Wu G. Agitation thrombolysis for fresh iatrogenic IVC thrombosis in patients with Budd-Chiari syndrome. 2010;52:782-4
Dodson TF (*see* Chung J et al). 2010;52:1435-41
Doerr K (*see* Beeman BR et al). 2010;52:1147-52
Dona B (*see* Boufi M et al). 2010;52:1211-7
Donas KP, Torsello G, Schwindt A, Schönefeld E, Boldt O, Pitoulias GA. Below knee bare nitinol stent placement in high-risk patients with critical limb ischemia is still durable after 24 months of follow-up. 2010;52:356-61
Donayre CE (*see* Khoynezhad A et al). 2010;52:562-8
Dong Z, Fu W, Wang Y, Wang C, Yan Z, Guo D, Xu X, Chen B. Stent graft-induced new entry after endovascular repair for Stanford type B aortic dissection. 2010;52:1450-8
Donovan MA (*see* Bismuth J et al). 2010;52:1072-80
Doré CJ (*see* Griffin MB et al). 2010;52:69-76
Doré CJ (*see* Nicolaides AN et al). 2010;52:1486-96
Dormandy J and the (*see* Belch JJF et al). 2010;52:825-33
Doucet L (*see* West CA et al). 2010;52:1164-72
Dougherty MJ (*see* Beeman BR et al). 2010;52:1147-52
Douville J (*see* Comerota AJ et al). 2010;52:723-9
Dragas M, Davidovic L, Pejkic S, Ilic N, Koncar I, Markovic M. Aorto-left renal vein fistula is a rare complication of abdominal aortic aneurysm with unique clinical presentation. 2010;52:1658-61
Duerschmied D, Maletzki P, Freund G, Olschewski M, Bode C, Hehrlein C. Success of arterial revascularization determined by

contrast ultrasound muscle perfusion imaging. 2010;52: 1531-6
Durham CA, McNally MM, Parker FM, Bogey WM, Powell CS, Goettler CE, Rotondo MF, Stoner MC. A contemporary rural trauma center experience in blunt traumatic aortic injury. 2010;52:884-90
Durham CA, Mohr MC, Parker FM, Bogey WM, Powell CS, Stoner MC. The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization. 2010;52:600-7
Dutson EP (*see* Tulloch AW et al). 2010;52:1283-9

E

Eagleton M (*see* Coscas R et al). 2010;52:272-81
Eagleton MJ. Invited Commentary. 2010;52:867-72
Edwards MS (*see* Becher RD et al). 2010;52:1041-4
Edwards MS (*see* Godshall CJ et al). 2010;52:775-81
Edwards MS (*see* Moore PS et al). 2010;52:118-26
Egashira K (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Egorova N (*see* Giacovelli JK et al). 2010;52:
Egorova NN (*see* Kent KC et al). 2010;52:539-48
Egorova NN (*see* Vouyouka AG et al). 2010;52:1196-203
Ekstrand J, Razuvaev A, Folkersen L, Roy J, Hedin U. Tissue factor pathway inhibitor-2 is induced by fluid shear stress in vascular smooth muscle cells and affects cell proliferation and survival. 2010;52:167-75 (Basic research studies)
El Sayed HF (*see* Bismuth J et al). 2010;52:1072-80
Eldrup-Jorgensen J (*see* Bertges DJ et al). 2010;52:674-83
Eldrup-Jorgensen J (*see* Goodney PP et al). 2010;52:897-905
Endean ED, Cavatassi W, Hansler J, Sorial E. Deglutition syncope: A manifestation of vagal hyperactivity following carotid endarterectomy. 2010;52:720-2 (Case rep.)
Endo M (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Eriksson L-G (*see* Mani K et al). 2010;52:1653-7 (Case rep.)
Erling N, Nakagawa NK, Cruz JWMC, Zanoni FL, Baptista-Silva JCC, Sannomiya P, Poli-de-Figueiredo LF. Microcirculatory effects of local and remote ischemic preconditioning in supraceliac aortic clamping. 2010;52:1321-9
Eton D, Yu H. Enhanced cell therapy strategy to treat chronic limb-threatening ischemia. 2010;52:199-204
European Virtual Reality Endovascular Research Team (EVEResT) (*see* Willaert W et al). 2010;52:1700-5
Exercise Therapy in Peripheral Arterial Disease (EXITPAD) study group (*see* Nicolaï SPA et al). 2010;52:348-55
EXITPAD study group (*see* Nicolaï SPA et al). 2010;52:1226-33

F

Fabricius M (*see* Thompson A et al). 2010;52:55-61
Fairman RM (*see* Brinster CJ et al). 2010;52:460-3
Fairman RM (*see* Foley PJ et al). 2010;52:850-3
Faraj KA (*see* Geutjes PJ et al). 2010;52:1330-8
Faries PL (*see* Han DK et al). 2010;52:746-50
Faries PL (*see* Vouyouka AG et al). 2010;52:1196-203
Fatini C (*see* Sticchi E et al). 2010;52:97-102
Feliciano B (*see* Crisostomo PR et al). 2010;52:1272-7
Feng Q (*see* De Marco Garcia LP et al). 2010;52:139-44
Fernandez JD (*see* Perera AD et al). 2010;52:1339-42 (Case rep.)
Fernandez N, McEnaney R, Marone LK, Rhee RY, Leers S, Makaroun M, Chaer RA. Predictors of failure and success of tibial interventions for critical limb ischemia. 2010;52:834-42
Ferran C (*see* Andersen ND et al). 2010;52:1608-15
Ferrigni RG (*see* Nakamura LY et al). 2010;52:453-5 (Case rep.)
Ferrucci L (*see* Singh N et al). 2010;52:624-31
FFBI Clinical Practice Workgroup (*see* Sales CM et al). 2010;52: 1699
Filis K (*see* Sigala F et al). 2010;52:704-13
Filizcan U (*see* Kurtoglu M et al). 2010;52:1262-71
Fillinger MF, Greenberg RK, McKinsey JF, Chaikof EL, Society for Vascular Surgery Ad Hoc Committee on TEVAR Reporting Standards. Reporting standards for thoracic endovascular aortic repair (TEVAR). 2010;52:1022-33
Fishman EW (*see* Han DK et al). 2010;52:746-50
Fleming SH (*see* Godshall CJ et al). 2010;52:775-81
Fleming SH, Davis RP, Craven TE, Deonanan JK, Godshall CJ, Hansen KJ. Accuracy of duplex sonography scans after renal artery stenting. 2010;52:953-8
Flynn CR (*see* Lopes LB et al). 2010;52:1596-607
Foley PJ, Morris RJ, Woo EY, Acker MA, Wang GJ, Fairman RM, Jackson BM. Limb ischemia during femoral cannulation for cardiopulmonary support. 2010;52:850-3
Folkersen L (*see* Ekstrand J et al). 2010;52:167-75 (Basic research studies)
Fonseca R, Rockman C, Pitti A, Cayne N, Maldonado TS, Lamparello PJ, Riles T, Adelman M. Intermediate-term EVAR outcomes in octogenarians. 2010;52:556-61
Forbes TL (*see* Tong MZ-Y et al). 2010;52:31-8
Forbes TL. Commentary. 2010;52:499-507
Forbes TL, Harris JR, Lawlor DK, DeRose G. Aortic dilatation after endovascular repair of blunt traumatic thoracic aortic injuries. 2010;52:45-8
Forbes TL, Ricco J-B. Commentary. 2010;52:1087-93
Forbes TL, Ricco J-B. Editors' Commentary. 2010;52:1716-22
Forster AJ (*see* van Walraven C et al). 2010;52:282-9
Foster WJ (*see* Kruger AJ et al). 2010;52:1518-24
Fowl RJ (*see* Nakamura LY et al). 2010;52:453-5 (Case rep.)
Freischlag JA (*see* Guzzo JL et al). 2010;52:658-63
Freund G (*see* Duerschmied D et al). 2010;52:1531-6
Fu W (*see* Dong Z et al). 2010;52:1450-8
Fukumoto H (*see* Yoshino M et al). 2010;52:1401-4
Fukumoto Y (*see* Tanaka H et al). 2010;52:479

G

Gable D (*see* McQuade K et al). 2010;52:584-91
Gabrielli R, Rosati MS, Costanzo A, Chiappa R, Siani A, Caselli G. Primary tibial vein aneurysm with recurrent pulmonary emboli. 2010;52:464-6
Galanaud J-P, Sevestre M-A, Genty C, Laroche J-P, Zizka V, Quéré I, Bosson J-L, OPTIMEV SFMV investigators. Comparison of the clinical history of symptomatic isolated muscular calf vein thrombosis versus deep calf vein thrombosis. 2010;52: 932-8
Gale SS, Lee JN, Walsh ME, Wojnarowski DL, Comerota AJ. A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein. 2010;52:645-50
Gallagher KA (*see* Goldstein LJ et al). 2010;52:1140-6
Galli F (*see* Bonforte G et al). 2010;52:1551-6
Gao X (*see* Matsumura JS et al). 2010;52:576-83
Garden OJ (*see* Burdess A et al). 2010;52:697-703
Gardiner KR (*see* Gordon P et al). 2010;52:467-70
Gardner AW, Parker DE, Montgomery PS, Khurana A, Ritti-Dias RM, Blevins SM. Gender differences in daily ambulatory activ-

ity patterns in patients with intermittent claudication. 2010;52: 1204-10
Garrett HE (*see* Perera AD et al). 2010;52:1339-42 (Case rep.)
Garzon K (*see* Obermayer A et al). 2010;52:1255-61
Gebhard MM (*see* Beller CJ et al). 2010;52:1572-9 (Basic research studies)
Geenen IL, Nyilas L, Stephen MS, Makeham V, White GH, Verran DJ. Prosthetic lower extremity hemodialysis access grafts have satisfactory patency despite a high incidence of infection. 2010; 52:1546-50
Gelabert H (*see* Lawrence PF et al). 2010;52:388-93
Gelijns A (*see* Giacovelli JK et al). 2010;52:
Gelijns AC (*see* Kent KC et al). 2010;52:539-48
Gemayel G (*see* Christenson JT et al). 2010;52:1234-41
Genovesi S (*see* Bonforte G et al). 2010;52:1551-6
Genty C (*see* Galanaud J-P et al). 2010;52:932-8
George SJ (*see* Lyon CA et al). 2010;52:1301-9
Georgiadis GS (*see* Trellopoulos G et al). 2010;52:751-7
Georgiou N (*see* Griffin MB et al). 2010;52:69-76
Geroulakos G (*see* Griffin MB et al). 2010;52:69-76
Geroulakos G (*see* Nicolaides AN et al). 2010;52:1486-96
Gessner A (*see* Bal A et al). 2010;52:456-9
Geutjes PJ, van der Vliet JA, Faraj KA, de Vries N, van Moerkerk HTB, Wismans RG, Hendriks T, Daamen WF, van Kuppevelt TH. Preparation and characterization of injectable fibrillar type I collagen and evaluation for pseudoaneurysm treatment in a pig model. 2010;52:1330-8
Ghandforoush A (*see* Hussain J et al). 2010;52:196-8 (Case rep.)
Giacovelli JK, Egorova N, Dayal R, Gelijns A, McKinsey J, Kent KC. Outcomes of carotid stenting compared with endarterectomy are equivalent in asymptomatic patients and inferior in symptomatic patients. 2010;52:
Gifford CSM (*see* Burkhardt CGE et al). 2010;52:91-6
Giles K (*see* Sachs T et al). 2010;52:860-6
Giles KA, Hamdan AD, Pomposelli FB, Wyers MC, Schermerhorn ML. Stroke and death after carotid endarterectomy and carotid artery stenting with and without high risk criteria. 2010;52: 1497-504
Giles KA, Wyers MC, Pomposelli FB, Hamdan AD, Ching YA, Schermerhorn ML. The impact of body mass index on perioperative outcomes of open and endovascular abdominal aortic aneurysm repair from the National Surgical Quality Improvement Program, 2005-2007. 2010;52:1471-7
Gillespie D (*see* Keefer A et al). 2010;52:873-7
Gillespie D (*see* Vasquez MA et al). 2010;52:1387-96
Gillot C (*see* Uhl J-F et al). 2010;52:714-9
Ginghina C (*see* Stoica AL et al). 2010;52:103-9
Godshall CJ (*see* Becher RD et al). 2010;52:1041-4
Godshall CJ (*see* Fleming SH et al). 2010;52:953-8
Godshall CJ (*see* Moore PS et al). 2010;52:118-26
Godshall CJ, Moore PS, Fleming SH, Andrews JS, Hansen KJ, Hoyle JR, Edwards MS. A vascular disease educational program in the preclinical years of medical school increases student interest in vascular disease. 2010;52:775-81
Goettler CE (*see* Durham CA et al). 2010;52:884-90
Goldin I (*see* Shemesh D et al). 2010;52:495-8
Goldstein LJ, Halpern JA, Rezayat C, Gallagher KA, Sambol EB, Bush HL, Karwowski JK. Endovascular aneurysm repair in nonagenarians is safe and effective. 2010;52:1140-6
Golfier V (*see* Mahe G et al). 2010;52:62-8
Goodney P (*see* Powell RJ et al). 2010;52:1525-30
Goodney PP (*see* Bertges DJ et al). 2010;52:674-83
Goodney PP (*see* De Martino RR et al). 2010;52:5-12
Goodney PP (*see* Stone DH et al). 2010;52:127-31
Goodney PP, Nolan BW, Eldrup-Jorgensen J, Likosky DS, Cronenwett JL, Vascular Study Group of Northern New England. Restenosis after carotid endarterectomy in a multicenter regional registry. 2010;52:897-905
Gopal K (*see* Usoh F et al). 2010;52:394-9
Gordon P, O'Donnell ME, Collins A, Gardiner KR, Irwin T. Successful endovascular treatment of multiple mesenteric arterial aneurysms associated with alpha-1-antitrypsin deficiency. 2010;52:467-70
Gorgoulis VG (*see* Sigala F et al). 2010;52:704-13
Gornik HL (*see* Marine L et al). 2010;52:1350-3
Goto S (*see* Takagi H et al). 2010;52:1103-8
Gouicem D (*see* Abouliatim I et al). 2010;52:1665-7
Grassi V (*see* Trimarchi S et al). 2010;52:261-6
Gray W (*see* Matsumura JS et al). 2010;52:576-83
Greco G (*see* Kent KC et al). 2010;52:539-48
Green SY (*see* Schwill S et al). 2010;52:1034-7 (Case rep.)
Greenberg RK (*see* Fillinger MF et al). 2010;52:1022-33
Greenberg RK, (*see* Coscas R et al). 2010;52:272-81
Greenfield L (*see* Criado E et al). 2010;52:406-11
Greenhalgh RM (*see* Sweeting MJ et al). 2010;52:1-4
Greisler HP (*see* Brewster LP et al). 2010;52:1697-8 (Surgical ethics challenges)
Grewal NK (*see* Comerota AJ et al). 2010;52:243-7
Griffin M (*see* Nicolaides AN et al). 2010;52:1486-96
Griffin MB, Kyriacou E, Pattichis C, Bond D, Kakkos SK, Sabetai M, Geroulakos G, Georgiou N, Doré CJ, Nicolaides A. Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms. 2010;52:69-76
Grigoryev DN (*see* Santora RJ et al). 2010;52:1003-14
Grisafi JL, Boiteau G, Detschelt E, Potts J, Kiproff P, Muluk SC. Endoluminal treatment of type IA endoleak with Onyx. 2010; 52:1346-9
Gueddi S (*see* Christenson JT et al). 2010;52:1234-41
Guo D (*see* Dong Z et al). 2010;52:1450-8
Gupta R (*see* Marine L et al). 2010;52:1350-3
Gupta V, Shaik I, Abbas J, Nazzal M. Iliofemoral venous thrombosis from external compression by a vesical diverticulum. 2010;52:1671-3
Guralnik JM (*see* Singh N et al). 2010;52:624-31
Gutowski P (*see* Wiernicki I et al). 2010;52:691-6
Guzzo JL, Chang K, Demos J, Black JH, Freischlag JA. Preoperative thrombolysis and venoplasty affords no benefit in patency following first rib resection and scalenectomy for subacute and chronic subclavian vein thrombosis. 2010;52:658-63

H

Hackney LA (*see* Abularrage CJ et al). 2010;52:314-22
Hafez H (*see* Thompson A et al). 2010;52:55-61
Hagberg R (*see* Sachs T et al). 2010;52:860-6
Halbritter K (*see* Schwarz T et al). 2010;52:1246-50
Halmesmäki K (*see* Arvela E et al). 2010;52:1218-25
Halpern JA (*see* Goldstein LJ et al). 2010;52:1140-6
Hamady M (*see* Willaert W et al). 2010;52:1700-5
Hamaishi M (*see* Mizukami T et al). 2010;52:1580-6
Hamano K (*see* Kubo M et al). 2010;52:429-34 (Basic research studies)
Hamawy A (*see* Causey MW et al). 2010;52:1397-400
Hamblin M, Chang L, Zhang H, Yang K, Zhang J, Chen YE. Vascular smooth muscle cell peroxisome proliferator-activated receptor-γ deletion promotes abdominal aortic aneurysms. 2010;52:984-93

Hamdan A (*see* Sachs T et al). 2010;52:860-6
Hamdan AD (*see* Giles KA et al). 2010;52:1471-7
Hamming JF (*see* Bosman W-MPF et al). 2010;52:152-8
Hamulyák K (*see* ten Cate-Hoek AJ et al). 2010;52:132-8
Han X-W (*see* Ding P-X et al). 2010;52:782-4
Han DK, Fishman EW, Walkup MH, Olin JW, Marin ML, Faries PL. A rare case of familial carotid body tumor in a patient with bilateral fibromuscular dysplasia. 2010;52:746-50
Hancock A (*see* Mazari FAK et al). 2010;52:1564-71
Hancock H, Rasmussen TE, Walker AJ, Rich NM. History of temporary intravascular shunts in the management of vascular injury. 2010;52:1405-9
Hansen KJ (*see* Fleming SH et al). 2010;52:953-8
Hansen KJ (*see* Godshall CJ et al). 2010;52:775-81
Hansen KJ (*see* Moore PS et al). 2010;52:118-26
Hansler J (*see* Endean ED et al). 2010;52:720-2 (Case rep.)
Hanson LA (*see* Martin JD et al). 2010;52:1094-100
Haq F (*see* Sales CM et al). 2010;52:1251-4
Hara T (*see* Yoshino M et al). 2010;52:1401-4
Harding SA (*see* Burdess A et al). 2010;52:697-703
Harris JR (*see* Forbes TL et al). 2010;52:45-8
Harris PL (*see* Manning BJ et al). 2010;52:491-4
Harris PW (*see* Davies MG et al). 2010;52:1478-85
Hartog AG, Algra A, Moll FL, de Borst GJ. Mechanisms of gender-related outcome differences after carotid endarterectomy. 2010;52:1062-71 (Presidential address)
Hartung O (*see* Boufi M et al). 2010;52:1211-7
Hartung O, Azghari A, Barthelemy P, Boufi M, Alimi YS. Laparoscopic transposition of the left renal vein into the inferior vena cava for nutcracker syndrome. 2010;52:738-41
Hasan E (*see* Kurtoglu M et al). 2010;52:1262-71
Hasegawa T (*see* Munezane T et al). 2010;52:967-74 (Basic research studies)
Hass SM (*see* AbuRahma AF et al). 2010;52:1180-7
Hassoun HT (*see* Santora RJ et al). 2010;52:1003-14
Haumer M (*see* Hoke M et al). 2010;52:383-7
Haverich A (*see* Bisdas T et al). 2010;52:323-30
Haynatzki GR (*see* Koutakis P et al). 2010;52:340-7
Hechelhammer L, Rancic Z, Pfiffner R, Mayer D, Meier T, Lachat M, Pfammatter T. Midterm outcome of endovascular repair of ruptured isolated iliac artery aneurysms. 2010;52:1159-63
Hedin U (*see* Ekstrand J et al). 2010;52:167-75 (Basic research studies)
Hehrlein C (*see* Duerschmied D et al). 2010;52:1531-6
Heidenhain C (*see* Heise M et al). 2010;52:421-8 (Basic research studies)
Heidsieck D (*see* Jongkind V et al). 2010;52:760-7 (Review article)
Heise M, Kirschner P, Rabsch A, Zanow J, Settmacher U, Heidenhain C. In vitro testing of a newly developed arteriovenous double-outflow graft. 2010;52:421-8 (Basic research studies)
Heldman M (*see* West CA et al). 2010;52:1164-72
Hendriks EJM (*see* Nicolaï SPA et al). 2010;52:1226-33
Hendriks T (*see* Geutjes PJ et al). 2010;52:1330-8
Henke PK. Invited commentary. 2010;52:1262-71
Herlambang B (*see* Mizukami T et al). 2010;52:1580-6
Hernandez AV (*see* Coscas R et al). 2010;52:272-81
Hess PJ (*see* Lee WA et al). 2010;52:1459-63
Heyden JV (*see* Li Y et al). 2010;52:1716-22
HGF-0205 Trial Investigators (*see* Powell RJ et al). 2010;52:1525-30
Higashi M (*see* Tanaka H et al). 2010;52:479
Hinchliffe R (*see* Patterson BO et al). 2010;52:25-30
Hinchliffe RJ (*see* Malkawi AH et al). 2010;52:480-8 (Review article)
Hinchliffe RJ (*see* Veith FJ et al). 2010;52:1087-93
Hingorani A (*see* Usoh F et al). 2010;52:394-9
Hiramoto JS (*see* Monahan TS et al). 2010;52:303-7
Hirano K (*see* Soga Y et al). 2010;52:608-15
Hislop S (*see* Keefer A et al). 2010;52:873-7
Ho LA (*see* Criqui MH et al). 2010;52:85-90
Hoch JR (*see* Bobadilla JL et al). 2010;52:267-71
Hoke M, Koppensteiner R, Schillinger M, Haumer M, Minar E, Wiesbauer F, Huber CD, Mlekusch W. D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures. 2010;52:383-7
Holt PJ (*see* Malkawi AH et al). 2010;52:480-8 (Review article)
Holt PJ (*see* Patterson BO et al). 2010;52:25-30
Houbballah R, Majewski M, Becquemin J-P. Significant sac retraction after endovascular aneurysm repair is a robust indicator of durable treatment success. 2010;52:878-83
Howe F (*see* Choke E et al). 2010;52:159-66
Hoyle JR (*see* Godshall CJ et al). 2010;52:775-81
Hsiao J-F, Chou H-H, Hsu L-A, Wu L-S, Yang C-W, Hsu T-S, Chang C-J. Vascular changes at the puncture segments of arteriovenous fistula for hemodialysis access. 2010;52:669-73
Hsu L-A (*see* Hsiao J-F et al). 2010;52:669-73
Hsu S-L (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Hsu T-S (*see* Hsiao J-F et al). 2010;52:669-73
Hsu W-C (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Hu S (*see* Jiang J et al). 2010;52:1278-82
Huang H-S (*see* Chen Y-S et al). 2010;52:1310-20
Huang Z (*see* Xiao Y et al). 2010;52:684-90 (Basic research studies)
Huber CD (*see* Hoke M et al). 2010;52:383-7
Humphries SE (*see* Thompson A et al). 2010;52:55-61
Hung R (*see* Lee GS et al). 2010;52:1058-60
Hupp JA (*see* Martin JD et al). 2010;52:1094-100
Husmann M (*see* Willenberg T et al). 2010;52:664-8
Hussain J, Strumpf R, Ghandforoush A, Wheatley G, Sutherland J. Transcatheter closure of recurrent aortic pseudoaneurysm previously treated by Amplatzer occluder device. 2010;52:196-8 (Case rep.)
Huynh TM (*see* Liem TK et al). 2010;52:651-7

I

Ichikawa S (*see* Kumakura H et al). 2010;52:110-7
Iida O (*see* Soga Y et al). 2010;52:608-15
Ilarraza A (*see* Timaran CH et al). 2010;52:1188-95
Ilic N (*see* Dragas M et al). 2010;52:1658-61
Iliodromitis EK (*see* Sigala F et al). 2010;52:704-13
Illig KA (*see* Keefer A et al). 2010;52:873-7
Illig KA. An improved method of exposure for transaxillary first rib resection. 2010;52:248-9 (Basic research studies)
Indes JE, Mandawat A, Tuggle CT, Muhs B, Sosa JA. Endovascular procedures for aorto-iliac occlusive disease are associated with superior short-term clinical and economic outcomes compared with open surgery in the inpatient population. 2010;52:1173-9
Irrera N (*see* Bitto A et al). 2010;52:1015-21
Irwin T (*see* Gordon P et al). 2010;52:467-70
Ishiguro Y (*see* Kondo N et al). 2010;52:1713-5
Ishii H, Kumada Y, Toriyama T, Aoyama T, Takahashi H, Murohara T. Prognostic values of C-reactive protein levels on clinical

outcome after endovascular therapy in hemodialysis patients with peripheral artery disease. 2010;52:854-9
Ito Y (*see* Nakamura Y et al). 2010;52:1061
Ivancev K (*see* Manning BJ et al). 2010;52:491-4
Iwai T (*see* Kondo N et al). 2010;52:1713-5
Iwasaki T (*see* Kumakura H et al). 2010;52:110-7

J

Jackson BM (*see* Brinster CJ et al). 2010;52:460-3
Jackson BM (*see* Foley PJ et al). 2010;52:850-3
Jacob T (*see* Usoh F et al). 2010;52:394-9
Jacobs MJ (*see* Bosman W-MPF et al). 2010;52:152-8
Jacomella V (*see* Willenberg T et al). 2010;52:664-8
Jain M (*see* Andersen ND et al). 2010;52:1608-15
Jenkins J (*see* Kruger AJ et al). 2010;52:1518-24
Jenkins JS (*see* Kruger AJ et al). 2010;52:1518-24
Jennings A (*see* van Walraven C et al). 2010;52:282-9
Jetty P (*see* van Walraven C et al). 2010;52:282-9
Jiang J, Ding X, Zhang G, Su Q, Wang Z, Hu S. Spontaneous retroperitoneal hematoma associated with iliac vein rupture. 2010;52:1278-82
Jim J, Sanchez LA, Rubin BG. Use of a surgeon-modified branched thoracic endograft to preserve an aortorenal bypass during treatment of an intercostal patch aneurysm. 2010;52:730-3
Jimenez JC (*see* Lawrence PF et al). 2010;52:388-93
Jimenez JC (*see* Tulloch AW et al). 2010;52:1283-9
Johanning JM (*see* Koutakis P et al). 2010;52:340-7
Johnson LW (*see* West CA et al). 2010;52:1164-72
Jones C, Badger SA, Boyd CS, Soong CV. The impact of radiation dose exposure during endovascular aneurysm repair on patient safety. 2010;52:298-302
Jones D (*see* Crisostomo PR et al). 2010;52:1272-7
Jones JW, McCullough LB. I know about Jack and you're no Jack Kevorkian. 2010;52:489-90 (Surgical ethics challenges)
Jones JW, McCullough LB. The ethical hierarchy of do not resuscitate orders: Never say never. 2010;52:1384-6 (Surgical ethics challenges)
Jongkind V, Akkersdijk GJM, Yeung KK, Wisselink W. A systematic review of endovascular treatment of extensive aortoiliac occlusive disease. 2010;52:1376-83
Jongkind V, Yeung KK, Akkersdijk GJM, Heidsieck D, Reitsma JB, Tangelder GJ, Wisselink W. Juxtarenal aortic aneurysm repair. 2010;52:760-7 (Review article)
Jonker FHW (*see* Trimarchi S et al). 2010;52:261-6
Jonker FHW, Verhagen HJM, Mojibian H, Davis KA, Moll FL, Muhs BE. Aortic endograft sizing in trauma patients with hemodynamic instability. 2010;52:39-44
Joven J (*see* Alegret JM et al). 2010;52:867-72
JUVENTAS and SMART study groups (*see* Sprengers RW et al). 2010;52:843-9

K

Kakkos SK (*see* Griffin MB et al). 2010;52:69-76
Kakkos SK (*see* Nicolaides AN et al). 2010;52:1486-96
Kamota T (*see* Kubo M et al). 2010;52:429-34 (Basic research studies)
Kanai H (*see* Kumakura H et al). 2010;52:110-7
Kang J, Albadawi H, Casey PJ, Abbruzzese TA, Patel VI, Yoo H-J, Cambria RP, Watkins MT. The effects of systemic hypothermia on a murine model of thoracic aortic ischemia reperfusion. 2010;52:435-43 (Basic research studies)
Kang WC (*see* Coscas R et al). 2010;52:272-81
Kao H-L (*see* Lin M-S et al). 2010;52:308-13
Kapoulas KC (*see* Trellopoulos G et al). 2010;52:751-7
Karabay O (*see* Kurtoglu M et al). 2010;52:1262-71
Karck M (*see* Beller CJ et al). 2010;52:1572-9 (Basic research studies)
Karwowski JK (*see* Goldstein LJ et al). 2010;52:1140-6
Kasama S (*see* Kumakura H et al). 2010;52:110-7
Kasashima F (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Kasashima S, Zen Y, Kawashima A, Endo M, Matsumoto Y, Kasashima F, Ohtake H, Nakanuma Y. A clinicopathologic study of immunoglobulin G4-related sclerosing disease of the thoracic aorta. 2010;52:1587-95 (Basic research studies)
Kashyap VS (*see* Marine L et al). 2010;52:1350-3
Kashyap VS, Bishop PD, Bena JF, Rosa K, Sarac TP, Ouriel K. A pilot, prospective evaluation of a direct thrombin inhibitor, bivalirudin (Angiomax), in patients undergoing lower extremity bypass. 2010;52:369-74
Kasirajan K (*see* Chung J et al). 2010;52:1435-41
Katada Y, Ouchi K, Wakita T, Nozaki M. Liquid sclerotherapy for posttraumatic arteriovenous fistula of the radialis indicis artery. 2010;52:1343-5
Kawarada O, Yokoi Y, Takemoto K. Practical use of duplex echo-guided recanalization of chronic total occlusion in the iliac artery. 2010;52:475-8
Kawashima A (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Keefer A, Hislop S, Singh MJ, Gillespie D, Illig KA. The influence of aneurysm size on anatomic suitability for endovascular repair. 2010;52:873-7
Keilani ZM, Berne JD, Agko M. Bilateral internal carotid and vertebral artery dissection after a horse-riding injury. 2010;52:1052-7
Kent KC (*see* Giacovelli JK et al). 2010;52:
Kent KC, Zwolak RM, Egorova NN, Riles TS, Manganaro A, Moskowitz AJ, Gelijns AC, Greco G. Analysis of risk factors for abdominal aortic aneurysm in a cohort of more than 3 million individuals. 2010;52:539-48
Khan JA (*see* Mazari FAK et al). 2010;52:1564-71
Khoynezhad A, Walot I, Kruse MJ, Rapae T, Donayre CE, White RA. Distribution of intimomedial tears in patients with type B aortic dissection. 2010;52:562-8
Khurana A (*see* Gardner AW et al). 2010;52:1204-10
Kim JY (*see* Cho J-S et al). 2010;52:1127-34
Kim Y-W (*see* Park U-J et al). 2010;52:222
Kimura K (*see* Ohtake H et al). 2010;52:1464-70
Kinikini DV (*see* Shelgikar C et al). 2010;52:920-4
Kiproff P (*see* Grisafi JL et al). 2010;52:1346-9
Kirschner P (*see* Heise M et al). 2010;52:421-8 (Basic research studies)
Klar E (*see* Bünger CM et al). 2010;52:1357-61
Klein J (*see* Crisostomo PR et al). 2010;52:1272-7
Kleinman, L (*see* Vouyouka AG et al). 2010;52:1196-203
Klodell CT (*see* Lee WA et al). 2010;52:1459-63
Ko CY (*see* Osborne NH et al). 2010;52:400-5
Kobeiter H (*see* Abouliatim I et al). 2010;52:1665-7
Koka P (*see* Tong MZ-Y et al). 2010;52:31-8
Koksoy C (*see* Kurtoglu M et al). 2010;52:1262-71
Komalavilas P (*see* Lopes LB et al). 2010;52:1596-607
Koncar I (*see* Dragas M et al). 2010;52:1658-61
Kondo N, Koyama M, Iwai T, Ozaki T, Ishiguro Y, Noda H. A durable iliac-axillary and axillary-carotid bypass for cerebral

ischemia due to Takayasu arteritis allowed successful pregnancies and deliveries. 2010;52:1713-5
Konno H (*see* Unno N et al). 2010;52:946-52
Koppensteiner R (*see* Hoke M et al). 2010;52:383-7
Korhonen M (*see* Arvela E et al). 2010;52:1218-25
Kotsinas A (*see* Sigala F et al). 2010;52:704-13
Koutakis P, Johanning JM, Haynatzki GR, Myers SA, Stergiou N, Longo GM, Pipinos II. Abnormal joint powers before and after the onset of claudication symptoms. 2010;52:340-7
Koutsouki E (*see* Lyon CA et al). 2010;52:1301-9
Koyama M (*see* Kondo N et al). 2010;52:1713-5
Kröger J-C (*see* Bünger CM et al). 2010;52:1357-61
Kraiss LW (*see* Shelgikar C et al). 2010;52:920-4
Kranendonk S (*see* Smolderen KG et al). 2010;52:637-44
Kreienberg PB (*see* Mehta M et al). 2010;52:1442-9
Krishnasastry KV (*see* De Marco Garcia LP et al). 2010;52:139-44
Kruger AJ, Walker PJ, Foster WJ, Jenkins JS, Boyne NS, Jenkins J. Important observations made managing carotid body tumors during a 25-year experience. 2010;52:1518-24
Krummel TM (*see* Rich NM et al). 2010;52:508-11
Kruse MJ (*see* Khoynezhad A et al). 2010;52:562-8
Kubo M (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Kubo M, Li T-S, Kamota T, Ohshima M, Shirasawa B, Hamano K. Extracorporeal shock wave therapy ameliorates secondary lymphedema by promoting lymphangiogenesis. 2010;52:429-34 (Basic research studies)
Kumada Y (*see* Ishii H et al). 2010;52:854-9
Kumakura H, Kanai H, Aizaki M, Mitsui K, Araki Y, Kasama S, Iwasaki T, Ichikawa S. The influence of the obesity paradox and chronic kidney disease on long-term survival in a Japanese cohort with peripheral arterial disease. 2010;52:110-7
Kureck I (*see* Wiegand S et al). 2010;52:742-5
Kurtoglu M, Koksoy C, Hasan E, Akcalı Y, Karabay O, Filizcan U, TROMBOTEK Study Group. Long-term efficacy and safety of once-daily enoxaparin plus warfarin for the outpatient ambulatory treatment of lower-limb deep vein thrombosis in the TROMBOTEK trial. 2010;52:1262-71
Kuzniec S, Natal SRB, Werebe EC, Wolosker N. Videothoracoscopic-guided management of a central vein perforation during hemodialysis catheter placement. 2010;52:1354-6
Kwolek CJ (*see* Abularrage CJ et al). 2010;52:314-22
Kyriacou E (*see* Griffin MB et al). 2010;52:69-76
Kyriacou E (*see* Nicolaides AN et al). 2010;52:1486-96

L

Labropoulos N (*see* Nicolaides AN et al). 2010;52:1486-96
Labrosse MR (*see* Beller CJ et al). 2010;52:1572-9 (Basic research studies)
Lachat M (*see* Hechelhammer L et al). 2010;52:1159-63
Lamparello PJ (*see* Fonseca R et al). 2010;52:556-61
LaMuraglia GM (*see* Abularrage CJ et al). 2010;52:314-22
LaMuraglia GM. Invited Commentary. 2010;52:1518-24
Landry GJ (*see* Liem TK et al). 2010;52:651-7
Lang W (*see* Bal A et al). 2010;52:456-9
Lang JL, Minei JP, Modrall JG, Clagett GP, Valentine RJ. The limitations of thoracic endovascular aortic repair in altering the natural history of blunt aortic injury. 2010;52:290-7
Laroche J-P (*see* Galanaud J-P et al). 2010;52:932-8
Laviolle B (*see* Mahe G et al). 2010;52:62-8
Lawlor DK (*see* Forbes TL et al). 2010;52:45-8
Lawrence PF (*see* Tulloch AW et al). 2010;52:1283-9
Lawrence PF, Chandra A, Wu M, Rigberg D, DeRubertis B, Gelabert H, Jimenez JC, Carter V. Classification of proximal endovenous closure levels and treatment algorithm. 2010;52:388-93
Lazarides MK (*see* Trellopoulos G et al). 2010;52:751-7
Lee H (*see* Abularrage CJ et al). 2010;52:19-24
Lee JN (*see* Gale SS et al). 2010;52:645-50
Lee V (*see* Riles TS et al). 2010;52:914-9
Lee GS, Brawley J, Hung R. Complex subclavian artery pseudoaneurysm causing failure of endovascular stent repair with salvage by percutaneous thrombin injection. 2010;52:1058-60
Lee W-J, Ou H-C, Hsu W-C, Chou M-M, Tseng J-J, Hsu S-L, Tsai K-L, Sheu WH-H. Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells. 2010;52:1290-300 (Basic research studies)
Lee WA, Martin TD, Hess PJ, Beaver TM, Klodell CT. First United States experience of the TX2 Pro-Form thoracic delivery system. 2010;52:1459-63
Leers S (*see* Fernandez N et al). 2010;52:834-42
Leijdekkers VJ (*see* Tolenaar JL et al). 2010;52:1045-7
LeMaire SA (*see* Schwill S et al). 2010;52:1034-7 (Case rep.)
Lepäntalo M (*see* Arvela E et al). 2010;52:1218-25
Leverson GE (*see* Bobadilla JL et al). 2010;52:267-71
Li H (*see* Yao Y et al). 2010;52:1662-4
Li H-Y (*see* Lin M-S et al). 2010;52:308-13
Li T, Zhang WW, Bai W, Zhai S, Pang Z. Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction. 2010;52:1242-5
Li T-S (*see* Kubo M et al). 2010;52:429-34 (Basic research studies)
Li Y, Castaldo J, Heyden JV, Plokker HWM. Is carotid artery disease responsible for perioperative strokes after coronary artery bypass surgery. 2010;52:1716-22
Li Y-D (*see* Ding P-X et al). 2010;52:782-4
Liang H (*see* Ouyang C et al). 2010;52:1048-51
Liang HD (*see* Yao Y et al). 2010;52:1662-4
Liao Y (*see* Singh N et al). 2010;52:624-31
Lie ML (*see* Santora RJ et al). 2010;52:1003-14
Liem TK, Huynh TM, Moseley SE, Minjarez RC, Landry GJ, Mitchell EL, DeLoughery TG, Moneta GL. Symptomatic perioperative venous thromboembolism is a frequent complication in patients with a history of deep vein thrombosis. 2010;52:651-7
Ligero C (*see* Alegret JM et al). 2010;52:867-72
Likosky DS (*see* Bertges DJ et al). 2010;52:674-83
Likosky DS (*see* Goodney PP et al). 2010;52:897-905
Lilly MP. Invited commentary. 2010;52:539-48
Lin C-H (*see* Lin M-S et al). 2010;52:308-13
Lin PH (*see* Bechara CF et al). 2010;52:1682-96
Lin M-S, Chen Y-H, Chao C-C, Lin C-H, Li H-Y, Chao C-L, Chen M-F, Kao H-L. Catheter-based neurosalvage for acute embolic complication during carotid intervention. 2010;52:308-13
Link A (*see* Comerota AJ et al). 2010;52:723-9
Liu B (*see* Yao Y et al). 2010;52:1662-4
Liu K (*see* Singh N et al). 2010;52:624-31
Liu N-F, Lu Q, Yan Z-X. Lymphatic malformation is a common component of Klippel-Trenaunay syndrome. 2010;52:1557-63
Loftus IM (*see* Choke E et al). 2010;52:159-66
Loftus IM (*see* Malkawi AH et al). 2010;52:480-8 (Review article)
Loftus IM (*see* Patterson BO et al). 2010;52:25-30
LoGerfo FW (*see* Andersen ND et al). 2010;52:1608-15
Longo GM (*see* Koutakis P et al). 2010;52:340-7

Lopes LB, Brophy CM, Flynn CR, Yi Z, Bowen BP, Smoke C, Seal B, Panitch A, Komalavilas P. A novel cell permeant peptide inhibitor of MAPKAP kinase II inhibits intimal hyperplasia in a human saphenous vein organ culture model. 2010;52:1596-607
Lough E (*see* AbuRahma AF et al). 2010;52:1180-7
Lu M-J (*see* Chen Y-S et al). 2010;52:1310-20
Lu Q (*see* Liu N-F et al). 2010;52:1557-63
Lumsden AB (*see* Bismuth J et al). 2010;52:1072-80
Lumsden AB (*see* Davies MG et al). 2010;52:1478-85
Lyon CA, Koutsouki E, Aguilera CM, Blaschuk OW, George SJ. Inhibition of N-cadherin retards smooth muscle cell migration and intimal thickening via induction of apoptosis. 2010;52:1301-9

M

Ma M-C (*see* Chen Y-S et al). 2010;52:1310-20
Mahe G, Ronziere T, Laviolle B, Golfier V, Cochery T, De Bray J-M, Paillard F. An unfavorable dietary pattern is associated with symptomatic ischemic stroke and carotid atherosclerosis. 2010;52:62-8
Majeski M (*see* Abouliatim I et al). 2010;52:1665-7
Majewski M (*see* Houbballah R et al). 2010;52:878-83
Makaroun M (*see* Fernandez N et al). 2010;52:834-42
Makaroun MS (*see* Chaer RA et al). 2010;52:569-75
Makaroun MS (*see* Cho J-S et al). 2010;52:1127-34
Makeham V (*see* Geenen IL et al). 2010;52:1546-50
Maldonado TS (*see* Fonseca R et al). 2010;52:556-61
Malek JY (*see* Andersen ND et al). 2010;52:1608-15
Maletzki P (*see* Duerschmied D et al). 2010;52:1531-6
Malgor RD (*see* Ricotta JJ et al). 2010;52:785-9
Malkawi AH, Hinchliffe RJ, Xu Y, Holt PJ, Loftus IM, Thompson MM. Patient-specific biomechanical profiling in abdominal aortic aneurysm development and rupture. 2010;52:480-8 (Review article)
Mamdani M (*see* Altoijry A et al). 2010;52:1-2
Manabe H (*see* Takagi H et al). 2010;52:1103-8
Mandawat A (*see* Indes JE et al). 2010;52:1173-9
Mandolfo S (*see* Bonforte G et al). 2010;52:1551-6
Manganaro A (*see* Kent KC et al). 2010;52:539-48
Mangano S (*see* Bonforte G et al). 2010;52:1551-6
Mani K, Rantatalo M, Bjurholm A, Thelin S, Eriksson L-G, Wanhainen A. Aortic rupture after spinal correction for scoliosis in the presence of a thoracic stent graft. 2010;52:1653-7 (Case rep.)
Manning BJ, Ivancev K, Harris PL. In situ fenestration in the aortic arch. 2010;52:491-4
Mano Y (*see* Unno N et al). 2010;52:946-52
Mapes JE (*see* Martin JD et al). 2010;52:1094-100
Marin M (*see* Vouyouka AG et al). 2010;52:1196-203
Marin ML (*see* Han DK et al). 2010;52:746-50
Marine L, Gupta R, Gornik HL, Kashyap VS. Glue embolus complicating the endovascular treatment of a patient with Locys-Dietz syndrome. 2010;52:1350-3
Markantonis S (*see* Sigala F et al). 2010;52:704-13
Markose G, Miller FNAC, Bolia A. Subintimal angioplasty for femoro-popliteal occlusive disease. 2010;52:1410-6
Markovic M (*see* Dragas M et al). 2010;52:1658-61
Marks N (*see* Usoh F et al). 2010;52:394-9
Marone LK (*see* Fernandez N et al). 2010;52:834-42
Marone LK (*see* Wagner JK et al). 2010;52:216-8
Marone EM, Coppi G, Tshomba Y, Chiesa R. Eight-year experience with carotid artery stenting for correction of symptomatic and asymptomatic post-endarterectomy defects. 2010;52:1511-7
Marston WA (*see* Vasquez MA et al). 2010;52:1387-96
Martín-Paredero V (*see* Alegret JM et al). 2010;52:867-72
Martin TD (*see* Lee WA et al). 2010;52:1459-63
Martin JD, Warble PB, Hupp JA, Mapes JE, Stanziale SF, Weiss LL, Schiller TB, Hanson LA. A real world analysis of payment per unit time in a Maryland Vascular Practice. 2010;52:1094-100
Martinez AE (*see* Timaran CH et al). 2010;52:1188-95
Mastracci TM (*see* Coscas R et al). 2010;52:272-81
Masud F (*see* Davies MG et al). 2010;52:1478-85
Matoba T (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Matsui M (*see* Takagi H et al). 2010;52:1675-81 (Review article)
Matsui O (*see* Ohtake H et al). 2010;52:1464-70
Matsumoto Y (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Matsumura JS, Gray W, Chaturvedi S, Gao X, Cheng J, Verta P, CAPTURE 2 Investigators and Executive Committee. CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection. 2010;52:576-83
Mayer D (*see* Hechelhammer L et al). 2010;52:1159-63
Mazari FAK, Carradice D, Rahman MNAA, Khan JA, Mockford K, Mehta T, McCollum PT, Chetter IC. An analysis of relationship between quality of life indices and clinical improvement following intervention in patients with intermittent claudication due to femoropopliteal disease. 2010;52:77-84
Mazari FAK, Mockford K, Barnett C, Khan JA, Brown B, Smith L, Polman RC, Hancock A, Vanicek NK, Chetter IC. Hull early walking aid for rehabilitation of transtibial amputees - randomized controlled trial (HEART). 2010;52:1564-71
McAfee-Bennett S (*see* Beeman BR et al). 2010;52:1147-52
McCollum PT (*see* Mazari FAK et al). 2010;52:77-84
McCullough LB (*see* Jones JW et al). 2010;52:1384-6 (Surgical ethics challenges)
McDermott MM (*see* Criqui MH et al). 2010;52:85-90
McDermott MM (*see* Singh N et al). 2010;52:624-31
McEnaney R (*see* Fernandez N et al). 2010;52:834-42
McKinsey J (*see* Giacovelli JK et al). 2010;52:
McKinsey JF (*see* Fillinger MF et al). 2010;52:1022-33
McLafferty RB (*see* Vasquez MA et al). 2010;52:1387-96
McNally MM (*see* Durham CA et al). 2010;52:884-90
McQuade K, Gable D, Pearl G, Theune B, Black S. Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease. 2010;52:584-91
Mehta M (*see* Rahimi SA et al). 2010;52:891-6
Mehta M, Darling RC, Taggert JB, Roddy SP, Sternbach Y, Ozsvath KJ, Kreienberg PB, Paty PSK. Outcomes of planned celiac artery coverage during TEVAR. 2010;52:1153-8
Mehta M, Sternbach Y, Taggert JB, Kreienberg PB, Roddy SP, Paty PSK, Ozsvath KJ, Darling RC. Long-term outcomes of secondary procedures after endovascular aneurysm repair. 2010;52:1442-9
Mehta M. Endovascular aneurysm repair for ruptured abdominal aortic aneurysm: How I do it. 2010;52:1706-12
Mehta T (*see* Mazari FAK et al). 2010;52:77-84
Meier GH (*see* Schmieder GC et al). 2010;52:375-82
Meier T (*see* Hechelhammer L et al). 2010;52:1159-63
Meissner MH (*see* Vasquez MA et al). 2010;52:1387-96
Melissano G (*see* Civilini E et al). 2010;52:758-9

Mendelsohn FO (*see* Powell RJ et al). 2010;52:1525-30
Mestres C-A (*see* Pereda D et al). 2010;52:1362-4
Midy D (*see* Berard X et al). 2010;52:471-4
Mikhailidis DP (*see* Paraskevas KI et al). 2010;52:1367-75 (Review article)
Millá L (*see* Alegret JM et al). 2010;52:867-72
Miller FNAC (*see* Markose G et al). 2010;52:1410-6
Miller S (*see* Causey MW et al). 2010;52:734-7
Milner R (*see* Chung J et al). 2010;52:1435-41
Minar E (*see* Hoke M et al). 2010;52:383-7
Minei JP (*see* Lang JL et al). 2010;52:290-7
Minjarez RC (*see* Liem TK et al). 2010;52:651-7
Mitchell EL (*see* Liem TK et al). 2010;52:651-7
Mitsui K (*see* Kumakura H et al). 2010;52:110-7
Mizukami T, Orihashi K, Herlambang B, Takahashi S, Hamaishi M, Okada K, Sueda T. Sodium 4-phenylbutyrate protects against spinal cord ischemia by inhibition of endoplasmic reticulum stress. 2010;52:1580-6
Mizutani T (*see* Yoshino M et al). 2010;52:1401-4
Mlekusch W (*see* Hoke M et al). 2010;52:383-7
Mockford K (*see* Mazari FAK et al). 2010;52:1564-71
Modrall JG (*see* Lang JL et al). 2010;52:290-7
Modrall JG (*see* Timaran CH et al). 2010;52:1188-95
Moen EK (*see* Powell RJ et al). 2010;52:1525-30
Mohebali J (*see* Shelgikar C et al). 2010;52:920-4
Mohr MC (*see* Durham CA et al). 2010;52:600-7
Mojibian H (*see* Jonker FHW et al). 2010;52:39-44
Moll FL (*see* Derksen WJM et al). 2010;52:592-9
Moll FL (*see* Hartog AG et al). 2010;52:1062-71 (Presidential address)
Moll FL (*see* Jonker FHW et al). 2010;52:39-44
Moll FL (*see* Sprengers RW et al). 2010;52:843-9
Moll FL (*see* Tolenaar JL et al). 2010;52:1045-7
Moll FL (*see* van Keulen JW et al). 2010;52:1081-6
Monahan TS (*see* Andersen ND et al). 2010;52:1608-15
Monahan TS, Chuter TAM, Reilly LM, Rapp JH, Hiramoto JS. Long-term follow-up of neck expansion after endovascular aortic aneurysm repair. 2010;52:303-7
Moneta GL (*see* Liem TK et al). 2010;52:651-7
Montero M (*see* Alegret JM et al). 2010;52:867-72
Montgomery PS (*see* Gardner AW et al). 2010;52:1204-10
Moore FA (*see* Santora RJ et al). 2010;52:1003-14
Moore PS (*see* Godshall CJ et al). 2010;52:775-81
Moore WS (*see* Tulloch AW et al). 2010;52:1283-9
Moore PS, Andrews JS, Craven TE, Davis RP, Corriere MA, Godshall CJ, Edwards MS, Hansen KJ. Trends in vena caval interruption. 2010;52:118-26
Morales C (*see* Coscas R et al). 2010;52:272-81
Morant K (*see* van Walraven C et al). 2010;52:282-9
Morris RJ (*see* Foley PJ et al). 2010;52:850-3
Morris TP (*see* Nicolaides AN et al). 2010;52:1486-96
Moscowitz A (*see* Vouyouka AG et al). 2010;52:1196-203
Moseley SE (*see* Liem TK et al). 2010;52:651-7
Moskowitz AJ (*see* Kent KC et al). 2010;52:539-48
Mosti G, Partsch H. Inelastic bandages maintain their hemodynamic effectiveness over time despite significant pressure loss. 2010;52:925-31
Mottola MT (*see* Stabile E et al). 2010;52:1505-10
Mousa A (*see* AbuRahma AF et al). 2010;52:1180-7
Mueller MT (*see* Shelgikar C et al). 2010;52:920-4
Muhs B (*see* Indes JE et al). 2010;52:1173-9
Muhs BE (*see* Jonker FHW et al). 2010;52:39-44
Muhs BE (*see* Trimarchi S et al). 2010;52:261-6
Muluk SC (*see* Grisafi JL et al). 2010;52:1346-9
Munezane T, Hasegawa T, Suritala, Tanaka A, Okada K, Okita Y. Activation of transglutaminase type 2 for aortic wall protection in a rat abdominal aortic aneurysm formation. 2010;52:967-74 (Basic research studies)
Murohara T (*see* Ishii H et al). 2010;52:854-9
Murtha K (*see* Beeman BR et al). 2010;52:1147-52
Myers SA (*see* Koutakis P et al). 2010;52:340-7

N

Nagahama R (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Nakagawa NK (*see* Erling N et al). 2010;52:1321-9
Nakamura Y, Tagusari O, Seike Y, Ito Y. Persistence of intra-aortic air after circulatory arrest. 2010;52:1061
Nakamura LY, Ferrigni RG, Stone WM, Fowl RJ. Urinary bladder injuries during vascular surgery. 2010;52:453-5 (Case rep.)
Nakano K (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Nakanuma Y (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Nanjundappa A (*see* AbuRahma AF et al). 2010;52:1180-7
Nanto S (*see* Soga Y et al). 2010;52:608-15
Naoum JJ (*see* Bismuth J et al). 2010;52:1072-80
Naoum JJ (*see* Davies MG et al). 2010;52:331-9
Nasir O (*see* Santora RJ et al). 2010;52:1003-14
Natal SRB (*see* Kuzniec S et al). 2010;52:1354-6
Naylor AR (*see* Schneider PA et al). 2010;52:499-507
Naylor R (*see* Nicolaides AN et al). 2010;52:1486-96
Nazzal M (*see* Gupta V et al). 2010;52:1671-3
Newby DE (*see* Burdess A et al). 2010;52:697-703
Ng MKC (*see* Byrom MJ et al). 2010;52:176-95
Nguyen BT (*see* Yu P et al). 2010;52:444-52
Nguyen LL, Barshes NR. Analysis of large databases in vascular surgery. 2010;52:768-74
Nicolaï SPA, Hendriks EJM, Prins MH, Teijink JAW, EXITPAD study group. Optimizing supervised exercise therapy for patients with intermittent claudication. 2010;52:1226-33
Nicolaï SPA, Teijink JAW, Prins MH, Exercise Therapy in Peripheral Arterial Disease (EXITPAD) study group. Multicenter randomized clinical trial of supervised exercise therapy with or without feedback versus walking advice for intermittent claudication. 2010;52:348-55
Nicolaides A (*see* Griffin MB et al). 2010;52:69-76
Nicolaides AN, Kakkos SK, Kyriacou E, Griffin M, Sabetai M, Thomas DJ, Tegos T, Geroulakos G, Labropoulos N, Doré CJ, Morris TP, Naylor R, Abbott AL, Asymptomatic Carotid Stenosis and Risk of Stroke (ACSRS) Study Group. Asymptomatic internal carotid artery stenosis and cerebrovascular risk stratification. 2010;52:1486-96
Nimmo AF (*see* Burdess A et al). 2010;52:697-703
Nishiyama M (*see* Unno N et al). 2010;52:946-52
Nobuyoshi M (*see* Soga Y et al). 2010;52:608-15
Noda H (*see* Kondo N et al). 2010;52:1713-5
Nolan BW (*see* Bertges DJ et al). 2010;52:674-83
Nolan BW (*see* De Martino RR et al). 2010;52:5-12
Nolan BW (*see* Goodney PP et al). 2010;52:897-905
Nolan BW (*see* Stone DH et al). 2010;52:127-31
Nooren M (*see* Smolderen KG et al). 2010;52:637-44
Nordon IM (*see* Patterson BO et al). 2010;52:25-30
Nozaki M (*see* Katada Y et al). 2010;52:1343-5
Nyilas L (*see* Geenen IL et al). 2010;52:1546-50

O

O'Donnell ME (*see* Gordon P et al). 2010;52:467-70
O'Donnell TF. Invited Commentary. 2010;52:1094-100
O'Malley MK (*see* Bismuth J et al). 2010;52:1072-80
Obermayer A, Garzon K. Identifying the source of superficial reflux in venous leg ulcers using duplex ultrasound. 2010;52:1255-61
Oda S, Nagahama R, Nakano K, Matoba T, Kubo M, Sunagawa K, Tominaga R, Egashira K. Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia. 2010;52:412-20 (Basic research studies)
Ogino H (*see* Tanaka H et al). 2010;52:479
Oguntoye MO (*see* Causey MW et al). 2010;52:734-7
Ohashi H (*see* Takahashi Y et al). 2010;52:1365-6
Ohshima M (*see* Kubo M et al). 2010;52:429-34 (Basic research studies)
Ohtake H (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Ohtake H, Kimura K, Sanada J, Matsui O, Watanabe G. Risk factor analysis of thoracic endovascular repair using the Matsui-Kitamura stent graft for acute aortic emergencies in the descending thoracic aorta. 2010;52:1464-70
Okada K (*see* Mizukami T et al). 2010;52:1580-6
Okada K (*see* Munezane T et al). 2010;52:967-74 (Basic research studies)
Okita Y (*see* Munezane T et al). 2010;52:967-74 (Basic research studies)
Olin JW (*see* Han DK et al). 2010;52:746-50
Olschewski M (*see* Duerschmied D et al). 2010;52:1531-6
Olsha O (*see* Shemesh D et al). 2010;52:495-8
OPTIMEV SFMV investigators (*see* Galanaud J-P et al). 2010;52:932-8
Orihashi K (*see* Mizukami T et al). 2010;52:1580-6
Orsini B (*see* Boufi M et al). 2010;52:1211-7
Osborne NH, Ko CY, Upchurch GR, Dimick JB. Evaluating parsimonious risk-adjustment models for comparing hospital outcomes with vascular surgery. 2010;52:400-5
Ou H-C (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Ouchi K (*see* Katada Y et al). 2010;52:1343-5
Ouriel K (*see* Kashyap VS et al). 2010;52:369-74
Ouyang C, Liang H. Urgent interventional bilateral renal artery fenestration for giant pararenal abdominal aortic aneurysm with upper digestive tract obstruction. 2010;52:1048-51
Owens CD. *JAK2 V617F* mutation, mesenteric vein thrombosis, and myeloproliferative disorders. 2010;52:205-7
Ozaki CK (*see* Yu P et al). 2010;52:444-52
Ozaki T (*see* Kondo N et al). 2010;52:1713-5
Ozsvath KJ (*see* Mehta M et al). 2010;52:1442-9

P

Paillard F (*see* Mahe G et al). 2010;52:62-8
Palomares R (*see* Alegret JM et al). 2010;52:867-72
Panetta TF (*see* De Marco Garcia LP et al). 2010;52:139-44
Pang K (*see* Bradbury AW et al). 2010;52:939-45 (Case rep.)
Pang Z (*see* Li T et al). 2010;52:1242-5
Panitch A (*see* Lopes LB et al). 2010;52:1596-607
Panneton JM (*see* Schmieder GC et al). 2010;52:362-8
Panneton JM, (*see* Schmieder GC et al). 2010;52:375-82
Paraskevas KI, Mikhailidis DP, Veith FJ. Mechanisms to explain the poor results of carotid artery stenting (CAS) in symptomatic patients to date and options to improve CAS outcomes. 2010;52:1367-75 (Review article)
Park T (*see* Cho J-S et al). 2010;52:1127-34
Park U-J, Rho Y-N, Yun W-S, Kim Y-W. A giant femoral artery pseudoaneurysm in an infant boy. 2010;52:222
Parker DE (*see* Gardner AW et al). 2010;52:1204-10
Parker FM (*see* Durham CA et al). 2010;52:884-90
Parry DJ, Al-Barjas HS, Chappell L, Rashid ST, Ariëns RAS, Scott DJA. Markers of inflammation in men with small abdominal aortic aneurysm. 2010;52:145-51 (Basic research studies)
Partsch H (*see* Mosti G et al). 2010;52:925-31
Paruchuri V (*see* Abularrage CJ et al). 2010;52:314-22
Pasterkamp G (*see* Derksen WJM et al). 2010;52:592-9
Patel N (*see* Usoh F et al). 2010;52:394-9
Patel VI (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Patel MS, Carpenter JP. The value of the initial post-EVAR computed tomography angiography scan in predicting future secondary procedures using the Powerlink stent graft. 2010;52:1135-9
Patterson BO, Holt PJ, Hinchliffe R, Nordon IM, Loftus IM, Thompson MM. Existing risk prediction methods for elective abdominal aortic aneurysm repair do not predict short-term outcome following endovascular repair. 2010;52:25-30
Pattichis C (*see* Griffin MB et al). 2010;52:69-76
Paty PSK (*see* Mehta M et al). 2010;52:1442-9
Pearl G (*see* McQuade K et al). 2010;52:584-91
Peden EK (*see* Bismuth J et al). 2010;52:1072-80
Peden EK (*see* Davies MG et al). 2010;52:331-9
Pejkic S (*see* Dragas M et al). 2010;52:1658-61
Pereda D, Cartañá R, Sundt TM, Mestres C-A. Seventeen-year follow-up after ascending-to-infrarenal aorta bypass for recurrent coarctation in an adult. 2010;52:1362-4
Perera AD, Willis AK, Fernandez JD, Garrett HE, Wolf BA. Staged total exclusion of the aorta for chronic type B aortic dissection. 2010;52:1339-42 (Case rep.)
Perler BA (*see* Sidawy AN et al). 2010;52:1431-4
Pfammatter T (*see* Hechelhammer L et al). 2010;52:1159-63
Pfiffner R (*see* Hechelhammer L et al). 2010;52:1159-63
Piatek J (*see* Wiernicki I et al). 2010;52:691-6
Pichlmaier M (*see* Bisdas T et al). 2010;52:323-30
Pipinos II (*see* Koutakis P et al). 2010;52:340-7
Pitoulias GA (*see* Donas KP et al). 2010;52:356-61
Pitta X (*see* Trellopoulos G et al). 2010;52:751-7
Pitti A (*see* Fonseca R et al). 2010;52:556-61
Plokker HWM (*see* Li Y et al). 2010;52:1716-22
Podbielski FJ (*see* Willson TD et al). 2010;52:219-21
Pogliani D (*see* Bonforte G et al). 2010;52:1551-6
Poli-de-Figueiredo LF (*see* Erling N et al). 2010;52:1321-9
Polito F (*see* Bitto A et al). 2010;52:1015-21
Polman RC (*see* Mazari FAK et al). 2010;52:1564-71
Pomposelli F (*see* Sachs T et al). 2010;52:860-6
Pomposelli FB (*see* Giles KA et al). 2010;52:1471-7
Popusoi G (*see* Stabile E et al). 2010;52:1505-10
Porras C (*see* Burkhardt CGE et al). 2010;52:91-6
Potts J (*see* Grisafi JL et al). 2010;52:1346-9
Powell CS (*see* Durham CA et al). 2010;52:884-90
Powell JT (*see* Sweeting MJ et al). 2010;52:1-4
Powell JT (*see* Veith FJ et al). 2010;52:1087-93
Powell RJ, Goodney P, Mendelsohn FO, Moen EK, Annex BH, HGF-0205 Trial Investigators. Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: Results of the HGF-0205 trial. 2010;52:1525-30

Pradhan L (*see* Andersen ND et al). 2010;52:1608-15
Pratesi C (*see* Sticchi E et al). 2010;52:97-102
Pratesi G (*see* Sticchi E et al). 2010;52:97-102
Prins MH (*see* Nicolaï SPA et al). 2010;52:1226-33
Prins MH (*see* Ten Bosch JA et al). 2010;52:13-8
Prins MH (*see* ten Cate-Hoek AJ et al). 2010;52:132-8
Propper CBW (*see* Burkhardt CGE et al). 2010;52:91-6
Pulli R (*see* Sticchi E et al). 2010;52:97-102

Q

Quéré I (*see* Galanaud J-P et al). 2010;52:932-8
Quan R (*see* Causey MW et al). 2010;52:1397-400
Quinones-Baldrich WJ (*see* Tulloch AW et al). 2010;52:1283-9

R

Rabe E (*see* Vasquez MA et al). 2010;52:1387-96
Rabsch A (*see* Heise M et al). 2010;52:421-8 (Basic research studies)
Rahimi SA, Darling RC, Mehta M, Roddy SP, Taggert JB, Sternbach Y. Endovascular repair of thoracic aortic traumatic transections is a safe method in patients with complicated injuries. 2010;52:891-6
Rahman MNAA (*see* Mazari FAK et al). 2010;52:77-84
Rampoldi V (*see* Trimarchi S et al). 2010;52:261-6
Rancic Z (*see* Hechelhammer L et al). 2010;52:1159-63
Rantatalo M (*see* Mani K et al). 2010;52:1653-7 (Case rep.)
Rao A (*see* Chaer RA et al). 2010;52:569-75
Rapae T (*see* Khoynezhad A et al). 2010;52:562-8
Raphael M (*see* Adiseshiah M et al). 2010;52:49-54
Rapp JH (*see* Monahan TS et al). 2010;52:303-7
Rashid ST (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Rasmussen TE (*see* Hancock H et al). 2010;52:1405-9
Rasmussen TE (*see* Burkhardt CGE et al). 2010;52:91-6
Rastan A (*see* Schwarz T et al). 2010;52:1246-50
Razuvaev A (*see* Ekstrand J et al). 2010;52:167-75 (Basic research studies)
Reardon MJ (*see* Davies MG et al). 2010;52:1478-85
Reilly LM (*see* Monahan TS et al). 2010;52:303-7
Reitsma JB (*see* Jongkind V et al). 2010;52:760-7 (Review article)
Revesz E (*see* Willson TD et al). 2010;52:219-21
Rezayat C (*see* Goldstein LJ et al). 2010;52:1140-6
Rhee RY (*see* Cho J-S et al). 2010;52:1127-34
Rhee RY (*see* Fernandez N et al). 2010;52:834-42
Rhee RY (*see* Wagner JK et al). 2010;52:216-8
Rho Y-N (*see* Park U-J et al). 2010;52:222
Ricco J-B (*see* Forbes TL et al). 2010;52:1716-22
Rich NM (*see* Hancock H et al). 2010;52:1405-9
Rich NM (*see* Welling DR et al). 2010;52:790-3
Rich NM, Krummel TM, Burris DG. Halsted-Holman vascular trauma legacy. 2010;52:508-11
Richardson AI (*see* Schmieder GC et al). 2010;52:375-82
Ricotta JJ. Invited Commentary. 2010;52:1450-8
Ricotta JJ, Malgor RD. Novel technique for endovascular salvage of a folded aortic endograft. 2010;52:785-9
Ridker PM (*see* Criqui MH et al). 2010;52:85-90
Rigberg D (*see* Lawrence PF et al). 2010;52:388-93
Rigberg DA (*see* Tulloch AW et al). 2010;52:1283-9
Righini P (*see* Trimarchi S et al). 2010;52:261-6
Riles T (*see* Fonseca R et al). 2010;52:556-61
Riles TS (*see* Kent KC et al). 2010;52:539-48
Riles TS, Lee V, Cheever D, Stableford J, Rockman CB. Clinical course of asymptomatic patients with carotid duplex scan end diastolic velocities of 100 to 124 centimeters per second. 2010;52:914-9
Rinaldi M (*see* Rispoli P et al). 2010;52:212-5
Rispoli P, Santovito D, Tallia C, Varetto G, Conforti M, Rinaldi M. A one-stage approach to the treatment of intravenous leiomyomatosis extending to the right heart. 2010;52:212-5
Ritti-Dias RM (*see* Gardner AW et al). 2010;52:1204-10
Rockman C (*see* Fonseca R et al). 2010;52:556-61
Rockman CB (*see* Riles TS et al). 2010;52:914-9
Roddy SP (*see* Mehta M et al). 2010;52:1442-9
Roddy SP (*see* Rahimi SA et al). 2010;52:891-6
Roddy SP. A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions. 2010;52:1417-8
Roddy SP. A real world analysis of payment per unit time in a Maryland Vascular Practice. 2010;52:1110
Roddy SP. Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia. 2010;52:794-5
Roddy SP. D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures. 2010;52:517
Roddy SP. Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections. 2010;52:516
Roddy SP. Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome. 2010;52:1419
Roddy SP. Limb ischemia during femoral cannulation for cardiopulmonary support. 2010;52:1109
Roddy SP. Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair. 2010;52:250
Roddy SP. Prosthetic lower extremity hemodialysis access grafts have satisfactory patency despite high incidence of infection. 2010;52:1723
Roddy SP. The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease. 2010;52:1724-5
Roddy SP. Trends in vena caval interruption. 2010;52:251
Roddy SP. A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the Closure-PLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein. 2010;52:796
Romagnuolo I (*see* Sticchi E et al). 2010;52:97-102
Ronziere T (*see* Mahe G et al). 2010;52:62-8
Rosa K (*see* Kashyap VS et al). 2010;52:369-74
Rosati MS (*see* Gabrielli R et al). 2010;52:464-6
Rosero EB (*see* Timaran CH et al). 2010;52:1188-95
Rossi E (*see* Bonforte G et al). 2010;52:1551-6
Rotondo MF (*see* Durham CA et al). 2010;52:884-90
Roy J (*see* Ekstrand J et al). 2010;52:167-75 (Basic research studies)
Rubin BG (*see* Jim J et al). 2010;52:730-3
Rubino P (*see* Stabile E et al). 2010;52:1505-10
Rutherford RB (*see* Buckley CJ et al). 2010;52:237-41 (Special communications)
Rutherford RB (*see* Vasquez MA et al). 2010;52:1387-96
Rzucidlo E (*see* Stone DH et al). 2010;52:127-31
Rzucidlo EM. Invited commentary. 2010;52:1290-300 (Basic research studies)

S

Saad WE (*see* Davies MG et al). 2010;52:331-9
Sabetai M (*see* Griffin MB et al). 2010;52:69-76
Sabetai M (*see* Nicolaides AN et al). 2010;52:1486-96

Sachs T, Pomposelli F, Hagberg R, Hamdan A, Wyers M, Giles K, Schermerhorn M. Open and endovascular repair of type B aortic dissection in the Nationwide Inpatient Sample. 2010;52: 860-6
Safranow K (*see* Wiernicki I et al). 2010;52:691-6
Sagara D (*see* Unno N et al). 2010;52:946-52
Salam A (*see* Chung J et al). 2010;52:1435-41
Salam AA (*see* Chung J et al). 2010;52:549-55
Sales CM, Haq F, Bustami R, Sun F. Management of isolated soleal and gastrocnemius vein thrombosis. 2010;52:1251-4
Sales CM, Schmidt RJ, FFBI Clinical Practice Workgroup. Fistula First clarifies its message. 2010;52:1699
Salloum A (*see* Vouyouka AG et al). 2010;52:1196-203
Sambol EB (*see* Goldstein LJ et al). 2010;52:1140-6
Sanada J (*see* Ohtake H et al). 2010;52:1464-70
Sanchez LA (*see* Jim J et al). 2010;52:730-3
Sannomiya P (*see* Erling N et al). 2010;52:1321-9
Santora RJ, Lie ML, Grigoryev DN, Nasir O, Moore FA, Hassoun HT. Therapeutic distant organ effects of regional hypothermia during mesenteric ischemia-reperfusion injury. 2010;52:1003-14
Santovito D (*see* Rispoli P et al). 2010;52:212-5
Sarac TP (*see* Kashyap VS et al). 2010;52:369-74
Sarfati MR (*see* Shelgikar C et al). 2010;52:920-4
Sarveswaran J (*see* Spark JI et al). 2010;52:632-6
Sassoust G (*see* Berard X et al). 2010;52:471-4
Savari P (*see* Sigala F et al). 2010;52:704-13
Scali ST (*see* Stone DH et al). 2010;52:127-31
Schönefeld E (*see* Donas KP et al). 2010;52:356-61
Schönleben F (*see* Bal A et al). 2010;52:456-9
Schanzer A (*see* Bertges DJ et al). 2010;52:674-83
Schanzer A (*see* De Martino RR et al). 2010;52:5-12
Schareck W (*see* Bünger CM et al). 2010;52:1357-61
Schellong S (*see* Schwarz T et al). 2010;52:1246-50
Schermerhorn M (*see* Sachs T et al). 2010;52:860-6
Schermerhorn ML (*see* Giles KA et al). 2010;52:1471-7
Schiller TB (*see* Martin JD et al). 2010;52:1094-100
Schillinger M (*see* Hoke M et al). 2010;52:383-7
Schmidt RJ (*see* Sales CM et al). 2010;52:1699
Schmieder GC, Carroll M, Panneton JM. Poor outcomes with cryoplasty for lower extremity arterial occlusive disease. 2010; 52:362-8
Schmieder GC, Richardson AI, Scott EC, Stokes GK, Meier GH, Panneton JM,. Outcomes of reinterventions after subintimal angioplasty. 2010;52:375-82
Schneider PA, Naylor AR. Asymptomatic carotid artery stenosis—Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting. 2010;52:499-507
Schumacher A (*see* Willenberg T et al). 2010;52:664-8
Schwarz T, Buschmann L, Beyer J, Halbritter K, Rastan A, Schellong S. Therapy of isolated calf muscle vein thrombosis: A randomized, controlled study. 2010;52:1246-50
Schwill S, LeMaire SA, Green SY, Bakaeen FG, Coselli JS. Endovascular repair of thoracic aortic pseudoaneurysms and patch aneurysms. 2010;52:1034-7 (Case rep.)
Schwindt A (*see* Donas KP et al). 2010;52:356-61
Scott DJA (*see* Parry DJ et al). 2010;52:145-51 (Basic research studies)
Scott EC (*see* Schmieder GC et al). 2010;52:375-82
Seal B (*see* Lopes LB et al). 2010;52:1596-607
Seike Y (*see* Nakamura Y et al). 2010;52:1061
Sesterhenn AM (*see* Wiegand S et al). 2010;52:742-5
Settmacher U (*see* Heise M et al). 2010;52:421-8 (Basic research studies)
Sevestre M-A (*see* Galanaud J-P et al). 2010;52:932-8
Shah PK (*see* Song L et al). 2010;52:994-1002
Shaik I (*see* Gupta V et al). 2010;52:1671-3
Sharifi BG (*see* Song L et al). 2010;52:994-1002
Shelgikar C, Mohebali J, Sarfati MR, Mueller MT, Kinikini DV, Kraiss LW. A design modification to minimize tilting of an inferior vena cava filter does not deliver a clinical benefit. 2010;52:920-4
Shemesh D, Goldin I, Olsha O. Banding between dialysis puncture sites to treat severe ischemic steal syndrome in low flow autogenous arteriovenous access. 2010;52:495-8
Shen J (*see* Chaer RA et al). 2010;52:569-75
Sheu WH-H (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Shiferson A (*see* Usoh F et al). 2010;52:394-9
Shirakawa Y (*see* Takahashi Y et al). 2010;52:1365-6
Shirasawa B (*see* Kubo M et al). 2010;52:429-34 (Basic research studies)
Shortell CK (*see* Vasquez MA et al). 2010;52:1387-96
Siani A (*see* Gabrielli R et al). 2010;52:464-6
Sigala F, Kotsinas A, Savari P, Filis K, Markantonis S, Iliodromitis EK, Gorgoulis VG, Andreadou I. Oxidized LDL in human carotid plaques is related to symptomatic carotid disease and lesion instability. 2010;52:704-13
Singh MJ (*see* Keefer A et al). 2010;52:873-7
Singh N (*see* Causey MW et al). 2010;52:1397-400
Singh N, Liu K, Tian L, Criqui MH, Guralnik JM, Ferrucci L, Liao Y, McDermott MM. Leg strength predicts mortality in men but not in women with peripheral arterial disease. 2010;52: 624-31
Smith KJ (*see* Spangler EL et al). 2010;52:
Smith L (*see* Mazari FAK et al). 2010;52:1564-71
Smith PC. Invited commentary. 2010;52:118-26
Smoke C (*see* Lopes LB et al). 2010;52:1596-607
Smolderen KG, Spertus JA, Vriens PW, Kranendonk S, Nooren M, Denollet J. Younger women with symptomatic peripheral arterial disease are at increased risk of depressive symptoms. 2010; 52:637-44
Society for Vascular Surgery Ad Hoc Committee on TEVAR Reporting Standards (*see* Fillinger MF et al). 2010;52:1022-33
Söderström M (*see* Arvela E et al). 2010;52:1218-25
Sofi F (*see* Sticchi E et al). 2010;52:97-102
Soga Y, Iida O, Hirano K, Yokoi H, Nanto S, Nobuyoshi M. Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent. 2010;52:608-15
Song L, Wang L, Shah PK, Chaux A, Sharifi BG. Bioengineered vascular graft grown in the mouse peritoneal cavity. 2010;52: 994-1002
Soong CV (*see* Jones C et al). 2010;52:298-302
Sorial E (*see* Endean ED et al). 2010;52:720-2 (Case rep.)
Sorropago G (*see* Stabile E et al). 2010;52:1505-10
Sosa JA (*see* Indes JE et al). 2010;52:1173-9
Spangler EL, Dillavou ED, Smith KJ. Cost-effectiveness of guidelines for insertion of inferior vena cava filters in high-risk trauma patients. 2010;52:
Spark JI, Sarveswaran J, Blest N, Charalabidis P, Asthana S. An elevated neutrophil-lymphocyte ratio independently predicts mortality in chronic critical limb ischemia. 2010;52:632-6
Speirs RD (*see* West CA et al). 2010;52:1164-72
Spertus JA (*see* Smolderen KG et al). 2010;52:637-44

Spinelli F (*see* Bitto A et al). 2010;52:1015-21
Sprengers RW, Teraa M, Moll FL, de Wit GA, van der Graaf Y, Verhaar MC, JUVENTAS and SMART study groups. Quality of life in patients with no-option critical limb ischemia underlines the need for new effective treatment. 2010;52:843-9
Squadrito F (*see* Bitto A et al). 2010;52:1015-21
Stabile E, Sorropago G, Tesorio T, Popusoi G, Ambrosini V, Mottola MT, Biamino G, Rubino P. Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: Results from a prospective randomized study. 2010;52:1505-10
Stableford J (*see* Riles TS et al). 2010;52:914-9
Stanziale SF (*see* Martin JD et al). 2010;52:1094-100
Stephen MS (*see* Geenen IL et al). 2010;52:1546-50
Stergiou N (*see* Koutakis P et al). 2010;52:340-7
Sternbach Y (*see* Mehta M et al). 2010;52:1442-9
Sternbach Y (*see* Rahimi SA et al). 2010;52:891-6
Sticchi E, Sofi F, Romagnuolo I, Pratesi G, Pulli R, Pratesi C, Abbate R, Fatini C. *eNOS* and *ACE* genes influence peripheral arterial disease predisposition in smokers. 2010;52:97-102
Stoica E (*see* Stoica AL et al). 2010;52:103-9
Stoica AL, Stoica E, Constantinescu I, Uscatescu V, Ginghina C. Interleukin-6 and interleukin-10 gene polymorphism, endothelial dysfunction, and postoperative prognosis in patients with peripheral arterial disease. 2010;52:103-9
Stokes GK (*see* Schmieder GC et al). 2010;52:375-82
Stone PA (*see* AbuRahma AF et al). 2010;52:1180-7
Stone WM (*see* Nakamura LY et al). 2010;52:453-5 (Case rep.)
Stone DH, Scali ST, Bjerk AA, Rzucidlo E, Chang CK, Goodney PP, Nolan BW, Walsh DB. Aggressive treatment of idiopathic axillo-subclavian vein thrombosis provides excellent long-term function. 2010;52:127-31
Stoner MC (*see* Durham CA et al). 2010;52:884-90
Stoner MC, deFreitas DJ. Process of care for carotid endarterectomy: Perioperative medical management. 2010;52:223-31 (Review article)
Strumpf R (*see* Hussain J et al). 2010;52:196-8 (Case rep.)
Su Q (*see* Jiang J et al). 2010;52:1278-82
Suárez DR (*see* Bosman W-MPF et al). 2010;52:152-8
Sueda T (*see* Mizukami T et al). 2010;52:1580-6
Sun F (*see* Sales CM et al). 2010;52:1251-4
Sunagawa K (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Sundt TM (*see* Pereda D et al). 2010;52:1362-4
Suritala (*see* Munezane T et al). 2010;52:967-74 (Basic research studies)
Sutherland J (*see* Hussain J et al). 2010;52:196-8 (Case rep.)
Suzuki M (*see* Unno N et al). 2010;52:946-52
Sweeting MJ, Thompson SG, Brown LC, Greenhalgh RM, Powell JT. Use of angiotensin converting enzyme inhibitors is associated with increased growth rate of abdominal aortic aneurysms. 2010;52:1-4
Szarvas T (*see* West CA et al). 2010;52:1164-72
Szeto WY (*see* Brinster CJ et al). 2010;52:460-3

T

Taggert JB (*see* Mehta M et al). 2010;52:1442-9
Taggert JB (*see* Rahimi SA et al). 2010;52:891-6
Tagusari O (*see* Nakamura Y et al). 2010;52:1061
Takagi H, Goto S, Matsui M, Manabe H, Umemoto T. A contemporary meta-analysis of Dacron versus polytetrafluoroethylene grafts for femoropopliteal bypass grafting. 2010;52:232-6
Takagi H, Goto S, Matsui M, Manabe H, Umemoto T. A further meta-analysis of population-based screening for abdominal aortic aneurysm. 2010;52:1103-8
Takagi H, Matsui M, Umemoto T. A meta-analysis of clinical studies of statins for prevention of abdominal aortic aneurysm expansion. 2010;52:1675-81 (Review article)
Takahashi H (*see* Ishii H et al). 2010;52:854-9
Takahashi S (*see* Mizukami T et al). 2010;52:1580-6
Takahashi Y, Tsutsumi Y, Shirakawa Y, Ohashi H. Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques. 2010;52:1365-6
Takemoto K (*see* Kawarada O et al). 2010;52:475-8
Tallia C (*see* Rispoli P et al). 2010;52:212-5
Tanaka A (*see* Munezane T et al). 2010;52:967-74 (Basic research studies)
Tanaka H (*see* Unno N et al). 2010;52:946-52
Tanaka H, Higashi M, Fukumoto Y, Ogino H. Entrapment of the popliteal artery. 2010;52:479
Taneja M (*see* Anil G et al). 2010;52:1038-40
Tangelder GJ (*see* Jongkind V et al). 2010;52:760-7 (Review article)
Tao M (*see* Yu P et al). 2010;52:444-52
Taoufiq H (*see* Berard X et al). 2010;52:471-4
Taylor LM. Invited commentary. 2010;52:1478-85
Teebken OE (*see* Bisdas T et al). 2010;52:323-30
Tefera G (*see* Bobadilla JL et al). 2010;52:267-71
Tegos T (*see* Nicolaides AN et al). 2010;52:1486-96
Teijink JAW (*see* Nicolaï SPA et al). 2010;52:1226-33
Teijink JAW (*see* Ten Bosch JA et al). 2010;52:13-8
ten Cate H (*see* ten Cate-Hoek AJ et al). 2010;52:132-8
Ten Bosch JA, Teijink JAW, Willigendael EM, Prins MH. Endovascular aneurysm repair is superior to open surgery for ruptured abdominal aortic aneurysms in EVAR-suitable patients. 2010;52:13-8
ten Cate-Hoek AJ, ten Cate H, Tordoir J, Hamulyák K, Prins MH. Individually tailored duration of elastic compression therapy in relation to incidence of the postthrombotic syndrome. 2010;52:132-8
Teraa M (*see* Sprengers RW et al). 2010;52:843-9
Tesorio T (*see* Stabile E et al). 2010;52:1505-10
Thakur S (*see* Comerota AJ et al). 2010;52:243-7
Thakur S, Comerota AJ. Bilateral nonthrombotic subclavian vein obstruction causing upper extremity venous claudication. 2010;52:208-11
Thalhammer C (*see* Willenberg T et al). 2010;52:664-8
Thelin S (*see* Mani K et al). 2010;52:1653-7 (Case rep.)
Theune B (*see* McQuade K et al). 2010;52:584-91
Thomas DJ (*see* Nicolaides AN et al). 2010;52:1486-96
Thompson A, Cooper JA, Fabricius M, Humphries SE, Ashton HA, Hafez H. An analysis of drug modulation of abdominal aortic aneurysm growth through 25 years of surveillance. 2010;52:55-61
Thompson MM (*see* Choke E et al). 2010;52:159-66
Thompson MM (*see* Malkawi AH et al). 2010;52:480-8 (Review article)
Thompson MM (*see* Patterson BO et al). 2010;52:25-30
Thompson SG (*see* Sweeting MJ et al). 2010;52:1-4
Tian L (*see* Singh N et al). 2010;52:624-31
Timaran CH, Rosero EB, Martinez AE, Ilarraza A, Modrall JG, Clagett GP. Atherosclerotic plaque composition assessed by virtual histology intravascular ultrasound and cerebral embolization after carotid stenting. 2010;52:1188-95
Tolenaar JL, van Keulen JW, Leijdekkers VJ, Vonken E-J, Moll FL, van Herwaarden JA. A ruptured aneurysm after stent graft

puncture during computed tomography-guided thrombin injection. 2010;52:1045-7
Tominaga R (*see* Oda S et al). 2010;52:412-20 (Basic research studies)
Tong MZ-Y, Koka P, Forbes TL. Economic evaluation of open vs endovascular repair of blunt traumatic thoracic aortic injuries. 2010;52:31-8
Tordoir J (*see* ten Cate-Hoek AJ et al). 2010;52:132-8
Toriyama T (*see* Ishii H et al). 2010;52:854-9
Torsello G (*see* Donas KP et al). 2010;52:356-61
Trellopoulos G, Georgiadis GS, C. Kapoulas K, Pitta X, Zervidis I, Lazarides MK. Emergency endovascular treatment of early spontaneous nonaneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia. 2010;52:751-7
Trimarchi S, Jonker FHW, Muhs BE, Grassi V, Righini P, Upchurch GR, Rampoldi V. Long-term outcomes of surgical aortic fenestration for complicated acute type B aortic dissections. 2010;52:261-6
TROMBOTEK Study Group (*see* Kurtoglu M et al). 2010;52:1262-71
Tsai K-L (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Tseng J-J (*see* Lee W-J et al). 2010;52:1290-300 (Basic research studies)
Tshomba Y (*see* Marone EM et al). 2010;52:1511-7
Tsutsumi Y (*see* Takahashi Y et al). 2010;52:1365-6
Tuggle CT (*see* Indes JE et al). 2010;52:1173-9
Tulloch AW, Jimenez JC, Lawrence PF, Dutson EP, Moore WS, Rigberg DA, Derubertis BG, Quinones-Baldrich WJ. Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome. 2010;52:1283-9

U

Uhl J-F, Gillot C, Chahim M. Anatomical variations of the femoral vein. 2010;52:714-9
Umemoto T (*see* Takagi H et al). 2010;52:1675-81 (Review article)
Unno N, Nishiyama M, Suzuki M, Tanaka H, Yamamoto N, Sagara D, Mano Y, Konno H. A novel method of measuring human lymphatic pumping using indocyanine green fluorescence lymphography. 2010;52:946-52
Upchurch GR (*see* Osborne NH et al). 2010;52:400-5
Upchurch GR (*see* Trimarchi S et al). 2010;52:261-6
Uscatescu V (*see* Stoica AL et al). 2010;52:103-9
Usoh F, Hingorani A, Ascher E, Shiferson A, Patel N, Gopal K, Marks N, Jacob T. Prospective randomized study comparing the clinical outcomes between inferior vena cava Greenfield and TrapEase filters. 2010;52:394-9

V

Valentine RJ (*see* Lang JL et al). 2010;52:290-7
Valstar ER (*see* Bosman W-MPF et al). 2010;52:152-8
van den Berg JC. Thrombolysis for acute arterial occlusion. 2010;52:512-5
van der Graaf Y (*see* Sprengers RW et al). 2010;52:843-9
van der Steenhoven TJ (*see* Bosman W-MPF et al). 2010;52:152-8
van der Vliet JA (*see* Geutjes PJ et al). 2010;52:1330-8
van Herwaarden JA (*see* Tolenaar JL et al). 2010;52:1045-7
van Herwaarden JA (*see* van Keulen JW et al). 2010;52:1081-6
van Keulen JW (*see* Tolenaar JL et al). 2010;52:1045-7
van Kuppevelt TH (*see* Geutjes PJ et al). 2010;52:1330-8
van Moerkerk HTB (*see* Geutjes PJ et al). 2010;52:1330-8
van Keulen JW, Moll FL, van Herwaarden JA. Tips and techniques for optimal stent graft placement in angulated aneurysm necks. 2010;52:1081-6
van Walraven C, Wong J, Morant K, Jennings A, Jetty P, Forster AJ. Incidence, follow-up, and outcomes of incidental abdominal aortic aneurysms. 2010;52:282-9
Vanicek NK (*see* Mazari FAK et al). 2010;52:1564-71
Varetto G (*see* Rispoli P et al). 2010;52:212-5
Vascular Study Group of New England (*see* Bertges DJ et al). 2010;52:674-83
Vascular Study Group of Northern New England (*see* De Martino RR et al). 2010;52:5-12
Vascular Study Group of Northern New England (*see* Goodney PP et al). 2010;52:897-905
Vasquez MA, Rabe E, McLafferty RB, Shortell CK, Marston WA, Gillespie D, Meissner MH, Rutherford RB. Revision of the venous clinical severity score: Venous outcomes consensus statement: Special communication of the American Venous Forum Ad Hoc Outcomes Working Group. 2010;52:1387-96
Veeraswamy RK (*see* Chung J et al). 2010;52:1435-41
Veith FJ (*see* Paraskevas KI et al). 2010;52:1367-75 (Review article)
Veith FJ, Powell JT, Hinchliffe RJ. Is a randomized trial necessary to determine whether endovascular repair is the preferred management strategy in patients with ruptured abdominal aortic aneurysms. 2010;52:1087-93
Velema E (*see* Derksen WJM et al). 2010;52:592-9
Venermo M (*see* Arvela E et al). 2010;52:1218-25
Verhaar MC (*see* Sprengers RW et al). 2010;52:843-9
Verhagen HJM (*see* Jonker FHW et al). 2010;52:39-44
Vermassen F (*see* Willaert W et al). 2010;52:1700-5
Verran DJ (*see* Geenen IL et al). 2010;52:1546-50
Verta P (*see* Matsumura JS et al). 2010;52:576-83
Vink A (*see* Derksen WJM et al). 2010;52:592-9
Vonken E-J (*see* Tolenaar JL et al). 2010;52:1045-7
Vos J-A (*see* Derksen WJM et al). 2010;52:592-9
Vouyouka AG, Egorova NN, Salloum A, Kleinman, L, Marin M, Faries PL, Moscowitz A. Lessons learned from the analysis of gender effect on risk factors and procedural outcomes of lower extremity arterial disease. 2010;52:1196-203
Vriens PW (*see* Smolderen KG et al). 2010;52:637-44

W

Wagner JK, Chaer RA, Rhee RY, Marone LK. True lumen re-entry after extravascular recanalization of a superficial femoral artery chronic total occlusion. 2010;52:216-8
Wakita T (*see* Katada Y et al). 2010;52:1343-5
Walker PJ (*see* Kruger AJ et al). 2010;52:1518-24
Walker AJ (*see* Hancock H et al). 2010;52:1405-9
Walkup MH (*see* Han DK et al). 2010;52:746-50
Walot I (*see* Khoynezhad A et al). 2010;52:562-8
Walsh DB (*see* Stone DH et al). 2010;52:127-31
Walsh ME (*see* Gale SS et al). 2010;52:645-50
Wang C (*see* Dong Z et al). 2010;52:1450-8
Wang GJ (*see* Foley PJ et al). 2010;52:850-3
Wang J (*see* Yao Y et al). 2010;52:1662-4
Wang L (*see* Song L et al). 2010;52:994-1002
Wang S (*see* Xiao Y et al). 2010;52:684-90 (Basic research studies)
Wang Y (*see* Dong Z et al). 2010;52:1450-8
Wang Z (*see* Jiang J et al). 2010;52:1278-82
Wanhainen A (*see* Mani K et al). 2010;52:1653-7 (Case rep.)
Warble PB (*see* Martin JD et al). 2010;52:1094-100
Wassel CL (*see* Criqui MH et al). 2010;52:85-90
Watanabe G (*see* Ohtake H et al). 2010;52:1464-70

Watkins MT (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Weiss LL (*see* Martin JD et al). 2010;52:1094-100
Welling DR, Burris DG, Rich NM. The influence of Dominique Jean Larrey on the art and science of amputations. 2010;52:790-3
Werebe EC (*see* Kuzniec S et al). 2010;52:1354-6
Werner JA (*see* Wiegand S et al). 2010;52:742-5
West CA, Johnson LW, Doucet L, Caldito G, Heldman M, Szarvas T, Speirs RD, Carson S. A contemporary experience of open aortic reconstruction in patients with chronic atherosclerotic occlusion of the abdominal aorta. 2010;52:1164-72
Wheatley G (*see* Hussain J et al). 2010;52:196-8 (Case rep.)
White GH (*see* Byrom MJ et al). 2010;52:176-95
White GH (*see* Geenen IL et al). 2010;52:1546-50
White RA (*see* Khoynezhad A et al). 2010;52:562-8
Wiegand S, Kureck I, Chapot R, Sesterhenn AM, Bien S, Werner JA. Early side effects after embolization of a carotid body tumor using Onyx. 2010;52:742-5
Wiernicki I, Safranow K, Baranowska-Bosiacka I, Piatek J, Gutowski P. Haptoglobin 2-1 phenotype predicts rapid growth of abdominal aortic aneurysms. 2010;52:691-6
Wiesbauer F (*see* Hoke M et al). 2010;52:383-7
Wilhelmi M (*see* Bisdas T et al). 2010;52:323-30
Willaert W, Aggarwal R, Bicknell C, Hamady M, Darzi A, Vermassen F, Cheshire N, European Virtual Reality Endovascular Research Team (EVEResT). Patient-specific simulation in carotid artery stenting. 2010;52:1700-5
Willenberg T, Schumacher A, Amann-Vesti B, Jacomella V, Thalhammer C, Diehm N, Baumgartner I, Husmann M. Impact of obesity on venous hemodynamics of the lower limbs. 2010;52:664-8
Williams K (*see* Burkhardt CGE et al). 2010;52:91-6
Willigendael EM (*see* Ten Bosch JA et al). 2010;52:13-8
Willis AK (*see* Perera AD et al). 2010;52:1339-42 (Case rep.)
Willson TD, Revesz E, Podbielski FJ, Blecha MJ. External iliac artery dissection secondary to endofibrosis in a cyclist. 2010;52:219-21
Wilson WRW (*see* Choke E et al). 2010;52:159-66
Wismans RG (*see* Geutjes PJ et al). 2010;52:1330-8
Wisselink W (*see* Jongkind V et al). 2010;52:1376-83
Wojnarowski DL (*see* Gale SS et al). 2010;52:645-50
Wolf BA (*see* Perera AD et al). 2010;52:1339-42 (Case rep.)
Wolosker N (*see* Kuzniec S et al). 2010;52:1354-6
Wong J (*see* van Walraven C et al). 2010;52:282-9
Woo EY (*see* Brinster CJ et al). 2010;52:460-3
Woo EY (*see* Foley PJ et al). 2010;52:850-3
Wu G (*see* Ding P-X et al). 2010;52:782-4
Wu L-S (*see* Hsiao J-F et al). 2010;52:669-73
Wu M (*see* Lawrence PF et al). 2010;52:388-93
Wyers M (*see* Sachs T et al). 2010;52:860-6
Wyers MC (*see* Giles KA et al). 2010;52:1471-7

X

Xiao Y, Huang Z, Yin H, Zhang H, Wang S. Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells. 2010;52:684-90 (Basic research studies)
Xu X (*see* Dong Z et al). 2010;52:1450-8
Xu Y (*see* Malkawi AH et al). 2010;52:480-8 (Review article)

Y

Yamamoto N (*see* Unno N et al). 2010;52:946-52
Yan Z (*see* Dong Z et al). 2010;52:1450-8
Yan Z-X (*see* Liu N-F et al). 2010;52:1557-63
Yang C-W (*see* Hsiao J-F et al). 2010;52:669-73
Yang K (*see* Hamblin M et al). 2010;52:984-93
Yao JST. The first operation on the profunda femoris artery. 2010;52:1101-2
Yao Y, Liu B, Wang J, Li H, Liang HD. Tuberculosis should not be ignored in patients with peripheral gangrene. 2010;52:1662-4
Yeung KK (*see* Jongkind V et al). 2010;52:1376-83
Yi Z (*see* Lopes LB et al). 2010;52:1596-607
Yin H (*see* Xiao Y et al). 2010;52:684-90 (Basic research studies)
Yokoi H (*see* Soga Y et al). 2010;52:608-15
Yokoi Y (*see* Kawarada O et al). 2010;52:475-8
Yoo H-J (*see* Kang J et al). 2010;52:435-43 (Basic research studies)
Yoshino M, Fukumoto H, Mizutani T, Yuyama R, Hara T. Mandibular subluxation stabilized by mouthpiece for distal internal carotid artery exposure in carotid endarterectomy. 2010;52:1401-4
Younes HK (*see* Davies MG et al). 2010;52:1478-85
Yu H (*see* Eton D et al). 2010;52:199-204
Yu P, Nguyen BT, Tao M, Campagna C, Ozaki CK. Rationale and practical techniques for mouse models of early vein graft adaptations. 2010;52:444-52
Yun W-S (*see* Park U-J et al). 2010;52:222
Yuyama R (*see* Yoshino M et al). 2010;52:1401-4

Z

Zanoni FL (*see* Erling N et al). 2010;52:1321-9
Zanow J (*see* Heise M et al). 2010;52:421-8 (Basic research studies)
Zen Y (*see* Kasashima S et al). 2010;52:1587-95 (Basic research studies)
Zervidis I (*see* Trellopoulos G et al). 2010;52:751-7
Zhai S (*see* Li T et al). 2010;52:1242-5
Zhang G (*see* Jiang J et al). 2010;52:1278-82
Zhang H (*see* Hamblin M et al). 2010;52:984-93
Zhang H (*see* Xiao Y et al). 2010;52:684-90 (Basic research studies)
Zhang J (*see* Hamblin M et al). 2010;52:984-93
Zhang WW (*see* Li T et al). 2010;52:1242-5
Zhao Y (*see* Bertges DJ et al). 2010;52:674-83
Zizka V (*see* Galanaud J-P et al). 2010;52:932-8
Zwolak RM (*see* Kent KC et al). 2010;52:539-48

Subject Index

A

Abdominal aortic aneurysm; *see* Aortic aneurysm, abdominal

Abstracts

Abstracts. 2010;52:252-6, 518-32, 797-817, 1111-23, 1420-4, 1726-47, S39-74

ACCH; *see* American College of Cardiology Foundation

Acetylsalicylic acid; *see* Aspirin

ACR; *see* American College of Radiology

Activities of daily living

Gender differences in daily ambulatory activity patterns in patients with intermittent claudication (Gardner et al). 2010;52:1204-10

Adhesives

Glue embolus complicating the endovascular treatment of a patient with Loeys-Dietz syndrome (Marine et al). 2010;52:1350-3 (Case rep.)

Adoption

Consensus statement on the adoption of the COPE guidelines. 2010;52:242

Aged

Cerebral reserve is decreased in elderly patients with carotid stenosis (Chaer et al). 2010;52:569-75

Aged, 80 and over

Endovascular aneurysm repair in nonagenarians is safe and effective (Goldstein et al); (Bush) (Commentary). 2010;52:1140-6

Intermediate-term EVAR outcomes in octogenarians (Fonseca et al). 2010;52:556-61

AHA; *see* American Heart Association

Algorithms

Classification of proximal endovenous closure levels and treatment algorithm (Lawrence et al). 2010;52:388-93

Alloys

Below knee bare nitinol stent placement in high-risk patients with critical limb ischemia is still durable after 24 months of follow-up (Donas et al). 2010;52:356-61

Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease (McQuade et al). 2010;52:584-91, 584.e1-e8

Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent (Soga et al). 2010;52:608-15

alpha 1-antitrypsin deficiency

Successful endovascular treatment of multiple mesenteric arterial aneurysms associated with alpha-1-antitrypsin deficiency (Gordon et al). 2010;52:467-70 (Case rep.)

American College of Cardiology Foundation

ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

American College of Radiology

ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

American Heart Association

ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

Amplatzer occluder device; *see* Septal occluder device

Amputation

History of the team approach to amputation prevention: pioneers and milestones (Sanders et al). 2010;52:S3-16

Hull early walking aid for rehabilitation of transtibial amputees - randomized controlled trial (HEART) (Mazari et al). 2010;52:1564-71

The influence of Dominique Jean Larrey on the art and science of amputations (Welling et al). 2010;52:790-3

Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010;52:1425-7

Rescuing Sisyphus: the team approach to amputation prevention (Mills Sr et al). 2010;52:S1-2 (Editorial)

Toe and flow: essential components and structure of the amputation prevention team (Rogers et al). 2010;52:S23-7

Anatomy

Regarding "The anatomy of the small saphenous vein: fascial and neural relations, saphenofemoral junction, and valves" (Caggiati) (Letter); (Schweighofer et al) (Reply). 2010;52:1428-9

Aneurysm

The effect of injectable biocompatible elastomer (PDMS) on the strength of the proximal fixation of endovascular aneurysm repair grafts: an in vitro study (Bosman et al). 2010;52:152-8

Endovascular aneurysm repair in nonagenarians is safe and effective (Goldstein et al); (Bush) (Commentary). 2010;52:1140-6

The impact of radiation dose exposure during endovascular aneurysm repair on patient safety (Jones et al). 2010;52:298-302

Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Abularrage et al). 2010;52:19-24

Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Roddy). 2010;52:250 (Editorial)

Primary tibial vein aneurysm with recurrent pulmonary emboli (Gabrielli et al). 2010;52:464-6 (Case rep.)

Regarding "Endovascular management of iliac rupture during endovascular aneurysm repair" (Semel et al). 2010;52:533 (Letter)

Significant sac retraction after endovascular aneurysm repair is a robust indicator of durable treatment success (Houbballah et al). 2010;52:878-83

Twelve-year outcomes after endovascular aneurysm repair using earliest available endograft components from a single center (Adiseshiah et al). 2010;52:49-54

Aneurysm, dissecting

Distribution of intimomedial tears in patients with type B aortic dissection (Khoynezhad et al). 2010;52:562-8

Endovascular management of type B aortic dissections (DiMusto et al). 2010;52:S26-36

Long-term outcomes of surgical aortic fenestration for complicated acute type B aortic dissections (Trimarchi et al). 2010;52:261-6

Open and endovascular repair of type B aortic dissection in the Nationwide Inpatient Sample (Sachs et al). 2010;52:860-6

Staged total exclusion of the aorta for chronic type B aortic dissection (Dias Perera et al). 2010;52:1339-42 (Case rep.)

Stent graft-induced new entry after endovascular repair for

Stanford type B aortic dissection (Dong et al); (Ricotta) (Commentary). 2010;52:1450-8

Aneurysm, false

Complex subclavian artery pseudoaneurysm causing failure of endovascular stent repair with salvage by percutaneous thrombin injection (Lee et al). 2010;52:1058-60 (Case rep.)

D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Hoke et al). 2010;52:383-7

D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Roddy). 2010;52:517 (Editorial)

Endovascular repair of thoracic aortic pseudoaneurysms and patch aneurysms (Schwill et al). 2010;52:1034-7 (Case rep.)

A giant femoral artery pseudoaneurysm in an infant boy (Park et al). 2010;52:222

Preparation and characterization of injectable fibrillar type I collagen and evaluation for pseudoaneurysm treatment in a pig model (Geutjes et al). 2010;52:1330-8

Regarding "Right subclavian pseudoaneurysm secondary to blunt trauma in an arteriovenous malformation" (Lee) (Letter);(Quinones-Baldrich,) (Reply). 2010;52:257-8

Transcatheter closure of recurrent aortic pseudoaneurysm previously treated by Amplatzer occluder device (Hussain et al). 2010;52:196-8 (Case rep.)

True lumen re-entry after extravascular recanalization of a superficial femoral artery chronic total occlusion (Wagner et al). 2010;52:216-18 (Case rep.)

Aneurysm, infected

Listeria monocytogenes as a rare cause of mycotic aortic aneurysm (Bal et al). 2010;52:456-9 (Case rep.)

Thoracic aortic endovascular repair for mycotic aneurysms and fistulas (Patel et al). 2010;52:S37-40 (Editorial)

Aneurysm repair; *see* Blood vessel prosthesis implantation

Aneurysm, ruptured

Emergency endovascular treatment of early spontaneous nonaneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia (Trellopoulos et al). 2010;52:751-7 (Case rep.)

Midterm outcome of endovascular repair of ruptured isolated iliac artery aneurysms (Hechelhammer et al). 2010;52:1159-63

Novel technique for endovascular salvage of a folded aortic endograft (Ricotta II and Malgor). 2010;52:785-9

Prior endovascular abdominal aortic aneurysm repair provides no survival benefits when the aneurysm ruptures (Cho et al). 2010;52:1127-34

Regarding "A closer look at meta-analyses of observational data" (Navarese et al) (Letter); (Jonker et al) (Reply). 2010;52:819-20

Regarding "Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I" (Mendonça). 2010;52:1425 (Letter)

A ruptured aneurysm after stent graft puncture during computed tomography-guided thrombin injection (Tolenaar et al). 2010;52:1045-7 (Case rep.)

Successful endovascular treatment of multiple mesenteric arterial aneurysms associated with alpha-1-antitrypsin deficiency (Gordon et al). 2010;52:467-70 (Case rep.)

Traumatic thoracic aortic injury and ruptures (Tefera). 2010;52:S41-4 (Editorial)

Angiogenesis inhibitors

Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20

Angiography

Intraprocedural imaging: flat panel detectors, rotational angiography, FluoroCT, IVUS, or still the portable C-arm? (Eagleton). 2010;52:S50-9 (Editorial)

The value of the initial post-EVAR computed tomography angiography scan in predicting future secondary procedures using the Powerlink stent graft (Patel and Carpenter). 2010;52:1135-9

Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction (Li et al). 2010;52:1242-5

Angioplasty

Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9

Cerebral reserve is decreased in elderly patients with carotid stenosis (Chaer et al). 2010;52:569-75

D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Hoke et al). 2010;52:383-7

D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Roddy). 2010;52:517 (Editorial)

Endoluminal treatment of type IA endoleak with Onyx (Grisafi et al). 2010;52:1346-9 (Case rep.)

Outcomes of interventions for recurrent disease after endoluminal intervention for superficial femoral artery disease (Davies et al). 2010;52:331-9, 331.e1-e2

Outcomes of reinterventions after subintimal angioplasty (Schmieder et al). 2010;52:375-82

Subintimal angioplasty for femoro-popliteal occlusive disease (Markose et al). 2010;52:1410-16

Angioplasty, balloon

Access site management with vascular closure devices for percutaneous transarterial procedures (Bechara et al). 2010;52:1682-96 (Review)

Primary balloon angioplasty plus balloon angioplasty maturation to upgrade small-caliber veins (<3 mm) for arteriovenous fistulas (De Marco Garcia et al). 2010;52:139-44

Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Lee). 2010;52:533 (Letter); Skurnick) (Letter); (Zamboni) (Reply). 2010;52:1749-50

Angiotensin-converting enzyme inhibitors

eNOS and *ACE* genes influence peripheral arterial disease predisposition in smokers (Sticchi et al). 2010;52:97-102, 97.e1

Use of angiotensin converting enzyme inhibitors is associated with increased growth rate of abdominal aortic aneurysms (Sweeting et al). 2010;52:1-4

Angiotensin II

Vascular endothelial growth factor enhances angiotensin II-induced aneurysm formation in apolipoprotein E-deficient mice (Choke et al). 2010;52:159-66, 159.e1

Anticoagulants

Therapy of isolated calf muscle vein thrombosis: a randomized, controlled study (Schwarz et al). 2010;52:1246-50

Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction (Li et al). 2010;52:1242-5

Antithrombins

A pilot, prospective evaluation of a direct thrombin inhibitor, bivalirudin (Angiomax), in patients undergoing lower extremity bypass (Kashyap et al). 2010;52:369-74

Aorta

Aorto-left renal vein fistula is a rare complication of abdominal aortic aneurysm with unique clinical presentation (Dragas et al). 2010;52:1658-61 (Case rep.)

A contemporary rural trauma center experience in blunt traumatic aortic injury (Durham et al). 2010;52:884-90

Dilated aortic root is related to a global aortic dilating diathesis (Alegret et al); (Eagleton) (Commentary). 2010;52:867-72

Endovascular procedures for aorto-iliac occlusive disease are associated with superior short-term clinical and economic outcomes compared with open surgery in the inpatient population (Indes et al). 2010;52:1173-9, 1173.e1

Fenestrated and branched devices in the pipeline (Greenberg and Qureshi). 2010;52:S15-21

Microcirculatory effects of local and remote ischemic preconditioning in supraceliac aortic clamping (Erling et al). 2010;52:1321-9

Persistence of intra-aortic air after circulatory arrest (Nakamura et al). 2010;52:1061

Seventeen-year follow-up after ascending-to-infrarenal aorta bypass for recurrent coarctation in an adult (Pereda et al). 2010;52:1362-4 (Case rep.)

A systematic review of endovascular treatment of extensive aortoiliac occlusive disease (Jongkind et al). 2010;52:1376-83 (Review)

Aorta, abdominal

A contemporary experience of open aortic reconstruction in patients with chronic atherosclerotic occlusion of the abdominal aorta (West et al). 2010;52:1164-72

Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Bisdas et al). 2010;52:323-30; Roddy). 2010;52:516 (Editorial)

Endovascular repair of extent I thoracoabdominal aneurysms with landing zone extension into the aortic arch and mesenteric portion of the abdominal aorta (Brinster et al). 2010;52:460-3 (Case rep.)

Outcomes in open repair of the thoracic and thoracoabdominal aorta (Acher and Wynn). 2010;52:S3-9

Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques (Takahashi et al). 2010;52:1365-6

Aorta, thoracic

Aortic dilatation after endovascular repair of blunt traumatic thoracic aortic injuries (Forbes et al). 2010;52:45-8

Aortic endograft sizing in trauma patients with hemodynamic instability (Jonker et al). 2010;52:39-44

Aortic rupture after spinal correction for scoliosis in the presence of a thoracic stent graft (Mani et al). 2010;52:1653-7 (Case rep.)

A clinicopathologic study of immunoglobulin G4-related sclerosing disease of the thoracic aorta (Kasashima et al). 2010;52:1587-95

Conduits and endoconduits, percutaneous access (Peterson). 2010;52:S60-4

Current Procedural Terminology (CPT) coding for endovascular intervention in the descending thoracic aorta (Seabrook). 2010;52:S103-6 (Editorial)

Economic evaluation of open vs endovascular repair of blunt traumatic thoracic aortic injuries (Tong et al). 2010;52:31-8

The effects of systemic hypothermia on a murine model of thoracic aortic ischemia reperfusion (Kang et al). 2010;52:435-43

Endovascular repair of extent I thoracoabdominal aneurysms with landing zone extension into the aortic arch and mesenteric portion of the abdominal aorta (Brinster et al). 2010;52:460-3 (Case rep.)

Endovascular repair of thoracic aortic traumatic transections is a safe method in patients with complicated injuries (Rahimi et al). 2010;52:891-6

The limitations of thoracic endovascular aortic repair in altering the natural history of blunt aortic injury (Lang et al). 2010;52:290-7

Outcomes in open repair of the thoracic and thoracoabdominal aorta (Acher and Wynn). 2010;52:S3-9

Risk factors for late mortality after endovascular repair of the thoracic aorta (Chung et al). 2010;52:549-55

In situ fenestration in the aortic arch (Manning et al). 2010;52:491-4

Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques (Takahashi et al). 2010;52:1365-6

Aortic aneurysm

Associated factors, timing, and technical aspects of late failure following open surgical aneurysm repairs (Coscas et al). 2010;52:272-81

Dilated aortic root is related to a global aortic dilating diathesis 4(Alegret et al); (Eagleton) (Commentary). 2010;52:867-72

Distribution of intimomedial tears in patients with type B aortic dissection (Khoynezhad et al). 2010;52:562-8

Endovascular management of type B aortic dissections (DiMusto et al). 2010;52:S26-36

Juxtarenal aortic aneurysm repair (Jongkind et al). 2010;52:760-7 (Review)

Listeria monocytogenes as a rare cause of mycotic aortic aneurysm (Bal et al). 2010;52:456-9 (Case rep.)

Long-term follow-up of neck expansion after endovascular aortic aneurysm repair (Monahan et al). 2010;52:303-7

Long-term outcomes of surgical aortic fenestration for complicated acute type B aortic dissections (Trimarchi et al). 2010;52:261-6

In situ fenestration in the aortic arch (Manning et al). 2010;52:491-4

Staged total exclusion of the aorta for chronic type B aortic dissection (Dias Perera et al). 2010;52:1339-42 (Case rep.)

Stent graft-induced new entry after endovascular repair for Stanford type B aortic dissection (Dong et al); (Ricotta) (Commentary). 2010;52:1450-8

Thrombotic mesenteric ischemia due to aortic dissection (Bao and Chen). 2010;52:1674, 1674.e1-e2

Aortic aneurysm, abdominal

Activation of transglutaminase type 2 for aortic wall protection in a rat abdominal aortic aneurysm formation (Munezane et al). 2010;52:967-74

Analysis of risk factors for abdominal aortic aneurysm in a cohort of more than 3 million individuals (Kent et al); (Lilly) (Commentary). 2010;52:539-48

An analysis of drug modulation of abdominal aortic aneurysm

growth through 25 years of surveillance (Thompson et al). 2010;52:55-61, 55.e1-e2

Aorto-left renal vein fistula is a rare complication of abdominal aortic aneurysm with unique clinical presentation (Dragas et al). 2010;52:1658-61 (Case rep.)

Duplex ultrasound factors predicting persistent type II endoleak and increasing AAA sac diameter after EVAR (Beeman et al). 2010;52:1147-52

The effect of warfarin therapy on endoleak development after endovascular aneurysm repair (EVAR) of the abdominal aorta (Bobadilla et al). 2010;52:267-71

Endovascular aneurysm repair for ruptured abdominal aortic aneurysm: how I do it (Mehta). 2010;52:1706-12

Endovascular aneurysm repair is superior to open surgery for ruptured abdominal aortic aneurysms in EVAR-suitable patients (Ten Bosch et al). 2010;52:13-18

Existing risk prediction methods for elective abdominal aortic aneurysm repair do not predict short-term outcome following endovascular repair (Patterson et al). 2010;52:25-30

A further meta-analysis of population-based screening for abdominal aortic aneurysm (Takagi et al). 2010;52:1103-8

Haptoglobin 2-1 phenotype predicts rapid growth of abdominal aortic aneurysms (Wiernicki et al). 2010;52:691-6

Hyperglycemia limits experimental aortic aneurysm progression (Miyama et al). 2010;52:975-83

I know about Jack and you're no Jack Kevorkian (Jones and McCullough). 2010;52:489-90

The impact of body mass index on perioperative outcomes of open and endovascular abdominal aortic aneurysm repair from the National Surgical Quality Improvement Program, 2005-2007 (Giles et al). 2010;52:1471-7

Incidence, follow-up, and outcomes of incidental abdominal aortic aneurysms (van Walraven et al). 2010;52:282-9, 282.e1-e2

The influence of aneurysm size on anatomic suitability for endovascular repair (Keefer et al). 2010;52:873-7

Intermediate-term EVAR outcomes in octogenarians (Fonseca et al). 2010;52:556-61

Is a randomized trial necessary to determine whether endovascular repair is the preferred management strategy in patients with ruptured abdominal aortic aneurysms? (Veith et al); (Forbes and Ricco) (Commentary). 2010;52:1087-93

Long-term outcomes of secondary procedures after endovascular aneurysm repair (Mehta et al). 2010;52:1442-9

Markers of inflammation in men with small abdominal aortic aneurysm (Parry et al). 2010;52:145-51

A meta-analysis of clinical studies of statins for prevention of abdominal aortic aneurysm expansion (Takagi et al). 2010;52:1675-81 (Review)

Midterm results of adjunctive neck therapies performed during elective infrarenal aortic aneurysm repair (Chung et al). 2010;52:1435-41

Novel technique for endovascular salvage of a folded aortic endograft (Ricotta II and Malgor). 2010;52:785-9

Outcomes of symptomatic abdominal aortic aneurysm repair (De Martino et al). 2010;52:5-12

Patient-specific biomechanical profiling in abdominal aortic aneurysm development and rupture (Malkawi et al). 2010;52:480-8 (Review)

Prior endovascular abdominal aortic aneurysm repair provides no survival benefits when the aneurysm ruptures (Cho et al). 2010;52:1127-34

Regarding "Long-term outcomes and resource utilization of endovascular versus open repair of abdominal aortic aneurysms in Ontario" (Takagi et al). 2010;52:818-19 (Letter)

Regarding "Preoperative statin therapy is associated with improved outcomes and resource utilization in patients undergoing aortic aneurysm repair" (Takagi et al) (Letter); (Stoner) (Reply). 2010;52:820-2

Regarding "Serum carboxymethyllysine concentrations are reduced in diabetic men with abdominal aortic aneurysms: Health In Men Study" (Meerwaldt and Zeebregts) (Letter); (Norman et al) (Reply). 2010;52:535-6

Regarding "The impact of model assumptions on results of computational mechanics in abdominal aortic aneurysm" (Georgakarakos et al) (Letter); (Reeps and Gee) (Reply). 2010;52:1124-5

Tips and techniques for optimal stent graft placement in angulated aneurysm necks (van Keulen et al). 2010;52:1081-6

Urgent interventional bilateral renal artery fenestration for giant pararenal abdominal aortic aneurysm with upper digestive tract obstruction (Ouyang and Liang). 2010;52:1048-51 (Case rep.)

Use of angiotensin converting enzyme inhibitors is associated with increased growth rate of abdominal aortic aneurysms (Sweeting et al). 2010;52:1-4

Use of a surgeon-modified branched thoracic endograft to preserve an aortorenal bypass during treatment of an intercostal patch aneurysm (Jim et al). 2010;52:730-3 (Case rep.)

The value of the initial post-EVAR computed tomography angiography scan in predicting future secondary procedures using the Powerlink stent graft (Patel and Carpenter). 2010;52:1135-9

Vascular endothelial growth factor enhances angiotensin II-induced aneurysm formation in apolipoprotein E-deficient mice (Choke et al). 2010;52:159-66, 159.e1

Vascular smooth muscle cell peroxisome proliferator-activated receptor-γ deletion promotes abdominal aortic aneurysms (Hamblin et al). 2010;52:984-93

Aortic aneurysm, thoracic

Advanced thoracic endovascular aortic repair: late complications of thoracic endografts (Halandras and Milner). 2010;52:S100-2 (Editorial)

Description and comparison of Food and Drug Administration-approved thoracic endovascular aneurysm devices (Lyden). 2010;52:S10-14

Endoleak management and postoperative surveillance following endovascular repair of thoracic aortic aneurysms (Ricotta). 2010;52:S91-9

Endovascular repair of extent I thoracoabdominal aneurysms with landing zone extension into the aortic arch and mesenteric portion of the abdominal aorta (Brinster et al). 2010;52:460-3 (Case rep.)

Endovascular repair of thoracic aortic pseudoaneurysms and patch aneurysms (Schwill et al). 2010;52:1034-7 (Case rep.)

First United States experience of the TX2 Pro-Form thoracic delivery system (Lee et al). 2010;52:1459-63

Incidence, prevention, and management in spinal cord protection during TEVAR (Rizvi and Sullivan). 2010;52:S86-90 (Editorial)

Open and endovascular repair of type B aortic dissection in the Nationwide Inpatient Sample (Sachs et al). 2010;52:860-6

Open techniques for arch vessel reconstruction during thoracic

endovascular aneurysm repair (TEVAR) (Vallabhaneni and Sanchez). 2010;52:S71-6 (Editorial)
Outcomes of planned celiac artery coverage during TEVAR (Mehta et al). 2010;52:1153-8
Regarding "A closer look at meta-analyses of observational data" (Navarese et al) (Letter); (Jonker et al) (Reply). 2010;52:819-20
Regulatory TEVAR clinical trials (Garcia-Toca and Eskandari). 2010;52:S22-5 (Editorial)
Reporting standards for thoracic endovascular aortic repair (TEVAR) (Fillinger et al). 2010;52:1022-33, 1022.e1-e5
Required skills and interdisciplinary teams in starting a TEVAR practice (Starr). 2010;52:S45-9 (Editorial)
Risk factor analysis of thoracic endovascular repair using the Matsui-Kitamura stent graft for acute aortic emergencies in the descending thoracic aorta (Ohtake et al). 2010;52:1464-70
Thoracic aortic endovascular repair for mycotic aneurysms and fistulas (Patel et al). 2010;52:S37-40 (Editorial)
Transcatheter closure of recurrent aortic pseudoaneurysm previously treated by Amplatzer occluder device (Hussain et al). 2010;52:196-8 (Case rep.)
Traumatic thoracic aortic injury and ruptures (Tefera). 2010;52:S41-4 (Editorial)
Use of a surgeon-modified branched thoracic endograft to preserve an aortorenal bypass during treatment of an intercostal patch aneurysm (Jim et al). 2010;52:730-3 (Case rep.)

Aortic arch; *see* Aorta, thoracic
Endovascular techniques for arch vessel reconstruction (Longo and Pipinos). 2010;52:S77-81
Open techniques for arch vessel reconstruction during thoracic endovascular aneurysm repair (TEVAR) (Vallabhaneni and Sanchez). 2010;52:S71-6 (Editorial)

Aortic coarctation
Seventeen-year follow-up after ascending-to-infrarenal aorta bypass for recurrent coarctation in an adult (Pereda et al). 2010;52:1362-4 (Case rep.)

Aortic diseases
Outcomes before and after initiation of an acute aortic treatment center (Davies et al); (Taylor) (Commentary). 2010;52:1478-85
Usefulness and limitations of computational models in aortic disease risk stratification (Beller et al). 2010;52:1572-9

Aortic rupture
Aortic rupture after spinal correction for scoliosis in the presence of a thoracic stent graft (Mani et al). 2010;52:1653-7 (Case rep.)
Endovascular aneurysm repair for ruptured abdominal aortic aneurysm: how I do it (Mehta). 2010;52:1706-12
Endovascular aneurysm repair is superior to open surgery for ruptured abdominal aortic aneurysms in EVAR-suitable patients (Ten Bosch et al). 2010;52:13-18
I know about Jack and you're no Jack Kevorkian (Jones and McCullough). 2010;52:489-90
Incidence, follow-up, and outcomes of incidental abdominal aortic aneurysms (van Walraven et al). 2010;52:282-9, 282.e1-e2
Is a randomized trial necessary to determine whether endovascular repair is the preferred management strategy in patients with ruptured abdominal aortic aneurysms? (Veith et al); (Forbes and Ricco) (Commentary). 2010;52:1087-93
The limitations of thoracic endovascular aortic repair in altering the natural history of blunt aortic injury (Lang et al). 2010;52:290-7
Patient-specific biomechanical profiling in abdominal aortic aneurysm development and rupture (Malkawi et al). 2010;52:480-8 (Review)
Stent graft-induced new entry after endovascular repair for Stanford type B aortic dissection (Dong et al); (Ricotta) (Commentary). 2010;52:1450-8

Apolipoproteins E
Vascular endothelial growth factor enhances angiotensin II-induced aneurysm formation in apolipoprotein E-deficient mice (Choke et al). 2010;52:159-66, 159.e1

Apoptosis
Evidence for markers of hypoxia and apoptosis in explanted human carotid atherosclerotic plaques (Bitto et al). 2010;52:1015-21
Inhibition of N-cadherin retards smooth muscle cell migration and intimal thickening via induction of apoptosis (Lyon et al). 2010;52:1301-9

Arm
Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Arvela et al). 2010;52:616-23
Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Roddy). 2010;52:794-5 (Editorial)
Inherent problems with randomized clinical trials with observational/no treatment arms (Buckley et al). 2010;52:237-41
The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease (Bonforte et al). 2010;52:1551-6
The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease (Roddy). 2010;52:1724-5

Arterial occlusive diseases
A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Boufi et al). 2010;52:1211-17; Roddy). 2010;52:1417-18 (Editorial)
A contemporary experience of open aortic reconstruction in patients with chronic atherosclerotic occlusion of the abdominal aorta (West et al). 2010;52:1164-72
Current assessment of endovascular therapy for infrainguinal arterial occlusive disease in patients with diabetes (Ihnat and Mills). 2010;52:S92-5
Endovascular procedures for aorto-iliac occlusive disease are associated with superior short-term clinical and economic outcomes compared with open surgery in the inpatient population (Indes et al). 2010;52:1173-9, 1173.e1
External iliac artery dissection secondary to endofibrosis in a cyclist (Willson et al). 2010;52:219-21 (Case rep.)
Limb salvage after delayed diagnosis for blunt traumatic infrapopliteal occlusion (Causey et al). 2010;52:734-7 (Case rep.)
Poor outcomes with cryoplasty for lower extremity arterial occlusive disease (Schmieder et al). 2010;52:362-8
Practical use of duplex echo-guided recanalization of chronic total occlusion in the iliac artery (Kawarada et al). 2010;52:475-8 (Case rep.)
Regarding "High spatial resolution magnetic resonance imaging of cystic adventitial disease of the popliteal artery"

(Weimar et al) (Letter); (Hagspiel) (Reply). 2010;52:534-5

Subintimal angioplasty for femoro-popliteal occlusive disease (Markose et al). 2010;52:1410-16

A systematic review of endovascular treatment of extensive aortoiliac occlusive disease (Jongkind et al). 2010;52:1376-83 (Review)

Thrombolysis for acute arterial occlusion (van den Berg). 2010;52:512-15

True lumen re-entry after extravascular recanalization of a superficial femoral artery chronic total occlusion (Wagner et al). 2010;52:216-18 (Case rep.)

Arteriovenous fistula

Fistula First clarifies its message (Sales et al). 2010;52:1699 (Editorial)

Liquid sclerotherapy for posttraumatic arteriovenous fistula of the radialis indicis artery (Katada et al). 2010;52:1343-5 (Case rep.)

Primary balloon angioplasty plus balloon angioplasty maturation to upgrade small-caliber veins (<3 mm) for arteriovenous fistulas (De Marco Garcia et al). 2010;52:139-44

Superficialization of arteriovenous fistulae employing minimally invasive liposuction (Causey et al). 2010;52:1397-400

Vascular changes at the puncture segments of arteriovenous fistula for hemodialysis access (Hsiao et al). 2010;52:669-73

Arteriovenous malformations

Regarding "Right subclavian pseudoaneurysm secondary to blunt trauma in an arteriovenous malformation" (Lee) (Letter); (Quinones-Baldrich,) (Reply). 2010;52:257-8

Arteriovenous shunt, surgical

Banding between dialysis puncture sites to treat severe ischemic steal syndrome in low flow autogenous arteriovenous access (Shemesh et al). 2010;52:495-8

History of temporary intravascular shunts in the management of vascular injury (Hancock et al). 2010;52:1405-9

The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease (Roddy). 2010;52:1724-5

In vitro testing of a newly developed arteriovenous double-outflow graft (Heise et al). 2010;52:421-8

Aspirin

Results of the randomized, placebo-controlled clopidogrel and acetylsalicylic acid in bypass surgery for peripheral arterial disease (CASPAR) trial (Belch et al). 2010;52:825-33, 825.e1-e2

Atherosclerosis

Regarding "Statins, heme oxygenase-1, iron, and atherosclerosis" (Sullivan) (Letter); (DePalma et al) (Reply). 2010;52:536-7

Atherosclerotic plaque; *see* Plaque, atherosclerotic

Athletic injuries

Bilateral internal carotid and vertebral artery dissection after a horse-riding injury (Keilani et al). 2010;52:1052-7 (Case rep.)

Axillary vein

Aggressive treatment of idiopathic axillo-subclavian vein thrombosis provides excellent long-term function (Stone et al). 2010;52:127-31

B

Bacteremia

Emergency endovascular treatment of early spontaneous non-aneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia (Trellopoulos et al). 2010;52:751-7 (Case rep.)

Balloon angioplasty; *see* Angioplasty, balloon

Bandages

Inelastic bandages maintain their hemodynamic effectiveness over time despite significant pressure loss (Mosti and Partsch). 2010;52:925-31

Behcet syndrome

Don't trust a vein graft to treat carotid aneurysm in patients with Behçet disease (Berard et al). 2010;52:471-4 (Case rep.)

Benchmarking

CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection (Matsumura et al). 2010;52:576-83, 576.e1-e2

Biocompatible materials

Animal models for the assessment of novel vascular conduits (Byrom et al). 2010;52:176-95

The effect of injectable biocompatible elastomer (PDMS) on the strength of the proximal fixation of endovascular aneurysm repair grafts: an in vitro study (Bosman et al). 2010;52:152-8

Bioengineering

Bioengineered vascular graft grown in the mouse peritoneal cavity (Song et al). 2010;52:994-1002, 994.e1-e2

Biological markers

Biomarkers in peripheral arterial disease patients and near- and longer-term mortality (Criqui et al). 2010;52:85-90

Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

Evidence for markers of hypoxia and apoptosis in explanted human carotid atherosclerotic plaques (Bitto et al). 2010;52:1015-21

Markers of inflammation in men with small abdominal aortic aneurysm (Parry et al). 2010;52:145-51

Biomarkers; *see* Biological markers

Biomechanics

Patient-specific biomechanical profiling in abdominal aortic aneurysm development and rupture (Malkawi et al). 2010;52:480-8 (Review)

Regarding "The impact of model assumptions on results of computational mechanics in abdominal aortic aneurysm" (Georgakarakos et al) (Letter); (Reeps and Gee) (Reply). 2010;52:1124-5

Bivalirudin; *see* Hirudins

Blood flow velocity

Clinical course of asymptomatic patients with carotid duplex scan end diastolic velocities of 100 to 124 centimeters per second (Riles et al). 2010;52:914-19, 914.e1

Blood platelets

Perioperative platelet and monocyte activation in patients with critical limb ischemia (Burdess et al). 2010;52:697-703

Blood vessel prosthesis

Fenestrated and branched devices in the pipeline (Greenberg and Qureshi). 2010;52:S15-21

Regulatory TEVAR clinical trials (Garcia-Toca and Eskandari). 2010;52:S22-5 (Editorial)

Blood vessel prosthesis implantation

Advanced thoracic endovascular aortic repair: late complications of thoracic endografts (Halandras and Milner). 2010; 52:S100-2 (Editorial)

Animal models for the assessment of novel vascular conduits (Byrom et al). 2010;52:176-95

Aortic endograft sizing in trauma patients with hemodynamic instability (Jonker et al). 2010;52:39-44

Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Arvela et al). 2010;52: 616-23

Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Roddy). 2010;52:794-5 (Editorial)

Bioengineered vascular graft grown in the mouse peritoneal cavity (Song et al). 2010;52:994-1002, 994.e1-e2

A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Boufi et al). 2010;52:1211-17

A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Roddy). 2010;52:1417-18 (Editorial)

A contemporary meta-analysis of Dacron versus polytetrafluoroethylene grafts for femoropopliteal bypass grafting (Takagi et al). 2010;52:232-6 (Review)

Description and comparison of Food and Drug Administration-approved thoracic endovascular aneurysm devices (Lyden). 2010;52:S10-14

Don't trust a vein graft to treat carotid aneurysm in patients with Behçet disease (Berard et al). 2010;52:471-4 (Case rep.)

Duplex ultrasound factors predicting persistent type II endoleak and increasing AAA sac diameter after EVAR (Beeman et al). 2010;52:1147-52

Early type III endoleak with an Endurant endograft (Abouliatim et al). 2010;52:1665-7 (Case rep.)

The effect of injectable biocompatible elastomer (PDMS) on the strength of the proximal fixation of endovascular aneurysm repair grafts: an in vitro study (Bosman et al). 2010; 52:152-8

The effect of warfarin therapy on endoleak development after endovascular aneurysm repair (EVAR) of the abdominal aorta (Bobadilla et al). 2010;52:267-71

Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Bisdas et al). 2010;52:323-30

Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Roddy). 2010;52:516 (Editorial)

Endoluminal treatment of type IA endoleak with Onyx (Grisafi et al). 2010;52:1346-9 (Case rep.)

Endovascular management of type B aortic dissections (DiMusto et al). 2010;52:S26-36

Endovascular repair of extent I thoracoabdominal aneurysms with landing zone extension into the aortic arch and mesenteric portion of the abdominal aorta (Brinster et al). 2010;52:460-3 (Case rep.)

Endovascular repair of thoracic aortic pseudoaneurysms and patch aneurysms (Schwill et al). 2010;52:1034-7 (Case rep.)

Endovascular repair of thoracic aortic traumatic transections is a safe method in patients with complicated injuries (Rahimi et al). 2010;52:891-6

Endovascular techniques for arch vessel reconstruction (Longo and Pipinos). 2010;52:S77-81

First United States experience of the TX2 Pro-Form thoracic delivery system (Lee et al). 2010;52:1459-63

Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease (McQuade et al). 2010; 52:584-91, 584.e1-e8

Glue embolus complicating the endovascular treatment of a patient with Loeys-Dietz syndrome (Marine et al). 2010; 52:1350-3 (Case rep.)

Intermediate-term EVAR outcomes in octogenarians (Fonseca et al). 2010;52:556-61

Mechanisms to explain the poor results of carotid artery stenting (CAS) in symptomatic patients to date and options to improve CAS outcomes (Paraskevas et al). 2010;52:1367-75 (Review)

Midterm results of adjunctive neck therapies performed during elective infrarenal aortic aneurysm repair (Chung et al). 2010;52:1435-41

Novel technique for endovascular salvage of a folded aortic endograft (Ricotta II and Malgor). 2010;52:785-9

Open and endovascular repair of type B aortic dissection in the Nationwide Inpatient Sample (Sachs et al). 2010;52:860-6

Open techniques for arch vessel reconstruction during thoracic endovascular aneurysm repair (TEVAR) (Vallabhaneni and Sanchez). 2010;52:S71-6 (Editorial)

Outcomes of planned celiac artery coverage during TEVAR (Mehta et al). 2010;52:1153-8

Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Abularrage et al). 2010;52: 19-24

Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Roddy). 2010;52:250 (Editorial)

Prosthetic lower extremity haemodialysis access grafts have satisfactory patency despite high incidence of infection (Roddy). 2010;52:1723 (Editorial)

Prosthetic lower extremity hemodialysis access grafts have satisfactory patency despite a high incidence of infection (Geenen et al). 2010;52:1546-50

Rationale and practical techniques for mouse models of early vein graft adaptations (Yu et al). 2010;52:444-52

Regarding "Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I" (Mendonça). 2010;52:1425 (Letter)

Results of the randomized, placebo-controlled clopidogrel and acetylsalicylic acid in bypass surgery for peripheral arterial disease (CASPAR) trial (Belch et al). 2010;52:825-33, 825.e1-e2

Risk factor analysis of thoracic endovascular repair using the Matsui-Kitamura stent graft for acute aortic emergencies in the descending thoracic aorta (Ohtake et al). 2010;52:1464-70

Significant sac retraction after endovascular aneurysm repair is a robust indicator of durable treatment success (Houbballah et al). 2010;52:878-83

In situ fenestration in the aortic arch (Manning et al). 2010; 52:491-4

Stent graft-induced new entry after endovascular repair for

Stanford type B aortic dissection (Dong et al); (Ricotta) (Commentary). 2010;52:1450-8
Tips and techniques for optimal stent graft placement in angulated aneurysm necks (van Keulen et al). 2010;52:1081-6
Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques (Takahashi et al). 2010;52:1365-6
True lumen re-entry after extravascular recanalization of a superficial femoral artery chronic total occlusion (Wagner et al). 2010;52:216-18 (Case rep.)
Twelve-year outcomes after endovascular aneurysm repair using earliest available endograft components from a single center (Adiseshiah et al). 2010;52:49-54
Urgent interventional bilateral renal artery fenestration for giant pararenal abdominal aortic aneurysm with upper digestive tract obstruction (Ouyang and Liang). 2010;52:1048-51 (Case rep.)
Use of a surgeon-modified branched thoracic endograft to preserve an aortorenal bypass during treatment of an intercostal patch aneurysm (Jim et al). 2010;52:730-3 (Case rep.)
The value of the initial post-EVAR computed tomography angiography scan in predicting future secondary procedures using the Powerlink stent graft (Patel and Carpenter). 2010;52:1135-9
Visceral reconstruction techniques (Srivastava). 2010;52:S82-5 (Editorial)
In vitro testing of a newly developed arteriovenous double-outflow graft (Heise et al). 2010;52:421-8

Blunt injuries; *see* Wounds, nonpenetrating

Body mass index
The impact of body mass index on perioperative outcomes of open and endovascular abdominal aortic aneurysm repair from the National Surgical Quality Improvement Program, 2005-2007 (Giles et al). 2010;52:1471-7

Bone marrow
Upper extremity ischemia treated with tissue repair cells from adult bone marrow (Comerota et al). 2010;52:723-9 (Case rep.)

Brachiocephalic veins
Collateral pathways visualization of the innominate vein (Civilini et al). 2010;52:758-9
Revascularization of an occluded brachiocephalic vein using Outback-LTD re-entry catheter (Anil and Taneja). 2010;52:1038-40 (Case rep.)

Brain ischemia
A durable iliac-axillary and axillary-carotid bypass for cerebral ischemia due to Takayasu arteritis allowed successful pregnancies and deliveries (Kondo et al). 2010;52:1713-5

Budd-Chiari syndrome
Agitation thrombolysis for fresh iatrogenic IVC thrombosis in patients with Budd-Chiari syndrome (Ding et al). 2010;52:782-4
Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction (Li et al). 2010;52:1242-5

C

Cadherins
Inhibition of N-cadherin retards smooth muscle cell migration and intimal thickening via induction of apoptosis (Lyon et al). 2010;52:1301-9

Cannulation; *see* Catheterization

Cardiopulmonary resuscitation
Limb ischemia during femoral cannulation for cardiopulmonary support (Foley et al). 2010;52:850-3
Limb ischemia during femoral cannulation for cardiopulmonary support (Roddy). 2010;52:1109 (Editorial)

Cardiopulmonary support; *see* Cardiopulmonary resuscitation

Cardiovascular diseases
Combined effects of smoking and peripheral arterial disease on all-cause and cardiovascular disease mortality in a Chinese male cohort (Luo et al). 2010;51:673-8. Correction. 2010;52:260

Carotid arteries
Atherosclerotic plaque composition assessed by virtual histology intravascular ultrasound and cerebral embolization after carotid stenting (Timaran et al). 2010;52:1188-95
CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection (Matsumura et al). 2010;52:576-83, 576.e1-e2
Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis (AbuRahma et al). 2010;52:1180-7
A durable iliac-axillary and axillary-carotid bypass for cerebral ischemia due to Takayasu arteritis allowed successful pregnancies and deliveries (Kondo et al). 2010;52:1713-5
Eight-year experience with carotid artery stenting for correction of symptomatic and asymptomatic post-endarterectomy defects). 2010;52:1511-7
Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: results from a prospective randomized study (Stabile et al). 2010;52:1505-10
Mandibular subluxation stabilized by mouthpiece for distal internal carotid artery exposure in carotid endarterectomy (Yoshino et al). 2010;52:1401-4
Mechanisms to explain the poor results of carotid artery stenting (CAS) in symptomatic patients to date and options to improve CAS outcomes (Paraskevas et al). 2010;52:1367-75 (Review)
Outcomes of carotid stenting compared with endarterectomy are equivalent in asymptomatic patients and inferior in symptomatic patients (Giacovelli et al). 2010;52:906-13, 906.e1-e4
Patient-specific simulation in carotid artery stenting (Willaert et al). 2010;52:1700-5
Stroke and death after carotid endarterectomy and carotid artery stenting with and without high risk criteria (Giles et al). 2010;52:1497-504

Carotid artery diseases
An unfavorable dietary pattern is associated with symptomatic ischemic stroke and carotid atherosclerosis (Mahe et al). 2010;52:62-8
Catheter-based neurosalvage for acute embolic complication during carotid intervention (Lin et al). 2010;52:308-13
A contemporary experience of open aortic reconstruction in patients with chronic atherosclerotic occlusion of the abdominal aorta (West et al). 2010;52:1164-72
Don't trust a vein graft to treat carotid aneurysm in patients with Behçet disease (Berard et al). 2010;52:471-4 (Case rep.)
Evidence for markers of hypoxia and apoptosis in explanted human carotid atherosclerotic plaques (Bitto et al). 2010;52:1015-21

Is carotid artery disease responsible for perioperative strokes after coronary artery bypass surgery? (Li et al); (Forbes and Ricco) (Commentary). 2010;52:1716-22

Oxidized LDL in human carotid plaques is related to symptomatic carotid disease and lesion instability (Sigala et al). 2010;52:704-13

Carotid artery, internal

Asymptomatic internal carotid artery stenosis and cerebrovascular risk stratification (Nicolaides et al). 2010;52:1486-96, 1486.e1-e5

Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al). 2010;52:69-76

Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al) 2010;52:69-76. Correction. 2010;52:1430

Carotid artery, internal, dissection

Bilateral internal carotid and vertebral artery dissection after a horse-riding injury (Keilani et al). 2010;52:1052-7 (Case rep.)

Carotid atherosclerosis; *see* Carotid artery diseases

Carotid body tumor

Carotid body tumor presenting with carotid sinus syndrome (da Gama and Cabral). 2010;52:1668-70 (Case rep.)

Early side effects after embolization of a carotid body tumor using Onyx (Wiegand et al). 2010;52:742-5 (Case rep.)

Important observations made managing carotid body tumors during a 25-year experience (Kruger et al); (LaMuraglia) (Commentary). 2010;52:1518-24

A rare case of familial carotid body tumor in a patient with bilateral fibromuscular dysplasia (Han et al). 2010;52:746-50 (Case rep.)

Carotid endarterectomy; *see* Endarterectomy, carotid

Carotid plaque; *see* Carotid stenosis

Carotid sinus

Carotid body tumor presenting with carotid sinus syndrome (da Gama and Cabral). 2010;52:1668-70 (Case rep.)

Carotid stenosis

Asymptomatic carotid artery stenosis-Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting (Schneider and Naylor); (Forbes) (Commentary) 2010;52:499-507. Correction. 2010;52:823

Asymptomatic carotid artery stenosis—Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting (Schneider and Naylor); (Forbes) (Commentary). 2010;52:499-507

Asymptomatic internal carotid artery stenosis and cerebrovascular risk stratification (Nicolaides et al). 2010;52:1486-96, 1486.e1-e5

Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis (AbuRahma et al). 2010;52:1180-7

Cerebral reserve is decreased in elderly patients with carotid stenosis (Chaer et al). 2010;52:569-75

Clinical course of asymptomatic patients with carotid duplex scan end diastolic velocities of 100 to 124 centimeters per second (Riles et al). 2010;52:914-19, 914.e1

Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al). 2010;52:69-76

Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al) 2010;52:69-76. Correction. 2010;52:1430

Mechanisms to explain the poor results of carotid artery stenting (CAS) in symptomatic patients to date and options to improve CAS outcomes (Paraskevas et al). 2010;52:1367-75 (Review)

Oxidized LDL in human carotid plaques is related to symptomatic carotid disease and lesion instability (Sigala et al). 2010;52:704-13

Patient-specific simulation in carotid artery stenting (Willaert et al). 2010;52:1700-5

Case reports

Case reports. 2010;52:196-221, 453-78, 720-57, 1034-60, 1339-64, 1653-73

Catheter ablation

A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Gale et al). 2010;52:645-50

A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Roddy). 2010;52:796 (Editorial)

Catheterization

Catheter-based neurosalvage for acute embolic complication during carotid intervention (Lin et al). 2010;52:308-13

Limb ischemia during femoral cannulation for cardiopulmonary support (Foley et al). 2010;52:850-3

Limb ischemia during femoral cannulation for cardiopulmonary support (Roddy). 2010;52:1109 (Editorial)

Transcatheter closure of recurrent aortic pseudoaneurysm previously treated by Amplatzer occluder device (Hussain et al). 2010;52:196-8 (Case rep.)

Videothoracoscopic-guided management of a central vein perforation during hemodialysis catheter placement (Kuzniec et al). 2010;52:1354-6 (Case rep.)

Catheterization, central venous

Revascularization of an occluded brachiocephalic vein using Outback-LTD re-entry catheter (Anil and Taneja). 2010;52:1038-40 (Case rep.)

Celiac artery

Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Roddy). 2010;52:1419 (Editorial)

Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Tulloch et al). 2010;52:1283-9

Microcirculatory effects of local and remote ischemic preconditioning in supraceliac aortic clamping (Erling et al). 2010;52:1321-9

Outcomes of planned celiac artery coverage during TEVAR (Mehta et al). 2010;52:1153-8

Cell differentiation

Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

Cell migration; *see* Cell movement

Cell movement

Inhibition of N-cadherin retards smooth muscle cell migration and intimal thickening via induction of apoptosis (Lyon et al). 2010;52:1301-9

Cell proliferation
Tissue factor pathway inhibitor-2 is induced by fluid shear stress in vascular smooth muscle cells and affects cell proliferation and survival (Ekstrand et al). 2010;52:167-75
Cell survival
Tissue factor pathway inhibitor-2 is induced by fluid shear stress in vascular smooth muscle cells and affects cell proliferation and survival (Ekstrand et al). 2010;52:167-75
Cerebral embolization; *see* Intracranial embolism
Cerebral ischemia; *see* Brain ischemia
Cerebrospinal fluid (CSF)
Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9
Cerebrovascular circulation
Cerebral reserve is decreased in elderly patients with carotid stenosis (Chaer et al). 2010;52:569-75
Cerebrovascular disorders
Asymptomatic internal carotid artery stenosis and cerebrovascular risk stratification (Nicolaides et al). 2010;52:1486-96, 1486.e1-e5
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Lee). 2010;52:533 (Letter)
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Skurnick) (Letter); (Zamboni) (Reply). 2010;52:1749-50
Chronic disease
A contemporary experience of open aortic reconstruction in patients with chronic atherosclerotic occlusion of the abdominal aorta (West et al). 2010;52:1164-72
Prevention and treatment of venous ulcers in primary chronic venous insufficiency (Neglen). 2010;52:S15-20
Reducing venous stasis ulcers by fifty percent in 10 years: the next steps (Henke et al). 2010;52:S37-8
Circulatory arrest, deep hypothermia induced
Persistence of intra-aortic air after circulatory arrest (Nakamura et al). 2010;52:1061
Clamping; *see* Constriction
Clinical trials as topic
Regulatory TEVAR clinical trials (Garcia-Toca and Eskandari). 2010;52:S22-5 (Editorial)
Clopidogrel; *see* Ticlopidine
Collateral circulation
Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20
Compression bandages
Individually tailored duration of elastic compression therapy in relation to incidence of the postthrombotic syndrome (ten Cate-Hoek et al). 2010;52:132-8
Computed tomography; *see* Tomography, X-ray computed
Consensus
Consensus statement on the adoption of the COPE guidelines. 2010;52:242
Revision of the venous clinical severity score: venous outcomes consensus statement: special communication of the American Venous Forum Ad Hoc Outcomes Working Group (Vasquez et al). 2010;52:1387-96
Constriction
Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: results from a prospective randomized study (Stabile et al). 2010;52:1505-10
Constriction, pathologic
Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9
Coronary artery bypass
Is carotid artery disease responsible for perioperative strokes after coronary artery bypass surgery? (Li et al); (Forbes and Ricco) (Commentary). 2010;52:1716-22
Coronary restenosis
Restenosis after carotid endarterectomy in a multicenter regional registry (Goodney et al). 2010;52:897-905, 897.e1-e2
Corrections
Asymptomatic carotid artery stenosis-Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting (Schneider and Naylor); (Forbes) (Commentary) 2010;52:499-507). 2010;52:823
Combined effects of smoking and peripheral arterial disease on all-cause and cardiovascular disease mortality in a Chinese male cohort (Luo et al). 2010;51:673-8. Correction. 2010; 52:260
Elevated sex steroid hormones in great saphenous veins in men (Kendler et al); (Labropoulos) (Commentary). 2010;51: 639-47). 2010;52:538
Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al). 2010;52: 69-76. Correction. 2010;52:1430
Cost-benefit analysis
Cost-effectiveness of guidelines for insertion of inferior vena cava filters in high-risk trauma patients (Spangler et al). 2010;52:1537-45, 1537.e1-e2
C-reactive protein
Prognostic values of C-reactive protein levels on clinical outcome after endovascular therapy in hemodialysis patients with peripheral artery disease (Ishii et al). 2010;52:854-9
Cryopreservation
Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Bisdas et al). 2010;52:323-30
Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Roddy). 2010;52:516 (Editorial)
Cryotherapy
Poor outcomes with cryoplasty for lower extremity arterial occlusive disease (Schmieder et al). 2010;52:362-8
Current procedural terminology
Current Procedural Terminology (CPT) coding for endovascular intervention in the descending thoracic aorta (Seabrook). 2010;52:S103-6 (Editorial)
Cysts
Regarding "High spatial resolution magnetic resonance imaging of cystic adventitial disease of the popliteal artery" (Weimar et al) (Letter); (Hagspiel) (Reply). 2010;52:534-5

D

Dacron; *see* Polyethylene terephthalates
Databases as topic
Analysis of large databases in vascular surgery (Nguyen and Barshes). 2010;52:768-74 (Review)
D-dimer; *see* Fibrin fibrinogen degradation products

Death
Stroke and death after carotid endarterectomy and carotid artery stenting with and without high risk criteria (Giles et al). 2010;52:1497-504
Debridement
Débridement of the noninfected wound (Cornell et al). 2010; 52:S31-6
A step-wise approach for surgical management of diabetic foot infections (Fisher et al). 2010;52:S72-5
Decompression, surgical
The spectrum of arterial compression at the thoracic outlet (Criado et al). 2010;52:406-11
Surgical off-loading of the diabetic foot (Frykberg et al). 2010; 52:S44-58
Deep vein thrombosis; *see* Venous thrombosis
Delayed diagnosis
Limb salvage after delayed diagnosis for blunt traumatic infrapopliteal occlusion (Causey et al). 2010;52:734-7 (Case rep.)
Depression
Younger women with symptomatic peripheral arterial disease are at increased risk of depressive symptoms (Smolderen et al). 2010;52:637-44
Desmuslin; *see* Intermediate filament proteins
Diabetes mellitus
Arterial imaging in patients with lower extremity ischemia and diabetes mellitus (Pomposelli). 2010;52:S81-91
Challenges of distal bypass surgery in patients with diabetes: patient selection, techniques, and outcomes (Conte). 2010; 52:S96-103
Current assessment of endovascular therapy for infrainguinal arterial occlusive disease in patients with diabetes (Ihnat and Mills). 2010;52:S92-5
Long-term outcomes of diabetic patients undergoing endovascular infrainguinal interventions (Abularrage et al). 2010; 52:314-22, 314.e1-e4
Noninvasive assessment of lower extremity hemodynamics in individuals with diabetes mellitus (Andersen). 2010;52:S76-80
Regarding "Serum carboxymethyllysine concentrations are reduced in diabetic men with abdominal aortic aneurysms: Health In Men Study" (Meerwaldt and Zeebregts) (Letter); (Norman et al) (Reply). 2010;52:535-6
What you can't feel can hurt you (Boulton). 2010;52:S28-30
Diabetic foot
The costs of diabetic foot: the economic case for the limb salvage team (Driver et al). 2010;52:S17-22
History of the team approach to amputation prevention: pioneers and milestones (Sanders et al). 2010;52:S3-16
Medical therapy of diabetic foot infections (Joseph and Lipsky). 2010;52:S67-71
Off-loading the diabetic foot for ulcer prevention and healing (Cavanagh and Bus). 2010;52:S37-43
Rescuing Sisyphus: the team approach to amputation prevention (Mills Sr et al). 2010;52:S1-2 (Editorial)
A step-wise approach for surgical management of diabetic foot infections (Fisher et al). 2010;52:S72-5
Surgical off-loading of the diabetic foot (Frykberg et al). 2010; 52:S44-58
What you can't feel can hurt you (Boulton). 2010;52:S28-30
Diabetic neuropathies
What you can't feel can hurt you (Boulton). 2010;52:S28-30
Diagnostic imaging
Arterial imaging in patients with lower extremity ischemia and diabetes mellitus (Pomposelli). 2010;52:S81-91
Dialysis
Banding between dialysis puncture sites to treat severe ischemic steal syndrome in low flow autogenous arteriovenous access (Shemesh et al). 2010;52:495-8
Diastole
Clinical course of asymptomatic patients with carotid duplex scan end diastolic velocities of 100 to 124 centimeters per second (Riles et al). 2010;52:914-19, 914.e1
Diathesis; *see* Disease susceptibility
Diet
An unfavorable dietary pattern is associated with symptomatic ischemic stroke and carotid atherosclerosis (Mahe et al). 2010;52:62-8
Dilatation, pathologic
Aortic dilatation after endovascular repair of blunt traumatic thoracic aortic injuries (Forbes et al). 2010;52:45-8
Dilated aortic root is related to a global aortic dilating diathesis (Alegret et al); (Eagleton) (Commentary). 2010;52:867-72
Dimethyl sulfoxide
Endoluminal treatment of type IA endoleak with Onyx (Grisafi et al). 2010;52:1346-9 (Case rep.)
Direct thrombin inhibitor; *see* Antithrombins
Disease progression
Hyperglycemia limits experimental aortic aneurysm progression (Miyama et al). 2010;52:975-83
Management of isolated soleal and gastrocnemius vein thrombosis (Sales et al). 2010;52:1251-4
Disease susceptibility
Dilated aortic root is related to a global aortic dilating diathesis (Alegret et al); (Eagleton) (Commentary). 2010;52:867-72
Dissection
External iliac artery dissection secondary to endofibrosis in a cyclist (Willson et al). 2010;52:219-21 (Case rep.)
Diverticulum
Iliofemoral venous thrombosis from external compression by a vesical diverticulum (Gupta et al). 2010;52:1671-3 (Case rep.)
Do not resuscitate orders; *see* Resuscitation orders
Drug prescriptions
An analysis of drug modulation of abdominal aortic aneurysm growth through 25 years of surveillance (Thompson et al). 2010;52:55-61, 55.e1-e2
Duodenum
Late erosion of a prophylactic Celect IVC filter into the aorta, right renal artery, and duodenal wall (Becher et al). 2010; 52:1041-4 (Case rep.)

E

Earthquakes
A fellow's experience in Haiti (Lind). 2010;52:533-4 (Letter)
Echocardiography
Dilated aortic root is related to a global aortic dilating diathesis (Alegret et al); (Eagleton) (Commentary). 2010;52:867-72
Editorials
Acknowledgment of reviewers (Sidawy and Perler). 2010;52: 1431-4
Advanced thoracic endovascular aortic repair: late complica-

tions of thoracic endografts (Halandras and Milner). 2010; 52:S100-2
Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Roddy). 2010;52:794-5
A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Roddy). 2010;52:1417-18
Current Procedural Terminology (CPT) coding for endovascular intervention in the descending thoracic aorta (Seabrook). 2010;52:S103-6
D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Roddy). 2010;52:517
Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Roddy). 2010;52:516
Fistula First clarifies its message (Sales et al). 2010;52:1699
Incidence, prevention, and management in spinal cord protection during TEVAR (Rizvi and Sullivan). 2010;52:S86-90
Intraprocedural imaging: flat panel detectors, rotational angiography, FluoroCT, IVUS, or still the portable C-arm? (Eagleton). 2010;52:S50-9
Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Roddy). 2010;52: 1419
Left subclavian artery revascularization: Society for Vascular Surgery® Practice Guidelines (Matsumura and Rizvi). 2010; 52:S65-70
Limb ischemia during femoral cannulation for cardiopulmonary support (Roddy). 2010;52:1109
Open techniques for arch vessel reconstruction during thoracic endovascular aneurysm repair (TEVAR) (Vallabhaneni and Sanchez). 2010;52:S71-6
The Pacific Vascular Symposium 6: the Venous Ulcer Summit in perspective (Henke). 2010;52:S1-2
Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Roddy). 2010;52:250
Prosthetic lower extremity haemodialysis access grafts have satisfactory patency despite high incidence of infection (Roddy). 2010;52:1723
A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Roddy). 2010;52: 796
A real world analysis of payment per unit time in a Maryland Vascular Practice (Roddy). 2010;52:1110
Regulatory TEVAR clinical trials (Garcia-Toca and Eskandari). 2010;52:S22-5
Required skills and interdisciplinary teams in starting a TEVAR practice (Starr). 2010;52:S45-9
Rescuing Sisyphus: the team approach to amputation prevention (Mills Sr et al). 2010;52:S1-2
Thoracic aortic endovascular repair for mycotic aneurysms and fistulas (Patel et al). 2010;52:S37-40
Traumatic thoracic aortic injury and ruptures (Tefera). 2010; 52:S41-4
Trends in vena caval interruption (Roddy). 2010;52:251
Visceral reconstruction techniques (Srivastava). 2010;52: S82-5

Education, medical
A vascular disease educational program in the preclinical years of medical school increases student interest in vascular disease (Godshall et al). 2010;52:775-81, 775.e1-e2

Education, medical, graduate
Incorporating simulation in vascular surgery education (Bismuth et al). 2010;52:1072-80

Elastomers
The effect of injectable biocompatible elastomer (PDMS) on the strength of the proximal fixation of endovascular aneurysm repair grafts: an in vitro study (Bosman et al). 2010; 52:152-8

Elderly; *see* Aged

Ellagic acid
Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells (Lee et al); (Rzucidlo) (Commentary). 2010; 52:1290-300

Embolic protection devices
CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection (Matsumura et al). 2010;52:576-83, 576.e1-e2

Embolization, therapeutic
Early side effects after embolization of a carotid body tumor using Onyx (Wiegand et al). 2010;52:742-5 (Case rep.)
Glue embolus complicating the endovascular treatment of a patient with Loeys-Dietz syndrome (Marine et al). 2010; 52:1350-3 (Case rep.)

Emergencies
Emergency endovascular treatment of early spontaneous nonaneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia (Trellopoulos et al). 2010;52:751-7 (Case rep.)

Endarterectomy
Histologic atherosclerotic plaque characteristics are associated with restenosis rates after endarterectomy of the common and superficial femoral arteries (Derksen et al). 2010;52: 592-9

Endarterectomy, carotid
Asymptomatic carotid artery stenosis-Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting (Schneider and Naylor); (Forbes) (Commentary) 2010;52:499-507. Correction. 2010;52:823
Asymptomatic carotid artery stenosis—Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting (Schneider and Naylor); (Forbes) (Commentary). 2010;52:499-507
Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis (AbuRahma et al). 2010;52:1180-7
Deglutition syncope: a manifestation of vagal hyperactivity following carotid endarterectomy (Endean et al). 2010;52: 720-2 (Case rep.)
Eight-year experience with carotid artery stenting for correction of symptomatic and asymptomatic post-endarterectomy defects. 2010;52:1511-7
Mandibular subluxation stabilized by mouthpiece for distal internal carotid artery exposure in carotid endarterectomy (Yoshino et al). 2010;52:1401-4
Mechanisms of gender-related outcome differences after carotid endarterectomy (den Hartog et al). 2010;52:1062-71, 1062.e1-e6 (Presidential Address)
Outcomes of carotid stenting compared with endarterectomy are equivalent in asymptomatic patients and inferior in

symptomatic patients (Giacovelli et al). 2010;52:906-13, 906.e1-e4
Process of care for carotid endarterectomy: perioperative medical management (Stoner and deFreitas). 2010;52:223-31 (Review)
Restenosis after carotid endarterectomy in a multicenter regional registry (Goodney et al). 2010;52:897-905, 897.e1-e2
Stroke and death after carotid endarterectomy and carotid artery stenting with and without high risk criteria (Giles et al). 2010;52:1497-504

Endoleak
Duplex ultrasound factors predicting persistent type II endoleak and increasing AAA sac diameter after EVAR (Beeman et al). 2010;52:1147-52
Early type III endoleak with an Endurant endograft (Aboulian et al). 2010;52:1665-7 (Case rep.)
The effect of warfarin therapy on endoleak development after endovascular aneurysm repair (EVAR) of the abdominal aorta (Bobadilla et al). 2010;52:267-71
Endoleak management and postoperative surveillance following endovascular repair of thoracic aortic aneurysms (Ricotta). 2010;52:S91-9
Endoluminal treatment of type IA endoleak with Onyx (Grisafi et al). 2010;52:1346-9 (Case rep.)
Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Abularrage et al). 2010;52:19-24
Preoperative variables predict persistent type 2 endoleak after endovascular aneurysm repair (Roddy). 2010;52:250 (Editorial)

Endoluminal treatment; *see* Angioplasty

Endoplasmic reticulum
Sodium 4-phenylbutyrate protects against spinal cord ischemia by inhibition of endoplasmic reticulum stress (Mizukami et al). 2010;52:1580-6

Endothelial cells
Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells (Lee et al); (Rzucidlo) (Commentary). 2010;52:1290-300
Endothelial cells are susceptible to rapid siRNA transfection and gene silencing ex vivo (Andersen et al). 2010;52:1608-15
Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20

Endothelium, vascular
Interleukin-6 and interleukin-10 gene polymorphism, endothelial dysfunction, and postoperative prognosis in patients with peripheral arterial disease (Stoica et). 2010;52:103-9

End-stage renal disease; *see* Kidney failure, chronic

Enoxaparin
Long-term efficacy and safety of once-daily enoxaparin plus warfarin for the outpatient ambulatory treatment of lower-limb deep vein thrombosis in the TROMBOTEK trial (Kurtoglu et al); (Henke) (Commentary). 2010;52:1262-71

Ethics, medical
The ethical hierarchy of do not resuscitate orders: Never say never (Jones and McCullough). 2010;52:1384-6

Ethics, professional
Regarding "How do we guarantee trainee professional purity?" (Nakayama) (Letter); (Jones and McCullough) (Reply). 2010;52:257

Evidence-based medicine
Non-medical initiatives to decrease venous ulcers prevalence (Passman). 2010;52:S29-36

Exercise therapy
Multicenter randomized clinical trial of supervised exercise therapy with or without feedback versus walking advice for intermittent claudication (Nicolaï et al). 2010;52:348-55
Optimizing supervised exercise therapy for patients with intermittent claudication (Nicolaï et al). 2010;52:1226-33

Extracorporeal shock wave therapy; *see* Lithotripsy

Extremities
An elevated neutrophil-lymphocyte ratio independently predicts mortality in chronic critical limb ischemia (Spark et al). 2010;52:632-6
Comparison of the clinical history of symptomatic isolated muscular calf vein thrombosis versus deep calf vein thrombosis (Galanaud et al). 2010;52:932-8, 932.e1-e2
Finnvasc score and modified Prevent III score predict long-term outcome after infrainguinal surgical and endovascular revascularization for critical limb ischemia (Arvela et al). 2010;52:1218-25
Limb ischemia during femoral cannulation for cardiopulmonary support (Foley et al). 2010;52:850-3
Limb ischemia during femoral cannulation for cardiopulmonary support (Roddy). 2010;52:1109 (Editorial)
Outcomes of selective tibial artery repair following combat-related extremity injury (Burkhardt et al). 2010;52:91-6
Perioperative platelet and monocyte activation in patients with critical limb ischemia (Burdess et al). 2010;52:697-703
Predictors of failure and success of tibial interventions for critical limb ischemia (Fernandez et al). 2010;52:834-42
Quality of life in patients with no-option critical limb ischemia underlines the need for new effective treatment (Sprengers et al). 2010;52:843-9, 843.e1

F

Fascia
Regarding "The anatomy of the small saphenous vein: fascial and neural relations, saphenofemoral junction, and valves" (Caggiati) (Letter); (Schweighofer et al) (Reply). 2010;52:1428-9

Feedback
Multicenter randomized clinical trial of supervised exercise therapy with or without feedback versus walking advice for intermittent claudication (Nicolaï et al). 2010;52:348-55

Femoral artery
An analysis of relationship between quality of life indices and clinical improvement following intervention in patients with intermittent claudication due to femoropopliteal disease (Mazari et al). 2010;52:77-84
A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Boufi et al). 2010;52:1211-17
A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Roddy). 2010;52:1417-18 (Editorial)
A contemporary meta-analysis of Dacron versus polytetrafluo-

roethylene grafts for femoropopliteal bypass grafting (Takagi et al). 2010;52:232-6 (Review)
D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Hoke et al). 2010;52:383-7
D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Roddy). 2010;52:517 (Editorial)
The first operation on the profunda femoris artery (Yao). 2010;52:1101-2
Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease (McQuade et al). 2010;52:584-91, 584.e1-e8
A giant femoral artery pseudoaneurysm in an infant boy (Park et al). 2010;52:222
Histologic atherosclerotic plaque characteristics are associated with restenosis rates after endarterectomy of the common and superficial femoral arteries (Derksen et al). 2010;52:592-9
The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization (Durham et al). 2010;52:600-7
Limb ischemia during femoral cannulation for cardiopulmonary support (Foley et al). 2010;52:850-3
Limb ischemia during femoral cannulation for cardiopulmonary support (Roddy). 2010;52:1109 (Editorial)
Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent (Soga et al). 2010;52:608-15
Outcomes of interventions for recurrent disease after endoluminal intervention for superficial femoral artery disease (Davies et al). 2010;52:331-9, 331.e1-e2
Preparation and characterization of injectable fibrillar type I collagen and evaluation for pseudoaneurysm treatment in a pig model (Geutjes et al). 2010;52:1330-8
Subintimal angioplasty for femoro-popliteal occlusive disease (Markose et al). 2010;52:1410-16
True lumen re-entry after extravascular recanalization of a superficial femoral artery chronic total occlusion (Wagner et al). 2010;52:216-18 (Case rep.)

Femoral vein
Anatomical variations of the femoral vein (Uhl et al). 2010;52:714-19
Endovenectomy of the common femoral vein and intraoperative iliac vein recanalization for chronic iliofemoral venous occlusion (Comerota et al). 2010;52:243-7
Iliofemoral venous thrombosis from external compression by a vesical diverticulum (Gupta et al). 2010;52:1671-3 (Case rep.)
Period frequency of iliofemoral venous occlusive disease by Doppler ultrasound and corresponding treatment in a tertiary care facility (Crisostomo et al). 2010;52:1272-7

Fibrillar collagens
Preparation and characterization of injectable fibrillar type I collagen and evaluation for pseudoaneurysm treatment in a pig model (Geutjes et al). 2010;52:1330-8

Fibrin fibrinogen degradation products
D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Hoke et al). 2010;52:383-7
D-dimer testing in the diagnosis of transfemoral pseudoaneurysm after percutaneous transluminal procedures (Roddy). 2010;52:517 (Editorial)

Fibromuscular dysplasia
A rare case of familial carotid body tumor in a patient with bilateral fibromuscular dysplasia (Han et al). 2010;52:746-50 (Case rep.)

Fistula
Aorto-left renal vein fistula is a rare complication of abdominal aortic aneurysm with unique clinical presentation (Dragas et al). 2010;52:1658-61 (Case rep.)
Mechanosensitive transient receptor potential vanilloid type 1 channels contribute to vascular remodeling of rat fistula veins (Chen et al). 2010;52:1310-20
The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease (Bonforte et al). 2010;52:1551-6
Thoracic aortic endovascular repair for mycotic aneurysms and fistulas (Patel et al). 2010;52:S37-40 (Editorial)

Fluorescent dyes
A novel method of measuring human lymphatic pumping using indocyanine green fluorescence lymphography (Unno et al). 2010;52:946-52

Fluoroscopy
Intraprocedural imaging: flat panel detectors, rotational angiography, FluoroCT, IVUS, or still the portable C-arm? (Eagleton). 2010;52:S50-9 (Editorial)

Foot deformities
Surgical off-loading of the diabetic foot (Frykberg et al). 2010;52:S44-58

Foot ulcer
Off-loading the diabetic foot for ulcer prevention and healing (Cavanagh and Bus). 2010;52:S37-43
Surgical off-loading of the diabetic foot (Frykberg et al). 2010;52:S44-58

Foreign-body migration
Novel technique for endovascular salvage of a folded aortic endograft (Ricotta II and Malgor). 2010;52:785-9

G

Gait
Abnormal joint powers before and after the onset of claudication symptoms (Koutakis et al). 2010;52:340-7

Ganglionectomy
Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Roddy). 2010;52:1419 (Editorial); Tulloch et al). 2010;52:1283-9

Gangrene
Tuberculosis should not be ignored in patients with peripheral gangrene (Yao et al). 2010;52:1662-4 (Case rep.)

Gender identity
Gender differences in daily ambulatory activity patterns in patients with intermittent claudication (Gardner et al). 2010;52:1204-10
Lessons learned from the analysis of gender effect on risk factors and procedural outcomes of lower extremity arterial disease (Vouyouka et al); (Bush) (Commentary). 2010;52:1196-203

Gene knockdown techniques
Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

Gene polymorphism; *see* Polymorphism, genetic

Genes
eNOS and *ACE* genes influence peripheral arterial disease predisposition in smokers (Sticchi et al). 2010;52:97-102, 97.e1
Gene silencing
Endothelial cells are susceptible to rapid siRNA transfection and gene silencing ex vivo (Andersen et al). 2010;52:1608-15
Gene therapy
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
Glue; *see* Adhesives
Glycoproteins
Tissue factor pathway inhibitor-2 is induced by fluid shear stress in vascular smooth muscle cells and affects cell proliferation and survival (Ekstrand et al). 2010;52:167-75
Gonadal steroid hormones
Elevated sex steroid hormones in great saphenous veins in men (Kendler et al); (Labropoulos) (Commentary). 2010;51:639-47. Correction. 2010;52:538
Graft occlusion, vascular
Histologic atherosclerotic plaque characteristics are associated with restenosis rates after endarterectomy of the common and superficial femoral arteries (Derksen et al). 2010;52:592-9
Graft survival
Rationale and practical techniques for mouse models of early vein graft adaptations (Yu et al). 2010;52:444-52
Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010;52:1425-7
Guidelines as topic
Consensus statement on the adoption of the COPE guidelines. 2010;52:242

H

Haiti
A fellow's experience in Haiti (Lind). 2010;52:533-4 (Letter)
Haptoglobins
Haptoglobin 2-1 phenotype predicts rapid growth of abdominal aortic aneurysms (Wiernicki et al). 2010;52:691-6
Health care costs
The costs of diabetic foot: the economic case for the limb salvage team (Driver et al). 2010;52:S17-22
A real world analysis of payment per unit time in a Maryland Vascular Practice (Martin et al); (O'Donnell) (Commentary). 2010;52:1094-100
A real world analysis of payment per unit time in a Maryland Vascular Practice (Roddy). 2010;52:1110 (Editorial)
Health resources
Regarding "Long-term outcomes and resource utilization of endovascular versus open repair of abdominal aortic aneurysms in Ontario" (Takagi et al). 2010;52:818-19 (Letter)
Health status indicators
The Vascular Study Group of New England Cardiac Risk Index (VSG-CRI) predicts cardiac complications more accurately than the Revised Cardiac Risk Index in vascular 4surgery patients (Bertges et al). 2010;52:674-83, 674.e1-e3
Heart
A one-stage approach to the treatment of intravenous leiomyomatosis extending to the right heart (Rispoli et al). 2010;52:212-15 (Case rep.)
Heart diseases
The Vascular Study Group of New England Cardiac Risk Index (VSG-CRI) predicts cardiac complications more accurately than the Revised Cardiac Risk Index in vascular surgery patients (Bertges et al). 2010;52:674-83, 674.e1-e3
Hematoma
Spontaneous retroperitoneal hematoma associated with iliac vein rupture (Jiang et al). 2010;52:1278-82
Heme oxygenase-1
Regarding "Statins, heme oxygenase-1, iron, and atherosclerosis" (Sullivan) (Letter); (DePalma et al) (Reply). 2010;52:536-7
Hemodialysis; *see* Renal dialysis
Hemodynamics
Aortic endograft sizing in trauma patients with hemodynamic instability (Jonker et al). 2010;52:39-44
Impact of obesity on venous hemodynamics of the lower limbs (Willenberg et al). 2010;52:664-8
Inelastic bandages maintain their hemodynamic effectiveness over time despite significant pressure loss (Mosti and Partsch:). 2010;52:925-31
Noninvasive assessment of lower extremity hemodynamics in individuals with diabetes mellitus (Andersen). 2010;52:S76-80
Heparin
Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: results from a prospective randomized study (Stabile et al). 2010;52:1505-10
Hepatocyte growth factor
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
HGF; *see* Hepatocyte growth factor
Hindlimb
Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20
Hirudins
Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: results from a prospective randomized study (Stabile et al). 2010;52:1505-10
A pilot, prospective evaluation of a direct thrombin inhibitor, bivalirudin (Angiomax), in patients undergoing lower extremity bypass (Kashyap et al). 2010;52:369-74
Homografts; *see* Transplantation, homologous
Hospital costs
The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization (Durham et al). 2010;52:600-7
Hospitals
Evaluating parsimonious risk-adjustment models for comparing hospital outcomes with vascular surgery (Osborne et al). 2010;52:400-5

Hydroxymethylglutaryl-CoA reductase inhibitors
A meta-analysis of clinical studies of statins for prevention of abdominal aortic aneurysm expansion (Takagi et al). 2010; 52:1675-81 (Review)
Regarding "Preoperative statin therapy is associated with improved outcomes and resource utilization in patients undergoing aortic aneurysm repair" (Takagi et al) (Letter); (Stoner) (Reply). 2010;52:820-2
Regarding "Statins, heme oxygenase-1, iron, and atherosclerosis" (Sullivan) (Letter); (DePalma et al) (Reply). 2010;52: 536-7
Hyperbaric oxygenation
Wound care: the role of advanced wound healing technologies (Wu et al). 2010;52:S59-66
Hyperglycemia
Hyperglycemia limits experimental aortic aneurysm progression (Miyama et al). 2010;52:975-83
Hyperplasia
A novel cell permeant peptide inhibitor of MAPKAP kinase II inhibits intimal hyperplasia in a human saphenous vein organ culture model (Lopes et al). 2010;52:1596-607
Hypothermia
The effects of systemic hypothermia on a murine model of thoracic aortic ischemia reperfusion (Kang et al). 2010;52: 435-43
Therapeutic distant organ effects of regional hypothermia during mesenteric ischemia-reperfusion injury (Santora et al). 2010;52:1003-14
Hypoxia-inducible factor 1, alpha subunit
Evidence for markers of hypoxia and apoptosis in explanted human carotid atherosclerotic plaques (Bitto et al). 2010; 52:1015-21

I

Iatrogenic disease
Agitation thrombolysis for fresh iatrogenic IVC thrombosis in patients with Budd-Chiari syndrome (Ding et al). 2010;52: 782-4
Iliac artery
A durable iliac-axillary and axillary-carotid bypass for cerebral ischemia due to Takayasu arteritis allowed successful pregnancies and deliveries (Kondo et al). 2010;52:1713-5
Endovascular procedures for aorto-iliac occlusive disease are associated with superior short-term clinical and economic outcomes compared with open surgery in the inpatient population (Indes et al). 2010;52:1173-9, 1173.e1
External iliac artery dissection secondary to endofibrosis in a cyclist (Willson et al). 2010;52:219-21 (Case rep.)
Midterm outcome of endovascular repair of ruptured isolated iliac artery aneurysms (Hechelhammer et al). 2010;52:1159-63
Practical use of duplex echo-guided recanalization of chronic total occlusion in the iliac artery (Kawarada et al). 2010;52: 475-8 (Case rep.)
Regarding "Endovascular management of iliac rupture during endovascular aneurysm repair" (Semel et al). 2010;52:533 (Letter)
Staged total exclusion of the aorta for chronic type B aortic dissection (Dias Perera et al). 2010;52:1339-42 (Case rep.)
A systematic review of endovascular treatment of extensive aortoiliac occlusive disease (Jongkind et al). 2010;52:1376-83 (Review)
Iliac vein
Endovenectomy of the common femoral vein and intraoperative iliac vein recanalization for chronic iliofemoral venous occlusion (Comerota et al). 2010;52:243-7
Iliofemoral venous thrombosis from external compression by a vesical diverticulum (Gupta et al). 2010;52:1671-3 (Case rep.)
Period frequency of iliofemoral venous occlusive disease by Doppler ultrasound and corresponding treatment in a tertiary care facility (Crisostomo et al). 2010;52:1272-7
Spontaneous retroperitoneal hematoma associated with iliac vein rupture (Jiang et al). 2010;52:1278-82
Immunoglobulin G
A clinicopathologic study of immunoglobulin G4-related sclerosing disease of the thoracic aorta (Kasashima et al). 2010; 52:1587-95
Incidental findings
Incidence, follow-up, and outcomes of incidental abdominal aortic aneurysms (van Walraven et al). 2010;52:282-9, 282.e1-e2
Indocyanine green
A novel method of measuring human lymphatic pumping using indocyanine green fluorescence lymphography (Unno et al). 2010;52:946-52
Infant
A giant femoral artery pseudoaneurysm in an infant boy (Park et al). 2010;52:222
Infection
Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Bisdas et al). 2010;52:323-30
Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Roddy). 2010;52:516 (Editorial)
Prosthetic lower extremity haemodialysis access grafts have satisfactory patency despite high incidence of infection (Roddy). 2010;52:1723 (Editorial)
Prosthetic lower extremity hemodialysis access grafts have satisfactory patency despite a high incidence of infection (Geenen et al). 2010;52:1546-50
Inferior vena cava; *see* Vena cava, inferior
Inflammation
Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells (Lee et al); (Rzucidlo) (Commentary). 2010; 52:1290-300
Markers of inflammation in men with small abdominal aortic aneurysm (Parry et al). 2010;52:145-51
Infrainguinal bypass surgery; *see* Vascular surgical procedures
Infrarenal aortic aneurysm; *see* Aortic aneurysm, abdominal
Injections, intramuscular
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
Injury; *see* Wounds and injuries
Innominate vein; *see* Brachiocephalic veins
Inpatients
Endovascular procedures for aorto-iliac occlusive disease are associated with superior short-term clinical and economic outcomes compared with open surgery in the inpatient population (Indes et al). 2010;52:1173-9, 1173.e1

Open and endovascular repair of type B aortic dissection in the Nationwide Inpatient Sample (Sachs et al). 2010;52:860-6

Interdisciplinary teams; *see* Patient care team

Interleukin-6

Interleukin-6 and interleukin-10 gene polymorphism, endothelial dysfunction, and postoperative prognosis in patients with peripheral arterial disease (Stoica et). 2010;52:103-9

Intermediate filament proteins

Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

Intermittent claudication

Abnormal joint powers before and after the onset of claudication symptoms (Koutakis et al). 2010;52:340-7

An analysis of relationship between quality of life indices and clinical improvement following intervention in patients with intermittent claudication due to femoropopliteal disease (Mazari et al). 2010;52:77-84

Bilateral nonthrombotic subclavian vein obstruction causing upper extremity venous claudication (Thakur and Comerota). 2010;52:208-11 (Case rep.)

Gender differences in daily ambulatory activity patterns in patients with intermittent claudication (Gardner et al). 2010;52:1204-10

Multicenter randomized clinical trial of supervised exercise therapy with or without feedback versus walking advice for intermittent claudication (Nicolaï et al). 2010;52:348-55

Optimizing supervised exercise therapy for patients with intermittent claudication (Nicolaï et al). 2010;52:1226-33

Internship and residency

Incorporating simulation in vascular surgery education (Bismuth et al). 2010;52:1072-80

Regarding "How do we guarantee trainee professional purity?" (Nakayama) (Letter); (Jones and McCullough) (Reply). 2010;52:257

Intestinal obstruction

Urgent interventional bilateral renal artery fenestration for giant pararenal abdominal aortic aneurysm with upper digestive tract obstruction (Ouyang and Liang). 2010;52:1048-51 (Case rep.)

Intracellular signaling peptides and proteins

A novel cell permeant peptide inhibitor of MAPKAP kinase II inhibits intimal hyperplasia in a human saphenous vein organ culture model (Lopes et al). 2010;52:1596-607

Intracranial embolism

Atherosclerotic plaque composition assessed by virtual histology intravascular ultrasound and cerebral embolization after carotid stenting (Timaran et al). 2010;52:1188-95

Catheter-based neurosalvage for acute embolic complication during carotid intervention (Lin et al). 2010;52:308-13

Intravascular ultrasound imaging; *see* Ultrasonography, interventional

Iron

Regarding "Statins, heme oxygenase-1, iron, and atherosclerosis" (Sullivan) (Letter); (DePalma et al) (Reply). 2010;52:536-7

Ischemia

An elevated neutrophil-lymphocyte ratio independently predicts mortality in chronic critical limb ischemia (Spark et al). 2010;52:632-6

An unfavorable dietary pattern is associated with symptomatic ischemic stroke and carotid atherosclerosis (Mahe et al). 2010;52:62-8

Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Arvela et al). 2010;52:616-23

Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Roddy). 2010;52:794-5 (Editorial)

Arterial imaging in patients with lower extremity ischemia and diabetes mellitus (Pomposelli). 2010;52:S81-91

Banding between dialysis puncture sites to treat severe ischemic steal syndrome in low flow autogenous arteriovenous access (Shemesh et al). 2010;52:495-8

Below knee bare nitinol stent placement in high-risk patients with critical limb ischemia is still durable after 24 months of follow-up (Donas et al). 2010;52:356-61

Enhanced cell therapy strategy to treat chronic limb-threatening ischemia (Eton and Yu). 2010;52:199-204 (Case rep.)

Finnvasc score and modified Prevent III score predict long-term outcome after infrainguinal surgical and endovascular revascularization for critical limb ischemia (Arvela et al). 2010;52:1218-25

Limb ischemia during femoral cannulation for cardiopulmonary support (Foley et al). 2010;52:850-3; Roddy). 2010;52:1109 (Editorial)

Microcirculatory effects of local and remote ischemic preconditioning in supraceliac aortic clamping (Erling et al). 2010;52:1321-9

Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20

Perioperative platelet and monocyte activation in patients with critical limb ischemia (Burdess et al). 2010;52:697-703

Predictors of failure and success of tibial interventions for critical limb ischemia (Fernandez et al). 2010;52:834-42

Quality of life in patients with no-option critical limb ischemia underlines the need for new effective treatment (Sprengers et al). 2010;52:843-9, 843.e1

Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration:results of the HGF-0205 trial (Powell et al). 2010;52:1525-30

Thrombotic mesenteric ischemia due to aortic dissection (Bao and Chen). 2010;52:1674, 1674.e1-e2

Upper extremity ischemia treated with tissue repair cells from adult bone marrow (Comerota et al). 2010;52:723-9 (Case rep.)

Ischemia reperfusion; *see* Reperfusion injury

J

Janus kinase 2

JAK2 V617F mutation, mesenteric vein thrombosis, and myeloproliferative disorders (Owens). 2010;52:205-7 (Case rep.)

Joints

Abnormal joint powers before and after the onset of claudication symptoms (Koutakis et al). 2010;52:340-7

K

Kidney diseases

The influence of the obesity paradox and chronic kidney disease on long-term survival in a Japanese cohort with peripheral arterial disease (Kumakura et al). 2010;52:110-17

Juxtarenal aortic aneurysm repair (Jongkind et al). 2010;52:760-7 (Review)

The middle-arm fistula as a valuable surgical approach in pa-

tients with end-stage renal disease (Roddy). 2010;52: 1724-5

Kidney failure, chronic

The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease (Bonforte et al). 2010; 52:1551-6

Revascularization of an occluded brachiocephalic vein using Outback-LTD re-entry catheter (Anil and Taneja). 2010; 52:1038-40 (Case rep.)

Klippel-Trenaunay-Weber syndrome

Lymphatic malformation is a common component of Klippel-Trenaunay syndrome (Liu et al). 2010;52:1557-63

Knee

Below knee bare nitinol stent placement in high-risk patients with critical limb ischemia is still durable after 24 months of follow-up (Donas et al). 2010;52:356-61

L

Laparoscopy

Laparoscopic transposition of the left renal vein into the inferior vena cava for nutcracker syndrome (Hartung et al). 2010;52:738-41 (Case rep.)

Laparoscopic treatment of renal artery entrapment (Bünger et al). 2010;52:1357-61 (Case rep.)

Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Roddy). 2010;52: 1419 (Editorial)

Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Tulloch et al). 2010;52:1283-9

Laser ablation; *see* Laser therapy

Laser therapy

Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up (Christenson et al). 2010;52:1234-41

A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Gale et al). 2010; 52:645-50

A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Roddy). 2010;52: 796 (Editorial)

Leg

Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Arvela et al). 2010;52: 616-23

Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Roddy). 2010;52:794-5 (Editorial)

Identifying the source of superficial reflux in venous leg ulcers using duplex ultrasound (Obermayer and Garzon). 2010; 52:1255-61

Leg strength predicts mortality in men but not in women with peripheral arterial disease (Singh et al). 2010;52:624-31

Leiomyomatosis

A one-stage approach to the treatment of intravenous leiomyomatosis extending to the right heart (Rispoli et al). 2010; 52:212-15 (Case rep.)

Letters to the Editor

Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9

A fellow's experience in Haiti (Lind). 2010;52:533-4 (Letter)

Regarding "A closer look at meta-analyses of observational data" (Navarese et al) (Letter); (Jonker et al) (Reply). 2010;52:819-20

Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Lee). 2010;52:533 (Letter)

Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Skurnick) (Letter); (Zamboni) (Reply). 2010;52:1749-50

Regarding "Endovascular management of iliac rupture during endovascular aneurysm repair" (Semel et al). 2010;52:533 (Letter)

Regarding "Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I" (Mendonça). 2010;52:1425 (Letter)

Regarding "Good Samaritan statutes: a malpractice defense for doing the right thing" (Semel) (Letter); (Brown) (Reply). 2010;52:1125

Regarding "High spatial resolution magnetic resonance imaging of cystic adventitial disease of the popliteal artery" (Weimar et al) (Letter); (Hagspiel) (Reply). 2010;52:534-5

Regarding "How do we guarantee trainee professional purity?" (Nakayama) (Letter); (Jones and McCullough) (Reply). 2010;52:257

Regarding "Long-term outcomes and resource utilization of endovascular versus open repair of abdominal aortic aneurysms in Ontario" (Takagi et al). 2010;52:818-19 (Letter)

Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010; 52:1425-7

Regarding "Peripheral arterial interventions: Trends in market shareandoutcomesbyspecialty,1998-2005"(Sacks)(Letter); (Eslami) (Reply). 2010;52:528-9

Regarding "Preoperative statin therapy is associated with improved outcomes and resource utilization in patients undergoing aortic aneurysm repair" (Takagi et al) (Letter); (Stoner) (Reply). 2010;52:820-2

Regarding "Right subclavian pseudoaneurysm secondary to blunt trauma in an arteriovenous malformation" (Lee) (Letter);(Quinones-Baldrich,) (Reply). 2010;52:257-8

Regarding "Serum carboxymethyllysine concentrations are reduced in diabetic men with abdominal aortic aneurysms: Health In Men Study" (Meerwaldt and Zeebregts) (Letter); (Norman et al) (Reply). 2010;52:535-6

Regarding "Statins, heme oxygenase-1, iron, and atherosclerosis" (Sullivan) (Letter); (DePalma et al) (Reply). 2010;52: 536-7

Regarding "The anatomy of the small saphenous vein: fascial and neural relations, saphenofemoral junction, and valves" (Caggiati) (Letter); (Schweighofer et al) (Reply). 2010;52: 1428-9

Regarding "The impact of model assumptions on results of computational mechanics in abdominal aortic aneurysm" (Georgakarakos et al) (Letter); (Reeps and Gee) (Reply). 2010;52:1124-5

Ligaments
Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Roddy). 2010;52:1419 (Editorial)
Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome (Tulloch et al). 2010;52:1283-9
Limbs; *see* Extremities
Limb salvage
The costs of diabetic foot: the economic case for the limb salvage team (Driver et al). 2010;52:S17-22
Limb salvage after delayed diagnosis for blunt traumatic infrapopliteal occlusion (Causey et al). 2010;52:734-7 (Case rep.)
Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010;52:1425-7
Lipectomy
Superficialization of arteriovenous fistulae employing minimally invasive liposuction (Causey et al). 2010;52:1397-400
Lipoproteins, LDL
Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells (Lee et al); (Rzucidlo) (Commentary). 2010;52:1290-300
Oxidized LDL in human carotid plaques is related to symptomatic carotid disease and lesion instability (Sigala et al). 2010;52:704-13
Liposuction; *see* Lipectomy
Listeria monocytogenes
Listeria monocytogenes as a rare cause of mycotic aortic aneurysm (Bal et al). 2010;52:456-9 (Case rep.)
Lithotripsy
Extracorporeal shock wave therapy ameliorates secondary lymphedema by promoting lymphangiogenesis (Kubo et al). 2010;52:429-34
Loeys-Dietz Syndrome
Glue embolus complicating the endovascular treatment of a patient with Loeys-Dietz syndrome (Marine et al). 2010;52:1350-3 (Case rep.)
Long-term care
Regarding "Long-term outcomes and resource utilization of endovascular versus open repair of abdominal aortic aneurysms in Ontario" (Takagi et al). 2010;52:818-19 (Letter)
Lower extremity
Abnormal joint powers before and after the onset of claudication symptoms (Koutakis et al). 2010;52:340-7
Arterial imaging in patients with lower extremity ischemia and diabetes mellitus (Pomposelli). 2010;52:S81-91
Below knee bare nitinol stent placement in high-risk patients with critical limb ischemia is still durable after 24 months of follow-up (Donas et al). 2010;52:356-61
Enhanced cell therapy strategy to treat chronic limb-threatening ischemia (Eton and Yu). 2010;52:199-204 (Case rep.)
Hull early walking aid for rehabilitation of transtibial amputees - randomized controlled trial (HEART) (Mazari et al). 2010;52:1564-71
Impact of obesity on venous hemodynamics of the lower limbs (Willenberg et al). 2010;52:664-8
Lessons learned from the analysis of gender effect on risk factors and procedural outcomes of lower extremity arterial disease (Vouyouka et al); (Bush) (Commentary). 2010;52:1196-203
Long-term efficacy and safety of once-daily enoxaparin plus warfarin for the outpatient ambulatory treatment of lower-limb deep vein thrombosis in the TROMBOTEK trial (Kurtoglu et al); (Henke) (Commentary). 2010;52:1262-71
Noninvasive assessment of lower extremity hemodynamics in individuals with diabetes mellitus (Andersen). 2010;52:S76-80
A pilot, prospective evaluation of a direct thrombin inhibitor, bivalirudin (Angiomax), in patients undergoing lower extremity bypass (Kashyap et al). 2010;52:369-74
Poor outcomes with cryoplasty for lower extremity arterial occlusive disease (Schmieder et al). 2010;52:362-8
Prosthetic lower extremity haemodialysis access grafts have satisfactory patency despite high incidence of infection (Roddy). 2010;52:1723 (Editorial)
Prosthetic lower extremity hemodialysis access grafts have satisfactory patency despite a high incidence of infection (Geenen et al). 2010;52:1546-50
Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010;52:1425-7
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
LOX-1; *see* Scavenger receptors, class E
Lung
Therapeutic distant organ effects of regional hypothermia during mesenteric ischemia-reperfusion injury (Santora et al). 2010;52:1003-14
Lymphangiogenesis
Extracorporeal shock wave therapy ameliorates secondary lymphedema by promoting lymphangiogenesis (Kubo et al). 2010;52:429-34
Lymphatic system
Lymphatic malformation is a common component of Klippel-Trenaunay syndrome (Liu et al). 2010;52:1557-63
Lymphatic vessels
A novel method of measuring human lymphatic pumping using indocyanine green fluorescence lymphography (Unno et al). 2010;52:946-52
Lymphedema
Extracorporeal shock wave therapy ameliorates secondary lymphedema by promoting lymphangiogenesis (Kubo et al). 2010;52:429-34
Lymphocytes
An elevated neutrophil-lymphocyte ratio independently predicts mortality in chronic critical limb ischemia (Spark et al). 2010;52:632-6
Lymphography
A novel method of measuring human lymphatic pumping using indocyanine green fluorescence lymphography (Unno et al). 2010;52:946-52
Lysine
Regarding "Serum carboxymethyllysine concentrations are reduced in diabetic men with abdominal aortic aneurysms:

Health In Men Study" (Meerwaldt and Zeebregts) (Letter); (Norman et al) (Reply). 2010;52:535-6

M

Magnetic resonance imaging
Regarding "High spatial resolution magnetic resonance imaging of cystic adventitial disease of the popliteal artery" (Weimar et al) (Letter); (Hagspiel) (Reply). 2010;52:534-5

Malpractice
Regarding "Good Samaritan statutes: a malpractice defense for doing the right thing" (Semel) (Letter); (Brown) (Reply). 2010;52:1125

Mandible
Mandibular subluxation stabilized by mouthpiece for distal internal carotid artery exposure in carotid endarterectomy (Yoshino et al). 2010;52:1401-4

Manuscripts
There may be nothing new under the sun, but this is ridiculous (Brewster and Greisler). 2010;52:1697-8

MAPKAP kinase II; *see* Intracellular signaling peptides and proteins; Protein-serine-threonine kinases

Marfan syndrome
Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques (Takahashi et al). 2010;52:1365-6

Mass screening
An analysis of drug modulation of abdominal aortic aneurysm growth through 25 years of surveillance (Thompson et al). 2010;52:55-61, 55.e1-e2
A further meta-analysis of population-based screening for abdominal aortic aneurysm (Takagi et al). 2010;52:1103-8

Medical school; *see* Schools, medical

Men's health
Regarding "Serum carboxymethyllysine concentrations are reduced in diabetic men with abdominal aortic aneurysms: Health In Men Study" (Meerwaldt and Zeebregts) (Letter); (Norman et al) (Reply). 2010;52:535-6

Mesenteric arteries
Successful endovascular treatment of multiple mesenteric arterial aneurysms associated with alpha-1-antitrypsin deficiency (Gordon et al). 2010;52:467-70 (Case rep.)
Thrombotic mesenteric ischemia due to aortic dissection (Bao and Chen). 2010;52:1674, 1674.e1-e2

Mesenteric artery, superior
Endovascular and open surgery for acute occlusion of the superior mesenteric artery (Block et al). 2010;52:959-66
Regarding "Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I" (Mendonça). 2010;52:1425 (Letter)

Mesenteric vascular occlusion
Endovascular and open surgery for acute occlusion of the superior mesenteric artery (Block et al). 2010;52:959-66
Therapeutic distant organ effects of regional hypothermia during mesenteric ischemia-reperfusion injury (Santora et al). 2010;52:1003-14

Mesenteric veins
JAK2 V617F mutation, mesenteric vein thrombosis, and myeloproliferative disorders (Owens). 2010;52:205-7 (Case rep.)

Microcirculation
Microcirculatory effects of local and remote ischemic preconditioning in supraceliac aortic clamping (Erling et al). 2010;52:1321-9

Monocytes
Perioperative platelet and monocyte activation in patients with critical limb ischemia (Burdess et al). 2010;52:697-703

Morbidity
Reducing venous stasis ulcers by fifty percent in 10 years: the next steps (Henke et al). 2010;52:S37-8

Mortality
An elevated neutrophil-lymphocyte ratio independently predicts mortality in chronic critical limb ischemia (Spark et al). 2010;52:632-6
Biomarkers in peripheral arterial disease patients and near- and longer-term mortality (Criqui et al). 2010;52:85-90
Leg strength predicts mortality in men but not in women with peripheral arterial disease (Singh et al). 2010;52:624-31
Risk factors for late mortality after endovascular repair of the thoracic aorta (Chung et al). 2010;52:549-55

Mouth protectors
Mandibular subluxation stabilized by mouthpiece for distal internal carotid artery exposure in carotid endarterectomy (Yoshino et al). 2010;52:1401-4

Multiple sclerosis
Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Lee). 2010;52:533 (Letter)
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Skurnick) (Letter); (Zamboni) (Reply). 2010;52:1749-50

Muscle, smooth, vascular
Inhibition of N-cadherin retards smooth muscle cell migration and intimal thickening via induction of apoptosis (Lyon et al). 2010;52:1301-9
Tissue factor pathway inhibitor-2 is induced by fluid shear stress in vascular smooth muscle cells and affects cell proliferation and survival (Ekstrand et al). 2010;52:167-75
Vascular smooth muscle cell peroxisome proliferator-activated receptor-γ deletion promotes abdominal aortic aneurysms (Hamblin et al). 2010;52:984-93

Mutation
JAK2 V617F mutation, mesenteric vein thrombosis, and myeloproliferative disorders (Owens). 2010;52:205-7 (Case rep.)

Mycotic aneurysm; *see* Aneurysm, infected

Myeloproliferative disorders
JAK2 V617F mutation, mesenteric vein thrombosis, and myeloproliferative disorders (Owens). 2010;52:205-7 (Case rep.)

Myocytes, smooth muscle
Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

N

Nanoparticles
Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20

Neck
Long-term follow-up of neck expansion after endovascular aortic aneurysm repair (Monahan et al). 2010;52:303-7
Negative-pressure wound therapy
Wound care: the role of advanced wound healing technologies (Wu et al). 2010;52:S59-66
Neurofibromatosis 1
Regarding "Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I" (Mendonça). 2010;52:1425 (Letter)
Neuroprotective agents
Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: results from a prospective randomized study (Stabile et al). 2010;52:1505-10
Neutrophils
An elevated neutrophil-lymphocyte ratio independently predicts mortality in chronic critical limb ischemia (Spark et al). 2010;52:632-6
Nitinol; *see* Alloys
Nitric oxide synthase type III
eNOS and *ACE* genes influence peripheral arterial disease predisposition in smokers (Sticchi et al). 2010;52:97-102, 97.e1
Nonagenarians; *see* Aged, 80 and over

O

Obesity
Impact of obesity on venous hemodynamics of the lower limbs (Willenberg et al). 2010;52:664-8
The influence of the obesity paradox and chronic kidney disease on long-term survival in a Japanese cohort with peripheral arterial disease (Kumakura et al). 2010;52:110-17
Octogenarians; *see* Aged, 80 and over
Onyx; *see* Dimethyl sulfoxide; Polyvinyls
Organ culture techniques
A novel cell permeant peptide inhibitor of MAPKAP kinase II inhibits intimal hyperplasia in a human saphenous vein organ culture model (Lopes et al). 2010;52:1596-607
Orthotic devices
Hull early walking aid for rehabilitation of transtibial amputees - randomized controlled trial (HEART) (Mazari et al). 2010;52:1564-71
Outcome and process assessment (health care)
The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization (Durham et al). 2010;52:600-7
Outpatients
Long-term efficacy and safety of once-daily enoxaparin plus warfarin for the outpatient ambulatory treatment of lower-limb deep vein thrombosis in the TROMBOTEK trial (Kurtoglu et al); (Henke) (Commentary). 2010;52:1262-71
Oxidative stress
Oxidized LDL in human carotid plaques is related to symptomatic carotid disease and lesion instability (Sigala et al). 2010;52:704-13
Oxidized LDL; *see* Lipoproteins, LDL

P

Pain
What you can't feel can hurt you (Boulton). 2010;52:S28-30
Patient care team
The costs of diabetic foot: the economic case for the limb salvage team (Driver et al). 2010;52:S17-22
History of the team approach to amputation prevention: pioneers and milestones (Sanders et al). 2010;52:S3-16
Required skills and interdisciplinary teams in starting a TEVAR practice (Starr). 2010;52:S45-9 (Editorial)
Rescuing Sisyphus: the team approach to amputation prevention (Mills Sr et al). 2010;52:S1-2 (Editorial)
Percutaneous transluminal angioplasty; *see* Angioplasty, balloon
Perfusion
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
Perfusion Imaging
Success of arterial revascularization determined by contrast ultrasound muscle perfusion imaging (Duerschmied et al). 2010;52:1531-6
Perioperative care
The impact of body mass index on perioperative outcomes of open and endovascular abdominal aortic aneurysm repair from the National Surgical Quality Improvement Program, 2005-2007 (Giles et al). 2010;52:1471-7
Perioperative platelet and monocyte activation in patients with critical limb ischemia (Burdess et al). 2010;52:697-703
Preoperative thrombolysis and venoplasty affords no benefit in patency following first rib resection and scalenectomy for subacute and chronic subclavian vein thrombosis (Guzzo et al). 2010;52:658-63
Process of care for carotid endarterectomy: perioperative medical management (Stoner and deFreitas). 2010;52:223-31 (Review)
Regarding "Preoperative statin therapy is associated with improved outcomes and resource utilization in patients undergoing aortic aneurysm repair" (Takagi et al) (Letter); (Stoner) (Reply). 2010;52:820-2
Symptomatic perioperative venous thromboembolism is a frequent complication in patients with a history of deep vein thrombosis (Liem et al). 2010;52:651-7
Peripheral arterial disease
ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52
Biomarkers in peripheral arterial disease patients and near- and longer-term mortality (Criqui et al). 2010;52:85-90
Combined effects of smoking and peripheral arterial disease on all-cause and cardiovascular disease mortality in a Chinese male cohort (Luo et al). 2010;51:673-8. Correction. 2010; 52:260
eNOS and *ACE* genes influence peripheral arterial disease predisposition in smokers (Sticchi et al). 2010;52:97-102, 97.e1
The influence of the obesity paradox and chronic kidney disease on long-term survival in a Japanese cohort with peripheral arterial disease (Kumakura et al). 2010;52:110-17
Interleukin-6 and interleukin-10 gene polymorphism, endothelial dysfunction, and postoperative prognosis in patients with peripheral arterial disease (Stoica et). 2010;52:103-9
Leg strength predicts mortality in men but not in women with peripheral arterial disease (Singh et al). 2010;52:624-31
Lessons learned from the analysis of gender effect on risk factors and procedural outcomes of lower extremity arterial disease (Vouyouka et al); (Bush) (Commentary). 2010;52: 1196-203

Noninvasive assessment of lower extremity hemodynamics in individuals with diabetes mellitus (Andersen). 2010;52:S76-80

Prognostic values of C-reactive protein levels on clinical outcome after endovascular therapy in hemodialysis patients with peripheral artery disease (Ishii et al). 2010;52:854-9

Results of the randomized, placebo-controlled clopidogrel and acetylsalicylic acid in bypass surgery for peripheral arterial disease (CASPAR) trial (Belch et al). 2010;52:825-33, 825.e1-e2

Younger women with symptomatic peripheral arterial disease are at increased risk of depressive symptoms (Smolderen et al). 2010;52:637-44

Peripheral vascular diseases

Laparoscopic transposition of the left renal vein into the inferior vena cava for nutcracker syndrome (Hartung et al). 2010;52:738-41 (Case rep.)

Regarding "Peripheral arterial interventions: Trends in market shareandoutcomesbyspecialty,1998-2005"(Sacks)(Letter); (Eslami) (Reply). 2010;52:528-9

Peritoneal cavity

Bioengineered vascular graft grown in the mouse peritoneal cavity (Song et al). 2010;52:994-1002, 994.e1-e2

Phenotype

Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

Haptoglobin 2-1 phenotype predicts rapid growth of abdominal aortic aneurysms (Wiernicki et al). 2010;52:691-6

Phenylbutyrates

Sodium 4-phenylbutyrate protects against spinal cord ischemia by inhibition of endoplasmic reticulum stress (Mizukami et al). 2010;52:1580-6

Physicians

Regarding "Good Samaritan statutes: a malpractice defense for doing the right thing" (Semel) (Letter); (Brown) (Reply). 2010;52:1125

Pitavastatin; *see* Quinolines

Plaque, atherosclerotic

Atherosclerotic plaque composition assessed by virtual histology intravascular ultrasound and cerebral embolization after carotid stenting (Timaran et al). 2010;52:1188-95

Evidence for markers of hypoxia and apoptosis in explanted human carotid atherosclerotic plaques (Bitto et al). 2010;52:1015-21

Histologic atherosclerotic plaque characteristics are associated with restenosis rates after endarterectomy of the common and superficial femoral arteries (Derksen et al). 2010;52:592-9

Plasmids

Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30

Polyethylene terephthalates

A contemporary meta-analysis of Dacron versus polytetrafluoroethylene grafts for femoropopliteal bypass grafting (Takagi et al). 2010;52:232-6 (Review)

Polymorphism, genetic

Interleukin-6 and interleukin-10 gene polymorphism, endothelial dysfunction, and postoperative prognosis in patients with peripheral arterial disease (Stoica et). 2010;52:103-9

Polytetrafluoroethylene

A contemporary meta-analysis of Dacron versus polytetrafluoroethylene grafts for femoropopliteal bypass grafting (Takagi et al). 2010;52:232-6 (Review)

Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease (McQuade et al). 2010;52:584-91, 584.e1-e8

Polyvinyls

Early side effects after embolization of a carotid body tumor using Onyx (Wiegand et al). 2010;52:742-5 (Case rep.)

Popliteal artery

An analysis of relationship between quality of life indices and clinical improvement following intervention in patients with intermittent claudication due to femoropopliteal disease (Mazari et al). 2010;52:77-84

A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Boufi et al). 2010;52:1211-17

A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Roddy). 2010;52:1417-18 (Editorial)

A contemporary meta-analysis of Dacron versus polytetrafluoroethylene grafts for femoropopliteal bypass grafting (Takagi et al). 2010;52:232-6 (Review)

Emergency endovascular treatment of early spontaneous nonaneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia (Trellopoulos et al). 2010;52:751-7 (Case rep.)

Entrapment of the popliteal artery (Tanaka et al). 2010;52:479

Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease (McQuade et al). 2010;52:584-91, 584.e1-e8

The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization (Durham et al). 2010;52:600-7

Limb salvage after delayed diagnosis for blunt traumatic infrapopliteal occlusion (Causey et al). 2010;52:734-7 (Case rep.)

Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent (Soga et al). 2010;52:608-15

Regarding "High spatial resolution magnetic resonance imaging of cystic adventitial disease of the popliteal artery" (Weimar et al) (Letter); (Hagspiel) (Reply). 2010;52:534-5

Subintimal angioplasty for femoro-popliteal occlusive disease (Markose et al). 2010;52:1410-16

Postoperative complications

Advanced thoracic endovascular aortic repair: late complications of thoracic endografts (Halandras and Milner). 2010;52:S100-2 (Editorial)

Endoleak management and postoperative surveillance following endovascular repair of thoracic aortic aneurysms (Ricotta). 2010;52:S91-9

Interleukin-6 and interleukin-10 gene polymorphism, endothelial dysfunction, and postoperative prognosis in patients with peripheral arterial disease (Stoica et). 2010;52:103-9

Liquid sclerotherapy for posttraumatic arteriovenous fistula of the radialis indicis artery (Katada et al). 2010;52:1343-5 (Case rep.)

Postthrombotic syndrome
Endovenectomy of the common femoral vein and intraoperative iliac vein recanalization for chronic iliofemoral venous occlusion (Comerota et al). 2010;52:243-7
Individually tailored duration of elastic compression therapy in relation to incidence of the postthrombotic syndrome (ten Cate-Hoek et al). 2010;52:132-8
Prevention and treatment of the postthrombotic syndrome (Henke). 2010;52:S21-8

PPAR gamma
Vascular smooth muscle cell peroxisome proliferator-activated receptor-γ deletion promotes abdominal aortic aneurysms (Hamblin et al). 2010;52:984-93

Practice guidelines as topic
Left subclavian artery revascularization: Society for Vascular Surgery® Practice Guidelines (Matsumura and Rizvi). 2010;52:S65-70 (Editorial)

Practice management, medical
A real world analysis of payment per unit time in a Maryland Vascular Practice (Martin et al); (O'Donnell) (Commentary). 2010;52:1094-100
A real world analysis of payment per unit time in a Maryland Vascular Practice (Roddy). 2010;52:1110 (Editorial)

Pregnancy
A durable iliac-axillary and axillary-carotid bypass for cerebral ischemia due to Takayasu arteritis allowed successful pregnancies and deliveries (Kondo et al). 2010;52:1713-5

Protein-serine-threonine kinases
A novel cell permeant peptide inhibitor of MAPKAP kinase II inhibits intimal hyperplasia in a human saphenous vein organ culture model (Lopes et al). 2010;52:1596-607

Pseudoaneurysm; *see* Aneurysm, false

Pulmonary embolism
Primary tibial vein aneurysm with recurrent pulmonary emboli (Gabrielli et al). 2010;52:464-6 (Case rep.)
Prospective randomized study comparing the clinical outcomes between inferior vena cava Greenfield and TrapEase filters (Usoh et al). 2010;52:394-9

Punctures
Banding between dialysis puncture sites to treat severe ischemic steal syndrome in low flow autogenous arteriovenous access (Shemesh et al). 2010;52:495-8
A ruptured aneurysm after stent graft puncture during computed tomography-guided thrombin injection (Tolenaar et al). 2010;52:1045-7 (Case rep.)
Vascular changes at the puncture segments of arteriovenous fistula for hemodialysis access (Hsiao et al). 2010;52:669-73

Q

Quality of life
An analysis of relationship between quality of life indices and clinical improvement following intervention in patients with intermittent claudication due to femoropopliteal disease (Mazari et al). 2010;52:77-84
Hull early walking aid for rehabilitation of transtibial amputees - randomized controlled trial (HEART) (Mazari et al). 2010;52:1564-71
Quality of life in patients with no-option critical limb ischemia underlines the need for new effective treatment (Sprengers et al). 2010;52:843-9, 843.e1

Quinolines
Nanoparticle-mediated endothelial cell-selective delivery of pitavastatin induces functional collateral arteries (therapeutic arteriogenesis) in a rabbit model of chronic hind limb ischemia (Oda et al). 2010;52:412-20

R

Radial artery
Liquid sclerotherapy for posttraumatic arteriovenous fistula of the radialis indicis artery (Katada et al). 2010;52:1343-5 (Case rep.)

Radiation dosage
The impact of radiation dose exposure during endovascular aneurysm repair on patient safety (Jones et al). 2010;52:298-302

Radiation injuries
The impact of radiation dose exposure during endovascular aneurysm repair on patient safety (Jones et al). 2010;52:298-302

Randomized controlled trials as topic
Inherent problems with randomized clinical trials with observational/no treatment arms (Buckley et al). 2010;52:237-41
Is a randomized trial necessary to determine whether endovascular repair is the preferred management strategy in patients with ruptured abdominal aortic aneurysms? (Veith et al); (Forbes and Ricco) (Commentary). 2010;52:1087-93
Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up (Christenson et al). 2010;52:1234-41
Therapy of isolated calf muscle vein thrombosis: a randomized, controlled study (Schwarz et al). 2010;52:1246-50

Reactive oxygen species
Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells (Lee et al); (Rzucidlo) (Commentary). 2010;52:1290-300

Recurrence
Venous ulcer diagnosis, treatment, and prevention of recurrences (Gillespie). 2010;52:S8-14

Registries
Restenosis after carotid endarterectomy in a multicenter regional registry (Goodney et al). 2010;52:897-905, 897.e1-e2

Renal artery
Accuracy of duplex sonography scans after renal artery stenting (Fleming et al). 2010;52:953-8
Laparoscopic treatment of renal artery entrapment (Bünger et al). 2010;52:1357-61 (Case rep.)
Late erosion of a prophylactic Celect IVC filter into the aorta, right renal artery, and duodenal wall (Becher et al). 2010;52:1041-4 (Case rep.)
Urgent interventional bilateral renal artery fenestration for giant pararenal abdominal aortic aneurysm with upper digestive tract obstruction (Ouyang and Liang). 2010;52:1048-51 (Case rep.)
Use of a surgeon-modified branched thoracic endograft to preserve an aortorenal bypass during treatment of an intercostal patch aneurysm (Jim et al). 2010;52:730-3 (Case rep.)

Visceral reconstruction techniques (Srivastava). 2010;52:S82-5 (Editorial)

Renal dialysis

Prognostic values of C-reactive protein levels on clinical outcome after endovascular therapy in hemodialysis patients with peripheral artery disease (Ishii et al). 2010;52:854-9

Prosthetic lower extremity haemodialysis access grafts have satisfactory patency despite high incidence of infection (Roddy). 2010;52:1723 (Editorial)

Prosthetic lower extremity hemodialysis access grafts have satisfactory patency despite a high incidence of infection (Geenen et al). 2010;52:1546-50

Revascularization of an occluded brachiocephalic vein using Outback-LTD re-entry catheter (Anil and Taneja). 2010;52:1038-40 (Case rep.)

Vascular changes at the puncture segments of arteriovenous fistula for hemodialysis access (Hsiao et al). 2010;52:669-73

Videothoracoscopic-guided management of a central vein perforation during hemodialysis catheter placement (Kuzniec et al). 2010;52:1354-6 (Case rep.)

Renal veins

Aorto-left renal vein fistula is a rare complication of abdominal aortic aneurysm with unique clinical presentation (Dragas et al). 2010;52:1658-61 (Case rep.)

Laparoscopic transposition of the left renal vein into the inferior vena cava for nutcracker syndrome (Hartung et al). 2010;52:738-41 (Case rep.)

Reperfusion injury

The effects of systemic hypothermia on a murine model of thoracic aortic ischemia reperfusion (Kang et al). 2010;52:435-43

Therapeutic distant organ effects of regional hypothermia during mesenteric ischemia-reperfusion injury (Santora et al). 2010;52:1003-14

Resuscitation orders

The ethical hierarchy of do not resuscitate orders: Never say never (Jones and McCullough). 2010;52:1384-6

Review

Acknowledgment of reviewers (Sidawy and Perler). 2010;52:1431-4 (Editorial)

Ribs

An improved method of exposure for transaxillary first rib resection (Illig). 2010;52:248-9

Preoperative thrombolysis and venoplasty affords no benefit in patency following first rib resection and scalenectomy for subacute and chronic subclavian vein thrombosis (Guzzo et al). 2010;52:658-63

Risk adjustment

CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection (Matsumura et al). 2010;52:576-83, 576.e1-e2

Evaluating parsimonious risk-adjustment models for comparing hospital outcomes with vascular surgery (Osborne et al). 2010;52:400-5

Risk factors

Analysis of risk factors for abdominal aortic aneurysm in a cohort of more than 3 million individuals (Kent et al); (Lilly) (Commentary). 2010;52:539-48

Asymptomatic internal carotid artery stenosis and cerebrovascular risk stratification (Nicolaides et al). 2010;52:1486-96, 1486.e1-e5

Lessons learned from the analysis of gender effect on risk factors and procedural outcomes of lower extremity arterial disease (Vouyouka et al); (Bush) (Commentary). 2010;52:1196-203

Risk factor analysis of thoracic endovascular repair using the Matsui-Kitamura stent graft for acute aortic emergencies in the descending thoracic aorta (Ohtake et al). 2010;52:1464-70

Risk factors for late mortality after endovascular repair of the thoracic aorta (Chung et al). 2010;52:549-55

Symptomatic perioperative venous thromboembolism is a frequent complication in patients with a history of deep vein thrombosis (Liem et al). 2010;52:651-7

Usefulness and limitations of computational models in aortic disease risk stratification (Beller et al). 2010;52:1572-9

Younger women with symptomatic peripheral arterial disease are at increased risk of depressive symptoms (Smolderen et al). 2010;52:637-44

RNA, small interfering

Endothelial cells are susceptible to rapid siRNA transfection and gene silencing ex vivo (Andersen et al). 2010;52:1608-15

Rupture

Regarding "Endovascular management of iliac rupture during endovascular aneurysm repair" (Semel et al). 2010;52:533 (Letter)

Spontaneous retroperitoneal hematoma associated with iliac vein rupture (Jiang et al). 2010;52:1278-82

S

Safety

The impact of radiation dose exposure during endovascular aneurysm repair on patient safety (Jones et al). 2010;52:298-302

Salmonella infections

Emergency endovascular treatment of early spontaneous nonaneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia (Trellopoulos et al). 2010;52:751-7 (Case rep.)

Salvage therapy

Complex subclavian artery pseudoaneurysm causing failure of endovascular stent repair with salvage by percutaneous thrombin injection (Lee et al). 2010;52:1058-60 (Case rep.)

Saphenous vein

Classification of proximal endovenous closure levels and treatment algorithm (Lawrence et al). 2010;52:388-93

Desmuslin gene knockdown causes altered expression of phenotype markers and differentiation of saphenous vein smooth muscle cells (Xiao et al). 2010;52:684-90

Elevated sex steroid hormones in great saphenous veins in men (Kendler et al); (Labropoulos) (Commentary). 2010;51:639-47. Correction. 2010;52:538

A novel cell permeant peptide inhibitor of MAPKAP kinase II inhibits intimal hyperplasia in a human saphenous vein organ culture model (Lopes et al). 2010;52:1596-607

Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up (Christenson et al). 2010;52:1234-41

A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous in-

sufficiency of the great saphenous vein (Gale et al). 2010; 52:645-50
A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Roddy). 2010;52: 796 (Editorial)
Regarding "The anatomy of the small saphenous vein: fascial and neural relations, saphenofemoral junction, and valves" (Caggiati) (Letter); (Schweighofer et al) (Reply). 2010;52: 1428-9

SCAI; *see* Society for Cardiac Angiography and Interventions

Scavenger receptors, class E
Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells (Lee et al); (Rzucidlo) (Commentary). 2010; 52:1290-300

Schools, medical
A vascular disease educational program in the preclinical years of medical school increases student interest in vascular disease (Godshall et al). 2010;52:775-81, 775.e1-e2

Sclerosis
A clinicopathologic study of immunoglobulin G4-related sclerosing disease of the thoracic aorta (Kasashima et al). 2010; 52:1587-95

Sclerotherapy
Liquid sclerotherapy for posttraumatic arteriovenous fistula of the radialis indicis artery (Katada et al). 2010;52:1343-5 (Case rep.)
Ultrasound-guided foam sclerotherapy is a safe and clinically effective treatment for superficial venous reflux (Bradbury et al). 2010;52:939-45

Scoliosis
Aortic rupture after spinal correction for scoliosis in the presence of a thoracic stent graft (Mani et al). 2010;52:1653-7 (Case rep.)

Septal occluder device
Transcatheter closure of recurrent aortic pseudoaneurysm previously treated by Amplatzer occluder device (Hussain et al). 2010;52:196-8 (Case rep.)

Severity of illness index
Revision of the venous clinical severity score: venous outcomes consensus statement: special communication of the American Venous Forum Ad Hoc Outcomes Working Group (Vasquez et al). 2010;52:1387-96

Sex factors
Mechanisms of gender-related outcome differences after carotid endarterectomy (den Hartog et al). 2010;52:1062-71, 1062.e1-e6 (Presidential Address)

SIR; *see* Society for Interventional Radiology

siRNA; *see* RNA, small interfering

Smoking
Combined effects of smoking and peripheral arterial disease on all-cause and cardiovascular disease mortality in a Chinese male cohort (Luo et al). 2010;51:673-8. Correction. 2010; 52:260
eNOS and *ACE* genes influence peripheral arterial disease predisposition in smokers (Sticchi et al). 2010;52:97-102, 97.e1

Smooth muscle cells; *see* Myocytes, smooth muscle

Society for Cardiac Angiography and Interventions
ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

Society for Interventional Radiology
ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

Society for Vascular Medicine
ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

Society for Vascular Nursing
ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

Society for Vascular Surgery
ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease (Olin et al). 2010;52:1616-52

Socioeconomic factors
The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization (Durham et al). 2010;52:600-7

Sodium 4-phenylbutyrate; *see* Phenylbutyrates

Specialties, medical
Regarding "Peripheral arterial interventions: Trends in market shareandoutcomesbyspecialty,1998-2005"(Sacks)(Letter); (Eslami) (Reply). 2010;52:528-9

Spinal cord
Aortic rupture after spinal correction for scoliosis in the presence of a thoracic stent graft (Mani et al). 2010;52:1653-7 (Case rep.)
Incidence, prevention, and management in spinal cord protection during TEVAR (Rizvi and Sullivan). 2010;52:S86-90 (Editorial)

Spinal cord ischemia
Sodium 4-phenylbutyrate protects against spinal cord ischemia by inhibition of endoplasmic reticulum stress (Mizukami et al). 2010;52:1580-6

Spinal cord vascular diseases
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Lee). 2010;52:533 (Letter)
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Skurnick) (Letter); (Zamboni) (Reply). 2010;52:1749-50

Statins; *see* Hydroxymethylglutaryl-CoA reductase inhibitors

Stem cells
Enhanced cell therapy strategy to treat chronic limb-threatening ischemia (Eton and Yu). 2010;52:199-204 (Case rep.)

Stenosis; *see* Constriction, pathologic

Stents
Accuracy of duplex sonography scans after renal artery stenting (Fleming et al). 2010;52:953-8
Aortic rupture after spinal correction for scoliosis in the presence of a thoracic stent graft (Mani et al). 2010;52:1653-7 (Case rep.)
Asymptomatic carotid artery stenosis-Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting (Schneider and Naylor); (Forbes) (Commentary) 2010;52:499-507. Correction. 2010;52:823
Asymptomatic carotid artery stenosis—Medical therapy alone versus medical therapy plus carotid endarterectomy or

stenting (Schneider and Naylor); (Forbes) (Commentary). 2010;52:499-507

Atherosclerotic plaque composition assessed by virtual histology intravascular ultrasound and cerebral embolization after carotid stenting (Timaran et al). 2010;52:1188-95

Below knee bare nitinol stent placement in high-risk patients with critical limb ischemia is still durable after 24 months of follow-up (Donas et al). 2010;52:356-61

CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection (Matsumura et al). 2010;52:576-83, 576.e1-e2

Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis (AbuRahma et al). 2010;52:1180-7

Cerebral reserve is decreased in elderly patients with carotid stenosis (Chaer et al). 2010;52:569-75

A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Boufi et al). 2010;52:1211-17

A comparison of the standard bolia technique versus subintimal recanalization plus Viabahn stent graft in the management of femoro-popliteal occlusions (Roddy). 2010;52:1417-18 (Editorial)

Complex subclavian artery pseudoaneurysm causing failure of endovascular stent repair with salvage by percutaneous thrombin injection (Lee et al). 2010;52:1058-60 (Case rep.)

Eight-year experience with carotid artery stenting for correction of symptomatic and asymptomatic post-endarterectomy defects. 2010;52:1511-7

Endovascular management of type B aortic dissections (DiMusto et al). 2010;52:S26-36

Endovascular repair of thoracic aortic traumatic transections is a safe method in patients with complicated injuries (Rahimi et al). 2010;52:891-6

Four-year randomized prospective comparison of percutaneous ePTFE/nitinol self-expanding stent graft versus prosthetic femoral-popliteal bypass in the treatment of superficial femoral artery occlusive disease (McQuade et al). 2010;52:584-91, 584.e1-e8

Heparin versus bivalirudin for carotid artery stenting using proximal endovascular clamping for neuroprotection: results from a prospective randomized study (Stabile et al). 2010;52:1505-10

Mechanisms to explain the poor results of carotid artery stenting (CAS) in symptomatic patients to date and options to improve CAS outcomes (Paraskevas et al). 2010;52:1367-75 (Review)

Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent (Soga et al). 2010;52:608-15

Outcomes of carotid stenting compared with endarterectomy are equivalent in asymptomatic patients and inferior in symptomatic patients (Giacovelli et al). 2010;52:906-13, 906.e1-e4

Patient-specific simulation in carotid artery stenting (Willaert et al). 2010;52:1700-5

Regarding "Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I" (Mendonça). 2010;52:1425 (Letter)

Risk factor analysis of thoracic endovascular repair using the Matsui-Kitamura stent graft for acute aortic emergencies in the descending thoracic aorta (Ohtake et al). 2010;52:1464-70

In situ fenestration in the aortic arch (Manning et al). 2010;52:491-4

Stent graft-induced new entry after endovascular repair for Stanford type B aortic dissection (Dong et al); (Ricotta) (Commentary). 2010;52:1450-8

Stroke and death after carotid endarterectomy and carotid artery stenting with and without high risk criteria (Giles et al). 2010;52:1497-504

Tips and techniques for optimal stent graft placement in angulated aneurysm necks (van Keulen et al). 2010;52:1081-6

Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques (Takahashi et al). 2010;52:1365-6

The value of the initial post-EVAR computed tomography angiography scan in predicting future secondary procedures using the Powerlink stent graft (Patel and Carpenter). 2010;52:1135-9

Stockings, compression

Individually tailored duration of elastic compression therapy in relation to incidence of the postthrombotic syndrome (ten Cate-Hoek et al). 2010;52:132-8

Inelastic bandages maintain their hemodynamic effectiveness over time despite significant pressure loss (Mosti and Partsch). 2010;52:925-31

Stroke

An unfavorable dietary pattern is associated with symptomatic ischemic stroke and carotid atherosclerosis (Mahe et al). 2010;52:62-8

CAPTURE 2 risk-adjusted stroke outcome benchmarks for carotid artery stenting with distal embolic protection (Matsumura et al). 2010;52:576-83, 576.e1-e2

Is carotid artery disease responsible for perioperative strokes after coronary artery bypass surgery? (Li et al); (Forbes and Ricco) (Commentary). 2010;52:1716-22

Stroke and death after carotid endarterectomy and carotid artery stenting with and without high risk criteria (Giles et al). 2010;52:1497-504

Subclavian artery

Complex subclavian artery pseudoaneurysm causing failure of endovascular stent repair with salvage by percutaneous thrombin injection (Lee et al). 2010;52:1058-60 (Case rep.)

Left subclavian artery revascularization: Society for Vascular Surgery® Practice Guidelines (Matsumura and Rizvi). 2010;52:S65-70 (Editorial)

Regarding "Right subclavian pseudoaneurysm secondary to blunt trauma in an arteriovenous malformation" (Lee) (Letter);(Quinones-Baldrich,) (Reply). 2010;52:257-8

The spectrum of arterial compression at the thoracic outlet (Criado et al). 2010;52:406-11

Subclavian vein

Aggressive treatment of idiopathic axillo-subclavian vein thrombosis provides excellent long-term function (Stone et al). 2010;52:127-31

Bilateral nonthrombotic subclavian vein obstruction causing upper extremity venous claudication (Thakur and Comerota). 2010;52:208-11 (Case rep.)

Superior mesenteric artery; *see* Mesenteric artery, superior

Surgical Ethics Challenges

The ethical hierarchy of do not resuscitate orders: Never say never (Jones and McCullough). 2010;52:1384-6

I know about Jack and you're no Jack Kevorkian (Jones and McCullough). 2010;52:489-90
Regarding "How do we guarantee trainee professional purity?" (Nakayama) (Letter); (Jones and McCullough) (Reply). 2010;52:257
There may be nothing new under the sun, but this is ridiculous (Brewster and Greisler). 2010;52:1697-8

Surgical procedures, elective
Midterm results of adjunctive neck therapies performed during elective infrarenal aortic aneurysm repair (Chung et al). 2010;52:1435-41

Surgical procedures, minimally invasive
Superficialization of arteriovenous fistulae employing minimally invasive liposuction (Causey et al). 2010;52:1397-400

Survivors
The influence of the obesity paradox and chronic kidney disease on long-term survival in a Japanese cohort with peripheral arterial disease (Kumakura et al). 2010;52:110-17

SVM; *see* Society for Vascular Medicine

SVN; *see* Society for Vascular Nursing

SVS; *see* Society for Vascular Surgery

Syncope
Deglutition syncope: a manifestation of vagal hyperactivity following carotid endarterectomy (Endean et al). 2010;52:720-2 (Case rep.)

T

Takayasu arteritis
A durable iliac-axillary and axillary-carotid bypass for cerebral ischemia due to Takayasu arteritis allowed successful pregnancies and deliveries (Kondo et al). 2010;52:1713-5

Thigh
Anatomical variations of the femoral vein (Uhl et al). 2010;52:714-19

Thoracic aortic aneurysm; *see* Aortic aneurysm, thoracic

Thoracic injuries
Aortic dilatation after endovascular repair of blunt traumatic thoracic aortic injuries (Forbes et al). 2010;52:45-8
Aortic endograft sizing in trauma patients with hemodynamic instability (Jonker et al). 2010;52:39-44
Economic evaluation of open vs endovascular repair of blunt traumatic thoracic aortic injuries (Tong et al). 2010;52:31-8

Thoracic outlet syndrome
The spectrum of arterial compression at the thoracic outlet (Criado et al). 2010;52:406-11

Thoracic surgery, video-assisted
Videothoracoscopic-guided management of a central vein perforation during hemodialysis catheter placement (Kuzniec et al). 2010;52:1354-6 (Case rep.)

Thrombin
Complex subclavian artery pseudoaneurysm causing failure of endovascular stent repair with salvage by percutaneous thrombin injection (Lee et al). 2010;52:1058-60 (Case rep.)
A ruptured aneurysm after stent graft puncture during computed tomography-guided thrombin injection (Tolenaar et al). 2010;52:1045-7 (Case rep.)

Thrombolysis; *see* Thrombolytic therapy

Thrombolytic therapy
Agitation thrombolysis for fresh iatrogenic IVC thrombosis in patients with Budd-Chiari syndrome (Ding et al). 2010;52:782-4
Preoperative thrombolysis and venoplasty affords no benefit in patency following first rib resection and scalenectomy for subacute and chronic subclavian vein thrombosis (Guzzo et al). 2010;52:658-63
Thrombolysis for acute arterial occlusion (van den Berg). 2010;52:512-15

Thrombosis
Thrombotic mesenteric ischemia due to aortic dissection (Bao and Chen). 2010;52:1674, 1674.e1-e2
Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction (Li et al). 2010;52:1242-5

Tibia
Primary tibial vein aneurysm with recurrent pulmonary emboli (Gabrielli et al). 2010;52:464-6 (Case rep.)

Tibial arteries
Outcomes of selective tibial artery repair following combat-related extremity injury (Burkhardt et al). 2010;52:91-6
Predictors of failure and success of tibial interventions for critical limb ischemia (Fernandez et al). 2010;52:834-42

Ticlopidine
Results of the randomized, placebo-controlled clopidogrel and acetylsalicylic acid in bypass surgery for peripheral arterial disease (CASPAR) trial (Belch et al). 2010;52:825-33, 825.e1-e2

Tissue engineering
Animal models for the assessment of novel vascular conduits (Byrom et al). 2010;52:176-95
Bioengineered vascular graft grown in the mouse peritoneal cavity (Song et al). 2010;52:994-1002, 994.e1-e2
Wound care: the role of advanced wound healing technologies (Wu et al). 2010;52:S59-66

Tissue factor pathway inhibitor-2; *see* Glycoproteins

Tissues
Upper extremity ischemia treated with tissue repair cells from adult bone marrow (Comerota et al). 2010;52:723-9 (Case rep.)

Toes
Toe and flow: essential components and structure of the amputation prevention team (Rogers et al). 2010;52:S23-7

Tomography, X-ray computed
Intraprocedural imaging: flat panel detectors, rotational angiography, FluoroCT, IVUS, or still the portable C-arm? (Eagleton). 2010;52:S50-9 (Editorial)
A ruptured aneurysm after stent graft puncture during computed tomography-guided thrombin injection (Tolenaar et al). 2010;52:1045-7 (Case rep.)
The value of the initial post-EVAR computed tomography angiography scan in predicting future secondary procedures using the Powerlink stent graft (Patel and Carpenter). 2010;52:1135-9

Transfection
Endothelial cells are susceptible to rapid siRNA transfection and gene silencing ex vivo (Andersen et al). 2010;52:1608-15

Transglutaminases
Activation of transglutaminase type 2 for aortic wall protection in a rat abdominal aortic aneurysm formation (Munezane et al). 2010;52:967-74

Transplantation, homologous
Eight-year experience with cryopreserved arterial homografts for the in situ reconstruction of abdominal aortic infections (Roddy). 2010;52:516 (Editorial)
Trauma; *see* Wounds and injuries
Trauma centers
A contemporary rural trauma center experience in blunt traumatic aortic injury (Durham et al). 2010;52:884-90
Treatment outcome
Carotid artery stenting outcomes are equivalent to carotid endarterectomy outcomes for patients with post-carotid endarterectomy stenosis (AbuRahma et al). 2010;52:1180-7
Challenges of distal bypass surgery in patients with diabetes: patient selection, techniques, and outcomes (Conte). 2010;52:S96-103
Evaluating parsimonious risk-adjustment models for comparing hospital outcomes with vascular surgery (Osborne et al). 2010;52:400-5
Finnvasc score and modified Prevent III score predict long-term outcome after infrainguinal surgical and endovascular revascularization for critical limb ischemia (Arvela et al). 2010;52:1218-25
The impact of body mass index on perioperative outcomes of open and endovascular abdominal aortic aneurysm repair from the National Surgical Quality Improvement Program, 2005-2007 (Giles et al). 2010;52:1471-7
Important observations made managing carotid body tumors during a 25-year experience (Kruger et al); (LaMuraglia) (Commentary). 2010;52:1518-24
Intermediate-term EVAR outcomes in octogenarians (Fonseca et al). 2010;52:556-61
Is carotid artery disease responsible for perioperative strokes after coronary artery bypass surgery? (Li et al); (Forbes and Ricco) (Commentary). 2010;52:1716-22
Lessons learned from the analysis of gender effect on risk factors and procedural outcomes of lower extremity arterial disease (Vouyouka et al); (Bush) (Commentary). 2010;52:1196-203
Long-term outcomes of diabetic patients undergoing endovascular infrainguinal interventions (Abularrage et al). 2010;52:314-22, 314.e1-e4
Long-term outcomes of secondary procedures after endovascular aneurysm repair (Mehta et al). 2010;52:1442-9
Long-term outcomes of surgical aortic fenestration for complicated acute type B aortic dissections (Trimarchi et al). 2010;52:261-6
Mechanisms of gender-related outcome differences after carotid endarterectomy (den Hartog et al). 2010;52:1062-71, 1062.e1-e6 (Presidential Address)
Mechanisms to explain the poor results of carotid artery stenting (CAS) in symptomatic patients to date and options to improve CAS outcomes (Paraskevas et al). 2010;52:1367-75 (Review)
Midterm outcome of endovascular repair of ruptured isolated iliac artery aneurysms (Hechelhammer et al). 2010;52:1159-63
Outcomes before and after initiation of an acute aortic treatment center (Davies et al); (Taylor) (Commentary). 2010;52:1478-85
Outcomes in open repair of the thoracic and thoracoabdominal aorta (Acher and Wynn). 2010;52:S3-9
Outcomes of interventions for recurrent disease after endoluminal intervention for superficial femoral artery disease (Davies et al). 2010;52:331-9, 331.e1-e2
Outcomes of planned celiac artery coverage during TEVAR (Mehta et al). 2010;52:1153-8
Outcomes of reinterventions after subintimal angioplasty (Schmieder et al). 2010;52:375-82
Outcomes of selective tibial artery repair following combat-related extremity injury (Burkhardt et al). 2010;52:91-6
Outcomes of symptomatic abdominal aortic aneurysm repair (De Martino et al). 2010;52:5-12
Poor outcomes with cryoplasty for lower extremity arterial occlusive disease (Schmieder et al). 2010;52:362-8
Prognostic values of C-reactive protein levels on clinical outcome after endovascular therapy in hemodialysis patients with peripheral artery disease (Ishii et al). 2010;52:854-9
Regarding "Long-term outcomes and resource utilization of endovascular versus open repair of abdominal aortic aneurysms in Ontario" (Takagi et al). 2010;52:818-19 (Letter)
Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010;52:1425-7
Revision of the venous clinical severity score: venous outcomes consensus statement: special communication of the American Venous Forum Ad Hoc Outcomes Working Group (Vasquez et al). 2010;52:1387-96
Significant sac retraction after endovascular aneurysm repair is a robust indicator of durable treatment success (Houbballah et al). 2010;52:878-83
Twelve-year outcomes after endovascular aneurysm repair using earliest available endograft components from a single center (Adiseshiah et al). 2010;52:49-54
TRPV cation channels
Mechanosensitive transient receptor potential vanilloid type 1 channels contribute to vascular remodeling of rat fistula veins (Chen et al). 2010;52:1310-20
Tuberculosis
Tuberculosis should not be ignored in patients with peripheral gangrene (Yao et al). 2010;52:1662-4 (Case rep.)

U

Ulcer
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
Ultrasonography
Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9
Success of arterial revascularization determined by contrast ultrasound muscle perfusion imaging (Duerschmied et al). 2010;52:1531-6
Ultrasound-guided foam sclerotherapy is a safe and clinically effective treatment for superficial venous reflux (Bradbury et al). 2010;52:939-45
Ultrasonography, Doppler, duplex
Accuracy of duplex sonography scans after renal artery stenting (Fleming et al). 2010;52:953-8
Clinical course of asymptomatic patients with carotid duplex scan end diastolic velocities of 100 to 124 centimeters per second (Riles et al). 2010;52:914-19, 914.e1

Duplex ultrasound factors predicting persistent type II endoleak and increasing AAA sac diameter after EVAR (Beeman et al). 2010;52:1147-52
Identifying the source of superficial reflux in venous leg ulcers using duplex ultrasound (Obermayer and Garzon). 2010;52:1255-61
Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al). 2010;52:69-76
Juxtaluminal hypoechoic area in ultrasonic images of carotid plaques and hemispheric symptoms (Griffin et al) 2010;52:69-76. Correction. 2010;52:1430
Period frequency of iliofemoral venous occlusive disease by Doppler ultrasound and corresponding treatment in a tertiary care facility (Crisostomo et al). 2010;52:1272-7
Practical use of duplex echo-guided recanalization of chronic total occlusion in the iliac artery (Kawarada et al). 2010;52:475-8 (Case rep.)

Ultrasonography, interventional
Atherosclerotic plaque composition assessed by virtual histology intravascular ultrasound and cerebral embolization after carotid stenting (Timaran et al). 2010;52:1188-95
Intraprocedural imaging: flat panel detectors, rotational angiography, FluoroCT, IVUS, or still the portable C-arm? (Eagleton). 2010;52:S50-9 (Editorial)

United States Food and Drug Administration
Description and comparison of Food and Drug Administration-approved thoracic endovascular aneurysm devices (Lyden). 2010;52:S10-14

Upper extremity
Bilateral nonthrombotic subclavian vein obstruction causing upper extremity venous claudication (Thakur and Comerota). 2010;52:208-11 (Case rep.)
Superficialization of arteriovenous fistulae employing minimally invasive liposuction (Causey et al). 2010;52:1397-400
Upper extremity ischemia treated with tissue repair cells from adult bone marrow (Comerota et al). 2010;52:723-9 (Case rep.)

Urinary bladder
Urinary bladder injuries during vascular surgery (Nakamura et al). 2010;52:453-5 (Case rep.)

V

Vagus nerve
Deglutition syncope: a manifestation of vagal hyperactivity following carotid endarterectomy (Endean et al). 2010;52:720-2 (Case rep.)

Varicose ulcer
Identifying the source of superficial reflux in venous leg ulcers using duplex ultrasound (Obermayer and Garzon). 2010;52:1255-61
Non-medical initiatives to decrease venous ulcers prevalence (Passman). 2010;52:S29-36
The Pacific Vascular Symposium 6: the Venous Ulcer Summit in perspective (Henke). 2010;52:S1-2 (Editorial)
Prevention and treatment of venous ulcers in primary chronic venous insufficiency (Neglen). 2010;52:S15-20
Reducing venous stasis ulcers by fifty percent in 10 years: the next steps (Henke et al). 2010;52:S37-8
Venous ulcer diagnosis, treatment, and prevention of recurrences (Gillespie). 2010;52:S8-14

Varicose veins
Elevated sex steroid hormones in great saphenous veins in men (Kendler et al); (Labropoulos) (Commentary). 2010;51:639-47. Correction. 2010;52:538
Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up (Christenson et al). 2010;52:1234-41

Vascular diseases
Reducing venous stasis ulcers by fifty percent in 10 years: the next steps (Henke et al). 2010;52:S37-8
The structure and processes of the Pacific Vascular Symposium 6 (Lurie et al). 2010;52:S3-7, S3.e1-e4
A vascular disease educational program in the preclinical years of medical school increases student interest in vascular disease (Godshall et al). 2010;52:775-81, 775.e1-e2

Vascular endothelial growth factor A
Vascular endothelial growth factor enhances angiotensin II-induced aneurysm formation in apolipoprotein E-deficient mice (Choke et al). 2010;52:159-66, 159.e1

Vascular Images
Collateral pathways visualization of the innominate vein (Civilini et al). 2010;52:758-9
Entrapment of the popliteal artery (Tanaka et al). 2010;52:479
A giant femoral artery pseudoaneurysm in an infant boy (Park et al). 2010;52:222
Persistence of intra-aortic air after circulatory arrest (Nakamura et al). 2010;52:1061
Thrombotic mesenteric ischemia due to aortic dissection (Bao and Chen). 2010;52:1674, 1674.e1-e2
Total aortic repair in Marfan syndrome using stent grafting with hybrid techniques (Takahashi et al). 2010;52:1365-6

Vascular patency
Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent (Soga et al). 2010;52:608-15

Vascular smooth muscle; *see* Muscle, smooth, vascular

Vascular surgical procedures
Access site management with vascular closure devices for percutaneous transarterial procedures (Bechara et al). 2010;52:1682-96 (Review)
Analysis of large databases in vascular surgery (Nguyen and Barshes). 2010;52:768-74 (Review)
An improved method of exposure for transaxillary first rib resection (Illig). 2010;52:248-9
Aortic dilatation after endovascular repair of blunt traumatic thoracic aortic injuries (Forbes et al). 2010;52:45-8
Associated factors, timing, and technical aspects of late failure following open surgical aneurysm repairs (Coscas et al). 2010;52:272-81
Challenges of distal bypass surgery in patients with diabetes: patient selection, techniques, and outcomes (Conte). 2010;52:S96-103
Classification of proximal endovenous closure levels and treatment algorithm (Lawrence et al). 2010;52:388-93
Conduits and endoconduits, percutaneous access (Peterson). 2010;52:S60-4
A contemporary experience of open aortic reconstruction in patients with chronic atherosclerotic occlusion of the abdominal aorta (West et al). 2010;52:1164-72
Current assessment of endovascular therapy for infrainguinal arterial occlusive disease in patients with diabetes (Ihnat and Mills). 2010;52:S92-5

Current Procedural Terminology (CPT) coding for endovascular intervention in the descending thoracic aorta (Seabrook). 2010;52:S103-6 (Editorial)

Economic evaluation of open vs endovascular repair of blunt traumatic thoracic aortic injuries (Tong et al). 2010;52:31-8

Emergency endovascular treatment of early spontaneous nonaneurysmal popliteal artery rupture in a patient with *Salmonella* bacteremia (Trellopoulos et al). 2010;52:751-7 (Case rep.)

Endoleak management and postoperative surveillance following endovascular repair of thoracic aortic aneurysms (Ricotta). 2010;52:S91-9

Endovascular and open surgery for acute occlusion of the superior mesenteric artery (Block et al). 2010;52:959-66

Endovascular aneurysm repair for ruptured abdominal aortic aneurysm: how I do it (Mehta). 2010;52:1706-12

Endovascular aneurysm repair in nonagenarians is safe and effective (Goldstein et al); (Bush) (Commentary). 2010;52:1140-6

Endovascular aneurysm repair is superior to open surgery for ruptured abdominal aortic aneurysms in EVAR-suitable patients (Ten Bosch et al). 2010;52:13-18

Endovascular procedures for aorto-iliac occlusive disease are associated with superior short-term clinical and economic outcomes compared with open surgery in the inpatient population (Indes et al). 2010;52:1173-9, 1173.e1

Endovenectomy of the common femoral vein and intraoperative iliac vein recanalization for chronic iliofemoral venous occlusion (Comerota et al). 2010;52:243-7

Evaluating parsimonious risk-adjustment models for comparing hospital outcomes with vascular surgery (Osborne et al). 2010;52:400-5

Existing risk prediction methods for elective abdominal aortic aneurysm repair do not predict short-term outcome following endovascular repair (Patterson et al). 2010;52:25-30

Fenestrated and branched devices in the pipeline (Greenberg and Qureshi). 2010;52:S15-21

Finnvasc score and modified Prevent III score predict long-term outcome after infrainguinal surgical and endovascular revascularization for critical limb ischemia (Arvela et al). 2010;52:1218-25

The first operation on the profunda femoris artery (Yao). 2010;52:1101-2

Halsted-Holman vascular trauma legacy (Rich et al). 2010;52:508-11

I know about Jack and you're no Jack Kevorkian (Jones and McCullough). 2010;52:489-90

The impact of body mass index on perioperative outcomes of open and endovascular abdominal aortic aneurysm repair from the National Surgical Quality Improvement Program, 2005-2007 (Giles et al). 2010;52:1471-7

The impact of radiation dose exposure during endovascular aneurysm repair on patient safety (Jones et al). 2010;52:298-302

The impact of socioeconomic factors on outcome and hospital costs associated with femoropopliteal revascularization (Durham et al). 2010;52:600-7

Incidence, prevention, and management in spinal cord protection during TEVAR (Rizvi and Sullivan). 2010;52:S86-90 (Editorial)

Incorporating simulation in vascular surgery education (Bismuth et al). 2010;52:1072-80

The influence of aneurysm size on anatomic suitability for endovascular repair (Keefer et al). 2010;52:873-7

The influence of Dominique Jean Larrey on the art and science of amputations (Welling et al). 2010;52:790-3

Is a randomized trial necessary to determine whether endovascular repair is the preferred management strategy in patients with ruptured abdominal aortic aneurysms? (Veith et al); (Forbes and Ricco) (Commentary). 2010;52:1087-93

Juxtarenal aortic aneurysm repair (Jongkind et al). 2010;52:760-7 (Review)

Left subclavian artery revascularization: Society for Vascular Surgery® Practice Guidelines (Matsumura and Rizvi). 2010;52:S65-70 (Editorial)

The limitations of thoracic endovascular aortic repair in altering the natural history of blunt aortic injury (Lang et al). 2010;52:290-7

Long-term follow-up of neck expansion after endovascular aortic aneurysm repair (Monahan et al). 2010;52:303-7

Long-term outcomes of diabetic patients undergoing endovascular infrainguinal interventions (Abularrage et al). 2010;52:314-22, 314.e1-e4

Long-term outcomes of secondary procedures after endovascular aneurysm repair (Mehta et al). 2010;52:1442-9

Long-term outcomes of surgical aortic fenestration for complicated acute type B aortic dissections (Trimarchi et al). 2010;52:261-6

Midterm outcome of endovascular repair of ruptured isolated iliac artery aneurysms (Hechelhammer et al). 2010;52:1159-63

Outcomes in open repair of the thoracic and thoracoabdominal aorta (Acher and Wynn). 2010;52:S3-9

Outcomes of interventions for recurrent disease after endoluminal intervention for superficial femoral artery disease (Davies et al). 2010;52:331-9, 331.e1-e2

Outcomes of reinterventions after subintimal angioplasty (Schmieder et al). 2010;52:375-82

Outcomes of symptomatic abdominal aortic aneurysm repair (De Martino et al). 2010;52:5-12

The Pacific Vascular Symposium 6: the Venous Ulcer Summit in perspective (Henke). 2010;52:S1-2 (Editorial)

A pilot, prospective evaluation of a direct thrombin inhibitor, bivalirudin (Angiomax), in patients undergoing lower extremity bypass (Kashyap et al). 2010;52:369-74

Preoperative thrombolysis and venoplasty affords no benefit in patency following first rib resection and scalenectomy for subacute and chronic subclavian vein thrombosis (Guzzo et al). 2010;52:658-63

Prior endovascular abdominal aortic aneurysm repair provides no survival benefits when the aneurysm ruptures (Cho et al). 2010;52:1127-34

Prognostic values of C-reactive protein levels on clinical outcome after endovascular therapy in hemodialysis patients with peripheral artery disease (Ishii et al). 2010;52:854-9

A real world analysis of payment per unit time in a Maryland Vascular Practice (Martin et al); (O'Donnell) (Commentary). 2010;52:1094-100

A real world analysis of payment per unit time in a Maryland Vascular Practice (Roddy). 2010;52:1110 (Editorial)

Regarding "A closer look at meta-analyses of observational data" (Navarese et al) (Letter); (Jonker et al) (Reply). 2010;52:819-20

Regarding "Endovascular management of iliac rupture during

endovascular aneurysm repair" (Semel et al). 2010;52:533 (Letter)
Regarding "Long-term outcomes and resource utilization of endovascular versus open repair of abdominal aortic aneurysms in Ontario" (Takagi et al). 2010;52:818-19 (Letter)
Regarding "Perioperative outcomes and amputation-free survival after lower extremity bypass surgery in California hospitals" (Takagi et al) (Letter); (Feinglass) (Reply). 2010;52:1425-7
Regarding "Preoperative statin therapy is associated with improved outcomes and resource utilization in patients undergoing aortic aneurysm repair" (Takagi et al) (Letter); (Stoner) (Reply). 2010;52:820-2
Regulatory TEVAR clinical trials (Garcia-Toca and Eskandari). 2010;52:S22-5 (Editorial)
Reporting standards for thoracic endovascular aortic repair (TEVAR) (Fillinger et al). 2010;52:1022-33, 1022.e1-e5
Required skills and interdisciplinary teams in starting a TEVAR practice (Starr). 2010;52:S45-9 (Editorial)
Risk factors for late mortality after endovascular repair of the thoracic aorta (Chung et al). 2010;52:549-55
Seventeen-year follow-up after ascending-to-infrarenal aorta bypass for recurrent coarctation in an adult (Pereda et al). 2010;52:1362-4 (Case rep.)
Staged total exclusion of the aorta for chronic type B aortic dissection (Dias Perera et al). 2010;52:1339-42 (Case rep.)
A step-wise approach for surgical management of diabetic foot infections (Fisher et al). 2010;52:S72-5
The structure and processes of the Pacific Vascular Symposium 6 (Lurie et al). 2010;52:S3-7, S3.e1-e4
Successful endovascular treatment of multiple mesenteric arterial aneurysms associated with alpha-1-antitrypsin deficiency (Gordon et al). 2010;52:467-70 (Case rep.)
Success of arterial revascularization determined by contrast ultrasound muscle perfusion imaging (Duerschmied et al). 2010;52:1531-6
A systematic review of endovascular treatment of extensive aortoiliac occlusive disease (Jongkind et al). 2010;52:1376-83 (Review)
Thoracic aortic endovascular repair for mycotic aneurysms and fistulas (Patel et al). 2010;52:S37-40 (Editorial)
Toe and flow: essential components and structure of the amputation prevention team (Rogers et al). 2010;52:S23-7
Urinary bladder injuries during vascular surgery (Nakamura et al). 2010;52:453-5 (Case rep.)
The Vascular Study Group of New England Cardiac Risk Index (VSG-CRI) predicts cardiac complications more accurately than the Revised Cardiac Risk Index in vascular surgery patients (Bertges et al). 2010;52:674-83, 674.e1-e3

Vascular system injuries
History of temporary intravascular shunts in the management of vascular injury (Hancock et al). 2010;52:1405-9

Veins
Arm vein conduit vs prosthetic graft in infrainguinal revascularization for critical leg ischemia (Arvela et al). 2010;52:616-23; Roddy). 2010;52:794-5 (Editorial)
Don't trust a vein graft to treat carotid aneurysm in patients with Behçet disease (Berard et al). 2010;52:471-4 (Case rep.)
Mechanosensitive transient receptor potential vanilloid type 1 channels contribute to vascular remodeling of rat fistula veins (Chen et al). 2010;52:1310-20
Primary balloon angioplasty plus balloon angioplasty maturation to upgrade small-caliber veins (<3 mm) for arteriovenous fistulas (De Marco Garcia et al). 2010;52:139-44
Primary tibial vein aneurysm with recurrent pulmonary emboli (Gabrielli et al). 2010;52:464-6 (Case rep.)
Rationale and practical techniques for mouse models of early vein graft adaptations (Yu et al). 2010;52:444-52
Videothoracoscopic-guided management of a central vein perforation during hemodialysis catheter placement (Kuzniec et al). 2010;52:1354-6 (Case rep.)

Vena cava filters
Cost-effectiveness of guidelines for insertion of inferior vena cava filters in high-risk trauma patients (Spangler et al). 2010;52:1537-45, 1537.e1-e2
A design modification to minimize tilting of an inferior vena cava filter does not deliver a clinical benefit (Shelgikar et al). 2010;52:920-4
Late erosion of a prophylactic Celect IVC filter into the aorta, right renal artery, and duodenal wall (Becher et al). 2010;52:1041-4 (Case rep.)
Prospective randomized study comparing the clinical outcomes between inferior vena cava Greenfield and TrapEase filters (Usoh et al). 2010;52:394-9
Trends in vena caval interruption (Moore et al); (Smith) (Commentary). 2010;52:118-26, 118.e1-e3
Trends in vena caval interruption (Roddy). 2010;52:251 (Editorial)

Vena cava, inferior
Agitation thrombolysis for fresh iatrogenic IVC thrombosis in patients with Budd-Chiari syndrome (Ding et al). 2010;52:782-4
Cost-effectiveness of guidelines for insertion of inferior vena cava filters in high-risk trauma patients (Spangler et al). 2010;52:1537-45, 1537.e1-e2
A design modification to minimize tilting of an inferior vena cava filter does not deliver a clinical benefit (Shelgikar et al). 2010;52:920-4
Laparoscopic transposition of the left renal vein into the inferior vena cava for nutcracker syndrome (Hartung et al). 2010;52:738-41 (Case rep.)
Late erosion of a prophylactic Celect IVC filter into the aorta, right renal artery, and duodenal wall (Becher et al). 2010;52:1041-4 (Case rep.)
Prospective randomized study comparing the clinical outcomes between inferior vena cava Greenfield and TrapEase filters (Usoh et al). 2010;52:394-9
Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction (Li et al). 2010;52:1242-5

Venous disease; *see* Vascular diseases

Venous insufficiency
Angioplasty of central venous stenosis in patients with multiple sclerosis with ultrasound evidence of chronic cerebrospinal venous insufficiency may be a sham (Requarth) (Letter); (Zamboni et al) (Reply). 2010;52:1748-9
Prevention and treatment of venous ulcers in primary chronic venous insufficiency (Neglen). 2010;52:S15-20
A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Gale et al). 2010;52:645-50
A randomized, controlled trial of endovenous thermal ablation using the 810-nm wavelength laser and the ClosurePLUS

radiofrequency ablation methods for superficial venous insufficiency of the great saphenous vein (Roddy). 2010;52:796 (Editorial)
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Lee). 2010;52:533 (Letter)
Regarding "A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency" (Skurnick) (Letter); (Zamboni) (Reply). 2010;52:1749-50
Revision of the venous clinical severity score: venous outcomes consensus statement: special communication of the American Venous Forum Ad Hoc Outcomes Working Group (Vasquez et al). 2010;52:1387-96
Ultrasound-guided foam sclerotherapy is a safe and clinically effective treatment for superficial venous reflux (Bradbury et al). 2010;52:939-45

Venous thromboembolism
Symptomatic perioperative venous thromboembolism is a frequent complication in patients with a history of deep vein thrombosis (Liem et al). 2010;52:651-7
Trends in vena caval interruption (Moore et al); (Smith) (Commentary). 2010;52:118-26, 118.e1-e3
Trends in vena caval interruption (Roddy). 2010;52:251 (Editorial)

Venous thrombosis
Aggressive treatment of idiopathic axillo-subclavian vein thrombosis provides excellent long-term function (Stone et al). 2010;52:127-31
Agitation thrombolysis for fresh iatrogenic IVC thrombosis in patients with Budd-Chiari syndrome (Ding et al). 2010;52:782-4
Comparison of the clinical history of symptomatic isolated muscular calf vein thrombosis versus deep calf vein thrombosis (Galanaud et al). 2010;52:932-8, 932.e1-e2
Iliofemoral venous thrombosis from external compression by a vesical diverticulum (Gupta et al). 2010;52:1671-3 (Case rep.)
Individually tailored duration of elastic compression therapy in relation to incidence of the postthrombotic syndrome (ten Cate-Hoek et al). 2010;52:132-8
JAK2 V617F mutation, mesenteric vein thrombosis, and myeloproliferative disorders (Owens). 2010;52:205-7 (Case rep.)
Long-term efficacy and safety of once-daily enoxaparin plus warfarin for the outpatient ambulatory treatment of lower-limb deep vein thrombosis in the TROMBOTEK trial (Kurtoglu et al); (Henke) (Commentary). 2010;52:1262-71
Management of isolated soleal and gastrocnemius vein thrombosis (Sales et al). 2010;52:1251-4
Period frequency of iliofemoral venous occlusive disease by Doppler ultrasound and corresponding treatment in a tertiary care facility (Crisostomo et al). 2010;52:1272-7
Preoperative thrombolysis and venoplasty affords no benefit in patency following first rib resection and scalenectomy for subacute and chronic subclavian vein thrombosis (Guzzo et al). 2010;52:658-63
Prevention and treatment of the postthrombotic syndrome (Henke). 2010;52:S21-8
Prospective randomized study comparing the clinical outcomes between inferior vena cava Greenfield and TrapEase filters (Usoh et al). 2010;52:394-9
Symptomatic perioperative venous thromboembolism is a frequent complication in patients with a history of deep vein thrombosis (Liem et al). 2010;52:651-7
Therapy of isolated calf muscle vein thrombosis: a randomized, controlled study (Schwarz et al). 2010;52:1246-50

Venous ulcer; *see* Varicose ulcer

Venous valves
Regarding "The anatomy of the small saphenous vein: fascial and neural relations, saphenofemoral junction, and valves" (Caggiati) (Letter); (Schweighofer et al) (Reply). 2010;52:1428-9

Vertebral artery dissection
Bilateral internal carotid and vertebral artery dissection after a horse-riding injury (Keilani et al). 2010;52:1052-7 (Case rep.)

Videothoracoscopy; *see* Thoracic surgery, video-assisted

W

Walking
Multicenter randomized clinical trial of supervised exercise therapy with or without feedback versus walking advice for intermittent claudication (Nicolaï et al). 2010;52:348-55

War
Outcomes of selective tibial artery repair following combat-related extremity injury (Burkhardt et al). 2010;52:91-6

Warfarin
The effect of warfarin therapy on endoleak development after endovascular aneurysm repair (EVAR) of the abdominal aorta (Bobadilla et al). 2010;52:267-71
Long-term efficacy and safety of once-daily enoxaparin plus warfarin for the outpatient ambulatory treatment of lower-limb deep vein thrombosis in the TROMBOTEK trial (Kurtoglu et al); (Henke) (Commentary). 2010;52:1262-71
Warfarin anticoagulation before angioplasty relieves thrombus burden in Budd-Chiari syndrome caused by inferior vena cava anatomic obstruction (Li et al). 2010;52:1242-5

Wound healing
Medical therapy of diabetic foot infections (Joseph and Lipsky). 2010;52:S67-71
Off-loading the diabetic foot for ulcer prevention and healing (Cavanagh and Bus). 2010;52:S37-43
Safety and efficacy of patient specific intramuscular injection of HGF plasmid gene therapy on limb perfusion and wound healing in patients with ischemic lower extremity ulceration: results of the HGF-0205 trial (Powell et al). 2010;52:1525-30
A step-wise approach for surgical management of diabetic foot infections (Fisher et al). 2010;52:S72-5
Wound care: the role of advanced wound healing technologies (Wu et al). 2010;52:S59-66

Wound infection
Débridement of the noninfected wound (Cornell et al). 2010;52:S31-6
Medical therapy of diabetic foot infections (Joseph and Lipsky). 2010;52:S67-71
A step-wise approach for surgical management of diabetic foot infections (Fisher et al). 2010;52:S72-5

Wounds and injuries
Cost-effectiveness of guidelines for insertion of inferior vena cava filters in high-risk trauma patients (Spangler et al). 2010;52:1537-45, 1537.e1-e2
Endovascular repair of thoracic aortic traumatic transections is

a safe method in patients with complicated injuries (Rahimi et al). 2010;52:891-6
Halsted-Holman vascular trauma legacy (Rich et al). 2010;52:508-11
Outcomes of selective tibial artery repair following combat-related extremity injury (Burkhardt et al). 2010;52:91-6
Traumatic thoracic aortic injury and ruptures (Tefera). 2010;52:S41-4 (Editorial)
Urinary bladder injuries during vascular surgery (Nakamura et al). 2010;52:453-5 (Case rep.)

Wounds, nonpenetrating
Aortic dilatation after endovascular repair of blunt traumatic thoracic aortic injuries (Forbes et al). 2010;52:45-8
A contemporary rural trauma center experience in blunt traumatic aortic injury (Durham et al). 2010;52:884-90
Economic evaluation of open vs endovascular repair of blunt traumatic thoracic aortic injuries (Tong et al). 2010;52:31-8
Limb salvage after delayed diagnosis for blunt traumatic infrapopliteal occlusion (Causey et al). 2010;52:734-7 (Case rep.)
The limitations of thoracic endovascular aortic repair in altering the natural history of blunt aortic injury (Lang et al). 2010;52:290-7
Regarding "Right subclavian pseudoaneurysm secondary to blunt trauma in an arteriovenous malformation" (Lee) (Letter);(Quinones-Baldrich,) (Reply). 2010;52:257-8